

## **REFERENCES**

## REFERENCES

- AACE/ACE. (1998). Position statement on the prevention, diagnosis and treatment of obesity. AACE/ACE Obesity Task Force. **Endocrine practice**, 4(5), 297-350.
- Abdellaoui, A, A.-K. (2007). C-reactive protein (CRP) as a marker in peripheral vascular disease. **Eur J Vasc Endovasc Surg**, 34(1), 18-22.
- Aekplakorn, W., Hogan, MC., Chongsuvivatwong, V., Tatsanavivat, P., Chariyalertsak, S., Boonthum, A., et al. (2007). Trends in obesity and associations with education and urban or rural residence in Thailand. **Obesity**, 15, 3113-3121.
- Albert, MA., Glynn, RJ., Buring, J. and Ridker, PM. (2004). C-reactive protein levels among women of various ethnic groups living in the United States (from the Women's Health Study). **Am J Cardiol**, 93, 1238-1242.
- Albert, W. Tsai, Aaron, R. Folsom, Wayne, D. Rosamond and Daniel, W. Jones. (2001). Ankle-brachial index and 7-year ischemic stroke incidence: The ARIC study. **Stroke**, 32, 1721-1724.
- Ali, R. Khoshdel, S. L. C., Balakrishnan, R. Nair and Alastair Gillies. (2007). Better management of cardiovascular diseases by pulse wave velocity: Combining clinical practice with clinical research using evidence-based medicine. **Clinical Medicine & Research**, 5(1), 45-52.
- Allison, MA., Hiatt, WR., Hirsch, AT., Coll, JR. and Criqui, MH. (2008). A high ankle-brachial index is associated with increased cardiovascular disease morbidity and lower quality of life. **J Am Coll ardiol**, 51(13), 1292-1298.
- American Diabetes Association. (2003). Peripheral arterial disease in people with diabetes. **Diabetes Care**, 26(12), 3333-3341.
- American Diabetes Association. (2006). Standards of medical care in diabetes-2006. **Diabetes Care**, 29(1), S4-S42.
- American Heart Association. (2002). **2002 heart and stroke statistical update**. Dallas, TX: American Heart Association.

- Andreas Pfützner, Thomas Schöndorf, Markolf Hanefeld and Thomas Forst. (2010). High-sensitivity C-reactive protein predicts cardiovascular risk in diabetic and nondiabetic patients: Effects of insulin-sensitizing treatment with pioglitazone. **J Diabetes Sci Technol**, 4(3), 706-716.
- Andrew, J. Hillman, Vitool Lohsoonthorn, Orrawadee Hanvivatvong, Wiroj Jiamjarasrangsi, Somrat Lertmaharit and Michelle, A. Williams. (2010). Association of high sensitivity C-reactive protein concentrations and metabolic syndrome among thai adults. **Asian Biomedicine**, 4(3), 385-393.
- Anthony, J.G., Hanley, K.W., Michael, P. Stern, Steven and M. Haffner. (2002). Homeostasis model assessment of insulin resistance in relation to the incidence of cardiovascular disease. **Diabetes Care**, 25, 1177-1184.
- Asia Pacific Cohort Studies Collaboration. (2003). The effects of diabetes on the risks of major cardiovascular diseases and death in the Asia Pacific region. **Diabetes Care**, 26(2), 360-366.
- Austin, MA., Hokanson, JE. and Edwards, KL. (1998). Hypertriglyceridemia as a cardiovascular risk factor. **Am J Cardiol**, 81, 7B-12B.
- Avolio, AP., Chen, S., Wang, R., Zhang, C., Li, M. and O'Rourke, MF. (1983). Effects of aging on changing arterial compliance and left ventricular load in a northern Chinese urban community. **Circulation**, 68, 50-58.
- Balkau, B. and Charles, MA. (1999). Comment on the provisional report from the WHO consultation. European group for the study of insulin resistance (EGIR). **Diabet Med**, 16, 442- 443.
- Bonora, E., Kiechl, S., Willeit, J., Oberhollenzer, F., Egger, G., Targher, G., et al. (1998). Prevalence of insulin resistance in metabolic disorders: The Bruneck Study. **Diabetes**, 47, 1643-1649.
- Bonora, E., Targher, G., Alberiche, M., Bonadonna, RC., Zenere, MB., Saggiani, F., et al. (2001). Intracellular partition of plasma glucose disposal in hypertensive and normotensive subjects with type 2 diabetes mellitus. **J Clin Endocrinol Metab**, 86, 2073-2079.

- Bonow, RO., Carabello, BA. and Kanu, C. (2006). ACC/AHA 2006 guidelines for the management of patients with valvular heart disease: A report of the American College of Cardiology/American Heart Association task force on practice guidelines (writing committee to revise the 1998 Guidelines for the management of patients with valvular heart disease): Developed in collaboration with the society of cardiovascular anesthesiologists: Endorsed by the society for cardiovascular angiography and interventions and the society of thoracic surgeons. **Circulation**, 114(5), 84-231.
- Booth, G. L., M. K. Kapral, K. Fung and J. V. Tu. (2006). Recent trends in cardiovascular complications among men and women with and without diabetes. **Diabetes Care**, 29(1), 32-37.
- Brown, MA., Davis, GK. and McHugh, L. (2001). The prevalence and clinical significance of nocturnal hypertension in pregnancy. **J Hypertens**, 19, 1437-1444.
- Camila, M. Manrique, James, L. Rosenzweig and Guillermo, E. Umpierrez. (2009). Diabetes, dyslipidemia and heart protection. **The Journal of Clinical Endocrinology and Metabolism**, 94(1), 1.
- Chambers, JC., Eda, S., Bassett, P., Karim, Y., Thompson, SG., Gallimore, JR., et al. (2001). C-Reactive protein, insulin resistance, central obesity, and coronary heart disease risk in Indian Asians from the United Kingdom compared with European whites. **Circulation**, 104(2), 145-150.
- Colin Medical Technology Corporation, Komaki, Japan. (n.d.). Non-invasive vascular screening device. **VP-1000/2000 TBI package manuscript**. Komaki, Japan: Colin Medical Technology Corporation.
- Criqui, MH. and Denenberg, JO. (1998). The generalized nature of atherosclerosis: How peripheral arterial disease may predict adverse events from coronary artery disease. **Vasc Med**, 3(3), 241-245.
- Criqui, MH., Langer, RD., Fronek, A. and Feigelson, HS. (1991). Coronary disease and stroke in patients with large-vessel peripheral arterial disease. **Drugs**, 42(5), 16-21.

- Darren Thompson, Mark, B. Pepys and Steve, P. Wood. (1999). The physiological structure of human C-reactive protein and its complex with phosphocholine. **Structure**, 7(2), 169-177.
- DeFronzo, R. A. (1988). The triumvirate: B-cell, muscle, liver: A collusion responsible for NIDDM. **Diabetes**, 37, 667-687.
- Dehnavi, RA., Roosb de, A., Rabelink, TJ., Pelt van, J., Wensink, MJ., Romijn, JA., et al. (2008). Elevated CRP levels are associated with increased carotid atherosclerosis independent of visceral obesity. **Atherosclerosis**, 200(2), 417-423.
- Du Clos, T. W. (2000). Function of C-reactive protein. **Ann Med**, 32, 274-278.
- Eknoyan, Garabed. (2008). Adolphe Quetelet (1796-1874)-the average man and indices of obesity. **Nephrol. Dial. Transplant**, 23(1), 47-51.
- Enzo, Bonora, G. F., Francesco Calcaterra, Simonetta Lombardi, Franco Marini, Luciano Zenari, Francesca Saggiani, et al. (2002). HOMA-Estimated Insulin Resistance Is an Independent Predictor of Cardiovascular Disease in Type 2 Diabetic Subjects. **Diabetes Care**, 25, 1135-1141.
- Enzo Bonora, S. K., Johann Willeit, Friedrich Oberhollenzer, Georg Egger, James, B. Meigs, Riccardo C. Bonadonna, et al. (2007). Insulin resistance as estimated by homeostasis model assessment predicts incident symptomatic cardiovascular disease in caucasian subjects from the general population. **Diabetes care**, 30, 318-324.
- Ethiraj Dhanaraj, Anil Bhansali, Shallu Jaggi, Pinaki Dutta, Shikha Jain, Pramila Tiwari, et al. (2009). Predictors of metabolic syndrome in Asian north Indians with newly detected type 2 diabetes. **Indian J Med Res**, 129, 506-514.
- Festa, A., D'Agostino, R. Jr, Howard, G., Mykkanen, L., Tracy, RP. and Haffner, SM. (2000). Chronic subclinical inflammation as part of the insulin resistance syndrome: the Insulin Resistance Atherosclerosis Study (IRAS). **Circulation**, 102(1), 42-47.
- Finn, AV., Nakano, M., Narula, J., Kolodgie, FD. and Virmani, R. (2010). Concept of vulnerable/unstable plaque. **Arterioscler. Thromb. Vasc. Biol**, 30(7), 1282-1292.

- Florez, H., Castillo-Florez, S., Mendez, A., Casanova-Romero, P., Larreal-Urdaneta, C., Lee, D., et al. (2006). C-reactive protein is elevated in obese patients with the metabolic syndrome. **Diabetes Res Clin Pract**, 71(1), 92-100.
- Gabay, C. and Kushner, I. (1999). Mechanisms of disease: Acute-phase proteins and other systemic responses to inflammation. **N Engl J Med**, 340, 448-454.
- Gary, L., Myers, N. R., Russell, P. Tracy, William, L. Roberts, R. Wayne Alexander, Luigi, M. Biasucci, et al. (2004). CDC/AHA workshop on markers of inflammation and cardiovascular disease: Application to clinical and public health practice: Report from the laboratory science discussion group. **Circulation**, 110, e545-e549.
- Giacomo Ruotolo, Barbara, V. Howard and David, C. Robbins. (2003). Dyslipdemia of obesity. Retrieved April 10, 2013, from <http://www.endotext.org/obesity/obesity10/obesityframe10.htm>
- Göran, K. Hansson (2005). Mechanisms of disease inflammation, atherosclerosis and coronary artery disease. **The new england journal of medicine (review article)**, 2005, 1685-1695.
- Guixiang Zhao, Earl S Ford, Chaoyang Li, James Tsai, Satvinder Dhingra and Lina S Balluz. (2011). Waist circumference, abdominal obesity, and depression among overweight and obese U.S. adults: national health and nutrition examination survey 2005-2006. **BMC Psychiatry**, 11(130), 1-9.
- Haffner, S. M., S. Lehto, T. Ronnemaa, K. Pyorala and M. Laakso. (1998). Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. **New England Journal of Medicine**, 339(4), 229-234.
- Hak, AE., Stehouwer, CD., Bots, ML., Polderman, KH., Schalkwijk, CG., Westendorp, IC., et al. (1999). Associations of C-reactive protein with measures of obesity, insulin resistance, and ubclinical atherosclerosis in healthy, middle-aged women. **Arterioscler Thromb Vasc Biol**, 19(8), 1986-1991.

- He, J. D. Gu, J. Chen, X. Wu, T. N. Kelly, J.-F. Huang, J.-C. Chen, C., et al. (2009). Premature deaths attributable to blood pressure in China: A prospective cohort study. *Lancet*, 374(9703), 1765-1772.
- Helaine, E., Resnick, R. S. L., Mary McGrae McDermott, Richard, B., Devereux, Kristina L., Jones, Richard R. Fabsitz, et al. (2004). Relationship of high and low ankle brachial index to all-cause and cardiovascular disease mortality: The strong heart study. *Circulation*, 109, 733-739.
- Hirofumi Tomiyama, Akira Yamashina and Tomio Arai. (2003). Influences of age and gender on results of noninvasive brachial-ankle pulse wave velocity measurement-a survey of 12517 subjects. *Atherosclerosis*, 166, 209-303.
- Hirsch, AT., Criqui, MH., Treat-Jacobson, D., Regensteiner, JG., Creager, MA., Olin, JW., et al. (2001). Peripheral arterial disease detection, awareness, and treatment in primary care. *JAMA*, 286(11), 1317-1324.
- Hosseini Fakhrazadeh and Ozra Tabatabaei Malazy. (2012). Dyslipidemia and cardiovascular disease. **Dyslipidemia - From Prevention to Treatment**, 2012, 303-320.
- Hotamisligil, G., Shargill, NS. and Spiegelman, BM. (1993). Adipose expression of tumor necrosis factor-  $\alpha$ : direct role in obesity-linked insulin resistant. *Science*, 259(5091), 87-91.
- IDF. (2013). **International diabetes federation atlas, sixth edition**. Retrieved September 12, 2013, from [http://www.idf.org/sites/default/files/EN\\_6E\\_Atlas\\_Full\\_0.pdf](http://www.idf.org/sites/default/files/EN_6E_Atlas_Full_0.pdf).
- IDF. (2005). **The IDF consensus worldwide definition of the metabolic syndrome**. N.P.: Promoting diabetes care, prevention and a cure worldwide.
- James, J. Maciejko. (2004). Atherosclerosis risk factors. *AACC Press*, 2004, 91-93.
- Joan, M. Hengst. (2003). The role of C-Reactive protein in the evaluation and management of infants with suspected sepsis. *Adv Neonatal Care*, 3(1), 1-8.
- John, B. Dixon and Paul, E. O'Brien. (2002). Lipid profile in the severely obese: Changes with weight loss after lap-band surgery. *Obesity Research*, 10(9), 19-25.

- Judith, A., Berliner, M. N., Alan, M., Fogelman, Joy S., Frankm Linda, L., Demer Peter, A., et al. (n.d.). **Atherosclerosis: Basic mechanisms oxidation, inflammation and Genetics**. Retrieved October 14, 2013, from <http://circ.ahajournals.org/content/91/9/2488.short>.
- Kagan, A., Gordon, T., Kannel, WB. and Dawber, TR. (1959). American heart association. In **Proceedings of the Council for High Blood Pressure Research** (p.81). New York: n.p.
- Kanokwan Srisupornkornkool, Kanjana Jittiporn and Wanvisa Boonlert. (2008). The occult prevalence of metabolic syndrome in Naresuan university staff: Comparison of serum low density lipoprotein levels in individuals of metabolic syndrome, at Risk and not at risk of metabolic syndrome. In **Poster presentation at the 4<sup>th</sup> Naresuan Research Conference**. Phitsanulok: Naresuan University.
- Kanokwan Srisupornkornkool, Sompiya Somthavil, Nidsarakul Wisetsut, Suthasinee Suebson and Suranchana Tongnak. (2007). Assessment of metabolic syndrome risk from central obesity and blood pressure among Naresuan university Personnel. **J. Naresuan University**, 15(2), 89-98.
- Kengne, A. P., A. Patel, F. Barzi, K. Jamrozik, T. H. Lam, H. Ueshima, et al. (2007). Systolic blood pressure, diabetes and the risk of cardiovascular diseases in the Asia-Pacific region. **Journal of Hypertension**, 25(6), 1205-1213.
- Kushner I. (1990). C-reactive protein and the acute phase response. **Hosp Pract**, 25, 13-28.
- Lawes Carlene, M.M., Stephen Vander Hoorn, Anthony Rodgers and International Society of Hypertension. (2008). Global burden of blood-pressure-related disease, 2001. **Lancet**, 371(9623), 1513-1518.
- Lawrence de Koning, Anwar, T. Merchant, Janice Pogue and Sonia S. Anand. (2007). Waist circumference and waist-to-hip ratio as predictors of cardiovascular events: meta-regression analysis of prospective studies. **European Heart Journal**, 28, 850-856.
- Lehmann, ED., Gosling, RG. and Sonksen, PH. (1992). Arterial wall compliance in diabetes. **Diabet Med**, 9, 114-119.

- Leng, GC., Lee, AJ., Fowkes, FG., Whiteman, M., Dunbar, J., Housley, E., et al. (1996). Incidence, natural history and cardiovascular events in symptomatic and asymptomatic peripheral arterial disease in the general population. **Int J Epidemiol**, 25(6), 1172-1181.
- Li Xu, Katashima Mitsuhiro, Yasumasu Takeshi and Li Ke Ji. (2012). Visceral fat area, waist circumference and metabolic risk factors in abdominally obese chinese Adults. **Biomed Environ Sci**, 25(2), 141-148.
- Libby, P. (2006). Inflammation and cardiovascular disease mechanisms. **The American Journal of Clinical Nutrition**, 83, 456S– 460S.
- Linda, J. Vorvick. (2011). **Developmental process of atherosclerosis**. Retrieved November 28, 2013, from <http://www.nlm.nih.gov/medlineplus/ency/imagepages/18020.htm>
- Liz Smith. (2005). New AHA recommendations for blood pressure measurement. **Am Fam Physician**, 72(7), 1391-1398.
- London, GM., Marchais, SJ. and Safar, ME. (1990). Aortic and large artery compliance in end-stage renal failure. **Kidney Int**, 37, 137-142.
- M. K. Poulsen, J. E. Henriksen and J. Dahl. (2010). Left ventricular diastolic function in type 2 diabetes mellitus prevalence and association with myocardial and vascular disease. **Circulation: Cardiovascular Imaging**, 3(1), 24-31.
- Mandel William, J. (1995). **Cardiac arrhythmias: Their mechanisms, diagnosis and management** (3<sup>rd</sup> ed.). N.P.: Lippincott Williams and Wilkins.
- Mayosi, B. M., A. J. Flisher, U. G. Lalloo, F. Sitas, S. M. Tollman and D. Bradshaw. (2009). The burden of non-communicable diseases in South Africa. **Lancet**, 374(9693), 934-947.
- Mbewu, A. and J. Mbanya. (2006). Cardiovascular disease. In D. Jamison, R. G. Feachem, M. W. Makgoba, E. R. Bos, F. K. Baingana, K. J. Hofman, et al. (Eds.), **Disease and mortality in Sub-Saharan Africa** (pp.305-328). Washington, DC: World Bank Group.
- Mora, S., Musunuru, K. and Blumenthal, RS. (2009). The clinical utility of high-sensitivity C-reactive protein in cardiovascular disease and the potential implication of JUPITER on current practice guidelines. **Clin Chem**, 55(2), 219-228.

- Mosca, L., Banka, CL. and Benjamin, EJ. (2007). Evidence-based guidelines for cardiovascular disease prevention in women: 2007 update. **Circulation**, 115, 1481-1501.
- Muna, W. F. T. (1993). Cardiovascular disorders in Africa. **World Health Statistics Quarterly**, 46(2), 125-133.
- Nader Rifai, Russell P. Tracy and Paul M. Ridker. (1999). Clinical efficacy of an automated high-sensitivity c-reactive protein assay. **Clinical Chemistry**, 45(12), 2136-2141.
- Navapun Charuruks, Bupayao Laohajinda , Suthee Rujiwanitgun and Mathuros Chaiworaporn. (2005). Reference value for C-reactive protein and its distribution in Thai adults. **Circulation Journal**, 69, 339-344.
- NCEP and ATP III. (2001). Expert panel on detection, evaluation and treatment of high blood cholesterol in adults. Executive summary of the third report of the national cholesterol education program (NCEP). Expert panel on detection, evaluation and treatment of high blood cholesterol in adults (Adult Treatment Panel III). **JAMA**, 285, 2486-2497.
- NCEP and ATP III. (2004). Third report of the national cholesterol education program (NCEP). Expert panel on detection, evaluation and treatment of high blood cholesterol in adults (Adult Treatment Panel III). Final report, national institutes of health, national heart, lung and blood institute. **NIH Publication**, 2004, 1-7.
- Newman, AB. (2000). Peripheral arterial disease: Insights from population studies of older adults. **J Am Geriatr Soc**, 48(9), 1157-1162.
- Nichols, WW. and O'Rourke, MF. (1990). Properties of the arterial wall. In **McDonald's Blood Flow in Arteries. Theoretical, Experimental and Clinical Principles** (3<sup>rd</sup> ed., pp.77-114). London: Edward Arnold.
- Päivi Lempiäinen, L. M., Kalevi Pyörälä, Markku Laakso and Johanna Kuusisto. (1999). Insulin resistance syndrome predicts coronary heart disease events in elderly nondiabetic men. **Circulation**, 100, 123-128.
- Paul M Ridker. (2001). High-sensitivity C-reactive protein: Potential adjunct for global risk assessment in the primary prevention of cardiovascular disease. **Circulation**, 103, 1813-1818.

- Paul, M. Ridker. (2003). Clinical application of C-reactive protein for cardiovascular detection and prevention. **Circulation**, 107, 363-369.
- Paul, S. Jellinger, Donald, A. Smith, Adi, E. Mehta, Om Ganda, Yehuda Handelsman, Helena, W. Rodbard, et al. (2012). American association of clinical endocrinologists' guidelines for management of dyslipidemia and prevention of atherosclerosis. **Endocrine practice**, 18(1), 1-78.
- Pekka Jousilahti, Erkki Vartiainen, Jaakko Tuomilehto and Pekka Puska. (1999). Sex, age, cardiovascular risk factors and coronary heart disease a prospective follow-up study of 14 786 middle-aged men and women in finland. **Circulation**, 99, 1165-1172.
- Pepys, M. (1981). C-reactive protein fifty years. **Lancet**, 1, 653-657.
- Pepys, MB., H. G. (2003). C-reactive protein: A critical update. **J. Clin. Invest**, 111(12), 1805-1812.
- Porapakham, Y. and Bunyaratpundu, P. (2006). **Health system research institute and bureau of policy and strategy**. Bangkok: Ministry of Public Health.
- Pouliot, M.C., Després, J.P., Lemieux, S., Moorjani, S., Bouchard, C., Tremblay A., et al. (1994). Waist circumference and abdominal sagittal diameter: Best simple anthropometric indexes of abdominal visceral adipose tissue accumulation and related cardiovascular risk in men and women. **Am J Cardiol**, 73, 460-468.
- PubMed Health, A.D.A.M., I. (2012). **Alcoholism and alcohol abuse**. N.P.: n.p.
- Rahim, MA., Azad Khan, AK., Sayeed, MA., Akhtar, B., Nahar, Q., Ali, SMK., et al. (2007). Metabolic syndrome in rural Bangladesh: Comparison of newly proposed IDF, modified ATP III and WHO criteria and their agreements. **Diabetes Metab Res Rev**, 1(4), 251-257.
- Razay George. (2006). **Obesity, abdominal obesity and alzheimer's disease**. N.P.: Dementia and Genetic Cognitive Disorders.
- Richard, C. Becker. (2005). Heart attack and stroke prevention in women. **Circulation**, 112, e273-e275.

- Rinkoo Dalan, Michelle Jong, Siew-Pang Chan, Robert Hawkins, Robin Choo, Brenda Lim, et al. (2010). High-sensitivity C-reactive protein concentrations among patients with and without diabetes in a multiethnic population of Singapore: CREDENCE Study. *Diabetes, metabolic syndrome and obesity: Targets and therapy*. Dovepress, 2010, 187-195.
- Robert, WL., Moulton, L., Law, TC., Farrow, G., Cooper-Anderson, M., Savory, J., et al. (2001). Evaluation of nine automated high-sensitivity C-Reactive protein methods: Implications for clinical and epidemiological applications (Part 2). *Clin Chem*, 47(3), 3418-3425.
- Samuel Klein, David, B. Allison, Steven, B. Heymsfield, David, E. Kelley, Rudolph, L. Leibel, Cathy Nonas, et al. (2007). Waist circumference and cardiometabolic risk: A consensus statement from shaping America's health: Association for weight management and obesity prevention; NAASO, the obesity society, the American society for nutrition and the American diabetes association1– 4. *Am J Clin Nutr*, 85, 1197-1202.
- Sarah Wild, Gojka Roglic, Anders Green, Richard Sicree and Hilary King. (2004). Global prevalence of diabetes estimates for the year 2000 and projections for 2030. *Diabetes Care*, 27, 1047-1053.
- Sarwar, N., Danesh, J., Eiriksdottir, G., Sigurdsson, G., Wareham, N., Bingham, S., et al. (2007). Triglycerides and the risk of coronary heart disease: 10,158 incident cases among 262,525 participants in 29 Western prospective studies. *Circulation*, 115, 450-458.
- Scott, M. Grundy, James, I. Cleeman, Stephen, R. Daniels, Karen, A. Donato, Robert, H. Eckel, Barry, A. Franklin, et al. (2005). Diagnosis and management of the metabolic syndrome an American heart association/national heart, lung and blood institute scientific statement. *Circulation*, 112, 2735-2752.
- Sims, N.R. and Muyderman, H. (2009). Mitochondria, oxidative metabolism and cell death in stroke. *Biochimica et Biophysica Acta*, 1802(1), 80-91.
- Sirtori, Cesare R. (2006). HDL and the progression of atherosclerosis: new insights. *European Heart Journal Supplements*, 19, 47-53.

- South African Department of Health and Medical Research Council. (2007).  
**South Africa demographic and health survey 2003**. Pretoria, South Africa:  
 South African Department of Health.
- Stamler, J., R. Stamler and J. D. Neaton. (1993). Blood pressure, systolic and diastolic,  
 and cardiovascular risks: US population data. **Archives of Internal Medicine**,  
 153(5), 598-615.
- Stanley, S. and Franklin, N. D. W. (2013). Hypertension and cardiovascular disease:  
 Contributions of the framingham heart study. **Global Heart**, 8(1), 49-57.
- Thom, T., Haase, N. and Rosamond, W. (2006). Heart disease and stroke statistics-  
 2006 update: A report from the American heart association statistics committee  
 and stroke statistics subcommittee. **Circulation**, 113, e85-e151.
- Thomas, A., Pearson George, A., Mensah, R., Wayne Alexander, Jeffrey, L.,  
 Anderson, Richard, O.Cannon III, et al. (2003). Markers of inflammation and  
 cardiovascular disease: Application to clinical and public health practice:  
 A Statement for healthcare professionals from the centers for disease control  
 and prevention and the American heart association. **Circulation**, 107,  
 499-511.
- Thomas, A. Pearson, George, A. Mensah, Yuling Hong and Sidney, C. Smith. (2004).  
 CDC/AHA Workshop on Markers of Inflammation and Cardiovascular  
 Disease: Application to Clinical and Public Health Practice: Report from the  
 Laboratory Science Discussion Group. **Circulation**, 110(25), e545-549.
- Thomas, R. J., P. J. Palumbo, L. J. Melton Iii, V. L. Roger, J. Ransom, P. C. O'Brien,  
 et al. (2003). Trends in the mortality burden associated with diabetes mellitus:  
 A population-based study in Rochester, Minn, 1970-1994. **Archives of  
 Internal Medicine**, 163(4), 445-451.
- Thygesen, K., Alpert, J.S. and White, H.D. (2007). Universal definition of myocardial  
 infarction. **Eur. Heart J**, 28(20), 2525-2538.
- Tomiyama, H., Yamashina, A., Arai, T., Hirose, K., Koji, Y., Chikamori, T., et al.  
 (2003). Influences of age and gender on results of noninvasive brachial-ankle  
 pulse wave velocity measurements-a survey 12,517 subjects. **Atherosclerosis**,  
 166(2), 303-309.

- Toth, Peter. (2005). The good cholesterol high-density lipoprotein. **Circulation**, 111(5), e89-e91.
- Trayhurn, P. and Wood, IS. (2005). Signalling role of adipose tissue: Adipokines and inflammation in obesity. **Biochemical Society Transaction**, 33(5), 1078-1081.
- Vague, J. (1947). La differenciation sexuelle, facteur determinant des formes de l'obesite. **La Presse Medicale**, 30, 339-340.
- Valentín Fuster and Bridget B. Kelly. (2010). **Promoting cardiovascular health in the developing world a critical challenge to achieve global health.** Washington, D.C.: The National Academies Press.
- Visser, M., Bouter, L.M., McQuillan, G.M., Wener, M.H. and Harris T.B. (1999). Elevated C-reactive protein levels in overweight and obese adults. **JAMA**, 282(22), 2131-2135.
- W., Aekplakorn and L., Mo-suwan. (2009). Prevalence of obesity in Thailand. **Obesity reviews**, 10, 589-592.
- Wada, T., Kodaira, K. and Fujishiro, K. (1994). Correlation of ultrasound-measured common carotid artery stiffness with pathological findings. **Arterioscler Thromb Vasc Biol**, 14, 479-482.
- Wanvisa Boonlert, Gerald, J. Kost, Aranya Jiraviriyakul, Surapon Tangvarasittichai. (2006). Point-of-care testing on a mobile medical unit in Northern Thailand: Screening for hyperglycemia, hyperlipidemia and thalassemia trait. **Point of care: The journal of near-patient testing & technology**, 5(4), 164 - 167.
- WHO. (1999). **Definition, diagnosis and classification of diabetes mellitus and its complications. report of a WHO consultation world health organization department of noncommunicable disease surveillance geneva.** Geneva: World Health Organization.
- WHO. (2000). **Obesity: Preventing and managing the global epidemic: WHO technical report series, No. 894.** Geneva: World Health Organization.
- WHO. (2010). Global status report on noncommunicable diseases 2010. In **WHO library cataloguing-in-publication data** (pp.1-31). Geneva: World Health Organization.
- WHO. (2011). **Global status report on noncommunicable disaeses 2010.** Geneva: World Health Organization.

- WHO. (2011). **Global atlas on cardiovascular disease prevention and control**. Geneva: World Health Organization.
- WHO. (2013). **Cardiovascular diseases (CVDs)**. Retrieved March 20, 2013, from <http://www.who.int/mediacentre/factsheets/fs317/en/index.html>
- WHO/IOTF/IASO. (2000). **The Asia-Pacific perspective: Redefining obesity and its treatment**. Melbourne, Australia: World Health Organization, International Obesity Task Force, International Association for the Study of Obesity.
- Wikipedia, t. f. e. "Low density lipoprotein."** (2008). Retrieved March 7, 2008, from [http://en.wikipedia.org/wiki/Low\\_density\\_lipoprotein](http://en.wikipedia.org/wiki/Low_density_lipoprotein)
- Yamashina, A., Tomiyama, H., Takeda, K., Tsuda, H., Arai, T., Hirose, K., et al. (2002). Validity, reproducibility and clinical significance of noninvasive brachial-ankle pulse wave velocity measurement. **Hypertens Res**, 25(3), 359-364.
- Yang, G., L. Kong, W. Zhao, X. Wan, Y. Zhai, L. C. Chen, et al. (2008). Emergence of chronic non-communicable diseases in China. **Lancet**, 372(9650), 1697-1705.
- Yasojima, K. S. C., McGeer, EG. and McGeer, PL. (2001). Generation of C-reactive protein and complement components in atherosclerotic plaques. **Am J Pathol**, 158, 1039-1051.
- Yi-Hao, Yu. and Henry, N. Ginsberg. (2005). Adipocyte signaling and lipid homeostasis sequelae of insulin-resistant adipose tissue. **Circulation Research**, 96(10), 1042-1052.
- Yuanxi, X.U., Jue, L.I. and Yingyi, L.U.O. (2007). The association between ankle-brachial index and cardiovascular or all-cause mortality in metabolic syndrome of elderly Chinese. **Hypertension research; The Japanese Society of Hypertension**, 30(7), 613-619.
- Yusuf, S., Hawken, S., Ounpuu, S., Dans, T., Avezum, A., Lanas, F., et al. (2004). Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): Case-control study. **Lancet**, 364(9438), 937-952.

Zwaka, TP., Hombach, V. and Torzewski, J. (2001). C-reactive protein-mediated low density lipoprotein uptake by macrophages: Implications for atherosclerosis. **Circulation**, 103, 1194-1197.

## **BIOGRAPHY**

## BIOGRAPHY

**Name – Surname** Patawee Ngamlert  
**Date of Birth** September 26, 1974  
**Address** 98/100 Moo 1, Kookot sub-district, Lam look ka district, Patumthani province, Thailand 12130  
**Work Place** Police General Hospital  
**Position** Medical Technologist (Professional)  
**Work Experiences**  
2002 Police Hospital, Bangkok  
1997 Bumrungrad Hospital, Bangkok  
**Education Background**  
1993 B.Sc. (Medical Technology), Mahidol University, Thailand

### Publication

- Treebuphachatsakul, WB. and Ngamlert, P. (2013). High sensitivity C–Reactive protein showing independent relationship with peripheral vascular atherosclerosis and waist circumference in healthy adults. *IJPBS*, 4(2), 918-926.
- Ngamlert, P., Chittamma, A. and Boonlert, W. (2010). Serum high sensitivity C-Reactive protein levels between central obesity and healthy normal waist circumference adults. In **Proceeding in the 11<sup>th</sup> graduate research conference** (pp.844-849). Khon Kaen: Khon Kaen University.