

REFERENCES

- Andreae, M.O. and Merlet, P. (2001), Emissions of trace gases and aerosols from biomass burning. *Global Biogeochemistry Cycles*, **15**, 4, pp. 955-966.
- Andreae, M.O. and Crutzen, P. J. (2001), Atmospheric aerosols: biogeochemical sources and role in atmospheric chemistry, *Science*, **276**, 5315, pp. 1052-1058.
- Akaakara, S., Viriya, K. and Tongtan, T. (2003). Fire behaviors in dry dipterocarp forest at Huay Kha Khaeng Wildlife Sanctuary, Research Report, Forest Fire Research Center, Uthai Thani' Forest Fire Control Office, National Park, Wildlife and Plant Conservation Department, Bangkok, Thailand, 43 pages. (in Thai)
- ASTM international (2007), Standard Test Method for Ash and Biomass, Designation: E 1755-01, United States, 3 pages.
- Baker, P. J. and Bunyavejchewin, S. (2009), Fire behavior and fire effects across the forest landscape of continental Southeast Asia, *Tropical Fire Ecology*, pp. 311-321.
- Bolin, B. (1970), The carbon cycle, *Scientific American*, **223**, pp. 125-132.
- Boonrodklab, C. and Teejuntuk, S. (2008), 'Structure and aboveground biomass of forest community in Doi Inthanon National Park, Chiangmai Province' Kasetsart University Annual Conference, Bangkok, Thailand. (in Thai)
- Brodowski, S., Amelung, W., Haumaier, L., Abetz, C., and Zech, W. (2005), Morphological and chemical properties of black carbon in physical soil fractions as revealed by scanning electron microscopy and energy-dispersive X-ray spectroscopy. *Geoderma Journal*, **128**, pp. 116-129.
- Byram, G. M. (1959), Combustion of forest fuels. In 'Forest Fire: Control and Use'. (Ed. KP Davis) 61-89. (McGraw – Hill : New York)
- Cachier, H., Liousse, C., Buat-Mernard, P. and Gaudichet, A. (1995), Particle content of savanna fire emissions, *Journal of Atmospheric Chemistry*, **22**, pp. 123-148.

- Cao, G., Zhang, X., Gong, S. and Zheng, F. (2008), Investigation on emission factors of particulate matter and gaseous pollutants from crop residue burning, *Journal of environmental sciences (China)*, **20**, pp. 50-55.
- Cao, J. J. et al., (2003), Characteristics of carbonaceous aerosol in Pearl River Delta Region, China, during 2001 winter period, *Atmospheric Environment*, **37**, pp. 1451-1460.
- Chaiyo, U., Garivait, S. and Wilairat, D. (2011). Trace Elements and Carbon Contents in Particulate Emissions from Tropical Deciduous Forest Fires in Chiangmai, Thailand, paper presented in the 2nd International Conference on Environmental Science and Technology (ICEST 2011), pp. 213-217, Singapore.
- Chen, X. G., Zhang, X. Q., Zhang, Y. P., Booth, T. and He, X. H. (2009), Changes of carbon stocks in bamboo stands in China during 100 years, *Forest Ecology and Management*, **258**, pp. 1489-1496.
- Chhabra, A. and Dadhwal, V. K. (2004), Assessment of major pools and fluxes of carbon in Indian forests, *Climate Change*, **64**, 341-360.
- Chow, J.C. and Watson, J.G. (1999), In *Elemental Analysis of Airborne Particles*, Landsberger, S. and Creatchman, M. (Eds.), Gordon and Breach Science, Amsterdam, **1**, p. 97.
- Cohen, D. (2004), Form and distribution of trace elements in biomass for power generation, research report 48. QCAT Technology Transfer Centre, Technology court Pullenvale Qld 4069 Australia.
- Cruz, C. N. and Pandiz, S. N. (1997), A study of the ability of pure secondary organic aerosol to act as cloud condensation nuclei, *Atmospheric Environment*, **31**, pp. 2205-2214.
- Crutzen, V. S. and Andreae, M. O. (1990), Biomass burning in the tropics: Impact on atmospheric chemistry and biogeochemical cycles, *Science*, **250**, pp. 1669-1678.

- Forest Fire Control Office, Forest fire control yearly report (2005), National Park Wildlife and Plant Conservation Department, Bangkok, Thailand (in Thai).
- Friedl A., Padouvas E., Rotter H., and Varmauza, K. (2005), Prediction of heating values of biomass fuel from elemental composition, *Analytica Chimica Acta*, **544**, 191-198.
- Glaser, B., Balashov, E., Haumaier, L., Guggenberger, G., Zech, W. (2000), Black carbon in density fractions of anthropogenic soils of the Brazilian Amazon region, *Organic Geochemistry*, **31**, pp. 669-678.
- Glaser, B., Guggenberger, G., Haumaier, L., Zech, W. (2001), Persistence of soil organic matter in archaeological soils (Terra Preta) of the Brazilian Amazon region. In: Rees, R. M., Ball, B.C., Campbell, C.D., Watson, C.A. (Eds.), Sustainable management of soil organic matter. CAB international, Wallingford, U.K. pp. 190-194.
- Hadley, O. L., Corrigan, C. E. and Kirchstetter, T. W. (2008), Modified thermal-optical analysis using spectro absorption selectivity to distinguish black carbon from pyrolyzed organic carbon. *Environ, Science Technology*, **42**, pp. 8459-8464.
- Hairiah, K., Sitonpul, S. M., Noordwijk, M. V. and Palm, C. (2001), Method for sampling carbon stocks above and below ground, ASB lecture note 4B. International Centre for Research in Agroforestry, Bogor, Indonesia. 32 pages.
- Himmapan, W. (2004), Behavior, effect and smoke composition of burning fire in dry dipterocarp forest at Huay Kha Khaeng Wildlife Sanctuary, Uthai Thani Province. Ph.D. Thesis, Kasetsart University, Bangkok, Thailand.
- Issaree, M. (1982), 'The primary productivity of plant communities in abandoned farm at the environmental research station, Sakaerat, Pak Thong Chai district, Nakhon Ratchasima' Master thesis, Graduated School, Kasetsart University, Bangkok, Thailand. (in Thai)
- Jacobson, M. Z. (2001), Strong radiative heating due to the mixing state of black carbon in atmospheric aerosols, *Nature*, **409**, pp. 695-697.

- Jampanin, S., and Gajasen, N. (2004), Assessment of carbon sequestration, litter production and litter decomposition in Kaeng Krachan National Park, Thailand. "The proceeding of Conference of Climate Change in Forest: Forest and Climate Change, organized by the National Park Wildlife and Plant Conservation Department, pp 1-15. (in Thai)
- Kaeskrom, P., Kaewkla, N., Thummikakpong, S. and Punsang, S. (2011), Evaluation of carbon storage in soil and plant biomass of primary and secondary mixed deciduous forests in the lower northern part of Thailand, *African Journal of Environmental Science and Technology*, **5**, 8-14.
- Kirchstetter T. W., Novakov, T., and Hobbs, P. (2004), Evidence that the spectral dependence of light absorption by aerosols is affected by organic carbon, *Journal of Geophysics and Research*, **109**, D21.
- Kruasuwun, J., Chotamornsuk, C., Ratjiranukul, P. and Wiralwechayan, A. (2008), Project of weather analysis and air monitoring. (in Thai).
- Larson, S. M. and Cass, G. R. (1989), Characteristics of summer midday low visibility events in the Los Angeles area, *Environmental Science and Technology*, **23**, pp. 281-288.
- Larson, S. M. and Cass, G. R. (2005), Characteristics of summer midday low visibility events in the Los Angeles area. *Environmental Science and Technology*, **23**, 397-407.
- Lee, S. C., Y. Cheng, K. F. Ho, J. J. Cao, P. K. K. Louie, J. C. Chow, and J. G. Watson (2005), PM_{1.0} and PM_{2.5} characteristics in the roadside environment of Hong Kong, *Aerosol Science and Technology*, **40**, pp. 157-165.
- Li, J., Huang, C., Zhang, G. (2006), Density, storage and spatial distribution of carbon in *Pleioblastus amarus* forest returned from farmland, *Journal of Zhejiang Forestry Science and Technology*, **26**, 4, pp. 2-5.
- Lin, Y., Lin P., and Wen, W. (1998), Studies on dynamics of carbon and nitrogen elements in *Dendrocalamopsis oldhami* forest, *Journal of Bamboo Research*, **17**, 4, pp. 25-30.

- McGroddy, M. E., Daufresne, T. and Hedin, L. O. (2004), Scaling of C:N:P stoichiometry in forests worldwide: Implications of terrestrial Redfield-type ratios. *Ecology*, **85**, pp. 2390-2401.
- Nelson, R. M. (1992), An evaluation of carbon mass technique for estimating emission factors and fuel consumption in forest fires. *U.S. Department of Agriculture, Forest Service, Southeastern Forest Experimental Station, Asheville, NC, Research Paper SE-231*.
- Neto, T.G.S. et al, (2009), Biomass consumption and CO₂, CO and main hydrocarbon gas emissions in an Amazonian forest clearing fire, *Atmospheric Environment*, **43**, pp. 438-446.
- Novokov, T. and Penner, J. E. (1993), Large contribution of organic aerosols to cloud-condensation nuclei concentrations, *Nature*, 365, pp. 323-365.
- Nuanurai. N. (2005), 'Comparison of leaf area index, above-ground biomass and carbon sequestration of forest ecosystems by forest inventory and remote sensing at Kaeng Krachan National Park, Thailand' Master Thesis, Graduated school, Chulalongkorn University, Bangkok, Thailand, 195 pages. (in Thai)
- Ogawa, H., Yoda, K., Ogino, K. and Kira, T. (1965), Comparative ecological studies on three main types of forest vegetation in Thailand, II, Plant Biomass, *Nature and Life in Southeast Asia*, **4**, pp. 49-80.
- Palheiro, P. M., Paulo, F., Miguel, G. Cruz (2006), A fire behavior-based fire danger classification for maritime pine stands: Comparison of two approaches, *Forest Ecology and Management* **234s**, s54.
- Penner, J. E., H. Eddleman, and T. Novokov. (1993), Towards the development of a global inventory for black carbon emissions, Part A. General Topics, *Atmospheric Environment*, **27**, pp. 1277.
- Permadi, D. A. and Kim Oanh, N. T. (2013), Assessment of biomass open burning emissions in Indonesia and potential climate forcing impact, *Atmospheric Environment*, **27**, pp. 250-258.

Pollutant Control Department, Misnistry of Natural Resources and Environment, Thailand, Air Quality and Noise, 2005.

Ramanathan, V. and Carmichael, G. (2008), Global and regional climate changes due to black carbon, *Nature Geoscience*, **1**, pp. 221-227.

Radke, L. F., Lyons, J. H., Hobbs, P.V., Hegg, D. A., Sandberg, D. V., and Ward D. E. (1990), Airborne monitoring and smoke characterization of prescribed fires on forest lands in western Washington and Oregon. Gen. Tech. Rep. PNW-GTR-251, U.S. Dept. of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, OR, 81pages.

Reid, J.S., Koppman, R., Eck, R.F., and Eluterio, D.P. (2005), A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmospheric Chemistry and Physical*, **5**, pp. 799-825.

Ryan, K. C. (2002), Dynamic interactions between forest structure and fire behavior in Boreal ecosystem, *Silva Fennica*, **36**, 1, pp. 13-39.

Samran, S. (2005). Effect of forest fire on change above ground biomass in the Maeklong mixed deciduous forest, Kanchanaburi province, Thailand, paper presented in the, *Conference on Climate Change in Forest: The Potential of Forest in Support of the Kyoto Protocol*, organized by the National Park Wildlife and Plant Conservation Department, Bangkok, Thailand, pp. 351-362 (in Thai).

Schmidl, C., Bauer, H., Datter, A., Hitzenberger, R., Weissenboek, G., Marr, J. L. and Puxbaum, H. (2008), Chemical characterization of particle emissions from burning leaves, *Atmospheric Environment*, **34**, pp. 1641-1653.

Sompoh, B. and Kaitpaneet, S. (2000). Fuel complex in deciduous forest at Huai Kha Khaeng Wildlife Sanctuary, paper presented in the, *38th Annual Meeting Conference*, Kasetsart University, pp 106-115. (in Thai)

Sompoh, B. (1998), Fuel complex in dry dipterocarp and mixed deciduous forests at Huai Kha Khaeng Wildlife Sanctuary, Changwat Uthai Thani, Master of Sciences

(Forestry), Major Field Silviculture, Department of Silviculture, Kasetsart University, Bangkok, Thailand, 83 pages.

Streets, D.G., et al. (2001), Black carbon emissions in China, *Atmospheric Environment*, **35**, 25, pp. 4281-4296.

Suwannapinunt, W. (1983), A study on the biomass of *Thyrsostachys siamensis* GAMBLE forest at Hin-Lap, Kanchanaburi, *Journal of Bamboo Research*, pp. 82-101.

Suthivanit, S. (1989), Effects of fire frequency on vegetation in dry dipterocarp forest at Sakarat, Changwat Nakorn Ratchasima, Master thesis, Forest Resource Administration, Kasetsart University, Bangkok, Thailand, 170 pages (in Thai).

Sudthichart, K. (1996), Impact of fire on soil and plant at Phu Kradueng National Park: Final report. Department of Silviculture, Faculty of Forestry, Kasetsart University, Bangkok, Thailand.

Terakunpisut, J., Gajaseni, N. and Ruankawe, N. (2007), Carbon sequestration potential in aboveground biomass of Thong Pha Phum National Forest Thailand, *Ecology and Environmental Research*, **5**, 2, pp. 93-102.

Terakulpisut, J., Naroechaikusol, S. and Pitdamkham, C. (2000), Climate change and carbondioxide removal by forest in Huay Kha Khaeng Wildlife Sanctuary, Kasetsart University, Bangkok, Thailand, 60 pages (in Thai).

The statistic of forest land in Thailand of department of forestry, Ministry of Natural Resources and Environment, Thailand.

Viriyabancha, C. (2003), Handbook of Stand Biomass Estimation, Department of Nation Park Wildlife and Plant Conservation, 101 pages. (in Thai)

Wanthongchai, K., Bauhus, J. and Goldammer, J. G. (2008), Nutrient losses through prescribed burning of aboveground litter and understorey in dry dipterocarp forests of different fire history, *Catena*, **74**, pp. 321-332.

- Wanthongchai, K., Goldammer, J.G., and Bauhus, J. (2011), Effects of fire frequency on prescribed fire behavior and soil temperatures in dry dipterocarp forests, *International Journal of Wildland Fire*, **20**, pp. 35-45.
- Wang, W., Yue, X., Liu, H.J., Ding, Y. and Tang, D. G. (2001), Study on emission factors of particles from traffic sources, *Research Environmental and Science*, **14**, pp. 36-40.
- Ward, D. E., and Hao, W. M. (1992). Air toxic emissions from burning of biomass globally-preliminary estimate, paper presented in the, 85th Annual Meeting and Exhibition of the Air and Waste Management Association, Kansas city, Missouri.
- Ward, D. E., Nelson, R. M.m and Adams, D. (1979). Forest fire smoke documentation, paper presented in the, 72nd Annual Meeting, June 21-29, Cincinnati, OH, Paper No. 79-6.3. Pittsburg, PA: Air Poll. Control Assn.
- William, T. "Jerry" Winbery, Jr., et al., (1999), Method IO-3.5 determination of metals in ambient particulate matter using inductively coupled plasma/mass spectrometry (ICP-MS), Center for environmental research information office of research and development U.S. Environmental Protection Agency Cincinnati, OH 45268. EPA/625/R-96/010a, pp. 1-31.
- William A G et al., (2008), Forest Structure and Downed Woody Debris in Boreal, Temperate, and Tropical Forest Fragments, *Ambio*, **7**, pp. 7-8.
- Yamasoe, M. A., Artaxo, P., Miguel, A.M., Allen, A.G. (2000), Chemical composition of aerosol particulates from direct emissions of vegetation fires in the Amazon Basin: water-soluble species and trace elements, *Journal of Atmospheric Environment*, **34**, pp. 1641-1653.
- Yen, T. M. and Lee, J. S. (2011), Comparing aboveground carbon sequestration between moso bamboo (*Phyllostachys heterocycla*) and China fir (*Cunninghamia lanceolata*) forests based on the allometric model, *Forest Ecology and Management*, **261**, pp 995-1002.
- Zhou, G. M., and Jiang, P. K. (2004), Density, storage and spatial distribution of carbon in *Phyllostachy pubescens* forest, *Scientia Silvae Sinicae*, **40**, pp. 20-24. (In Chinese)