

## REFERENCES

- Aldor, Ilana S and Jay D Keaslingy. (2003), Process design for microbial plastic factories: Metabolic engineering of polyhydroxyalkanoates, *Current Opinion in Biotechnology*, **14**, 5, pp. 475-483.
- Alias, Z. and Irene K.P.Tan. (2005), Isolation of palm oil-utilising, polyhydroxyalkanoate (pha)-producing bacteria by an enrichment technique, *Bioresource Technology*, **96**, 11, pp. 1229-1234.
- Allmendinger, M., R. Eberhardt, G. A. Luinstra, F. Molnar and B. URL Rieger. (2003), The alternating copolymerization of epoxides and carbon monoxide: A novel way to polyesters, *Beyond Metallocenes: Next Generation Polymerization Catalysts*, **857**, pp. 114-130.
- Almeida, A., P. Nickel, A. Giordano, B. Mendez and J. URL Pettinari. (2007), Effect of the granule associated protein phasin (phap) on cell growth and poly(3-hydroxybutyrate) (phb) accumulation from glycerol in bioreactor cultures of recombinant *e-coli*, *Journal of Biotechnology*, **131**, 2, pp. S167-S167.
- Almeida, A., P. I. Nickel, A. M. Giordano and M. J. URL Pettinari. (2007), Effects of granule-associated protein phap on glycerol-dependent growth and polymer production in poly(3-hydroxybutyrate)-producing *eschetichia coli*, *Applied and Environmental Microbiology*, **73**, 24, pp. 7912-7916.
- Altschul, SF, W Gish, W Miller, EW Myers and DJ Lipman. (1990), Basic local alignment search tool, *Journal of Molecular Biology*, **215**, pp. 403-410.
- Anderson, A.J. and Dawes E.A. (1990), Occurrence, metabolism, metabolic role, and industrial uses of bacterial polyhydroxyalkanoates, *Microbiology Review*, **54**, pp. 450-472.
- Ashby, Foglia TA, Solaiman DKY, Liu C-K, Nunez A and Eggink G URL. (2000), Viscoelastic properties of linseed oil-based medium chain length poly(hydroxyalkanoate) films: Effects of epoxidation and curing, *International Journal of Biology and Macromolecule*, **27**, pp. 355-361.
- Ashby, D. K. Solaiman and T. A. Foglia. (2005), Synthesis of short-/medium-chain-length poly(hydroxyalkanoate) blends by mixed culture fermentation of glycerol, *Biomacromolecules*, **6**, 4, pp. 2106-2112.

- Ashby, D. K. Y. Solaiman and T. A. Foglia. (2004), Bacterial poly(hydroxyalkanoate) polymer production from the biodiesel co-product stream, *Journal of Polymers and the Environment*, **12**, 3, pp. 105-112.
- Barnard, G.N. and J.K. Sanders. (1989), The poly- $\beta$ -hydroxybutyrate granule in vivo. A new insight based on nmr spectroscopy of whole cells, *The Journal of Biological Chemistry*, **264**, 6, pp. 3286-3291.
- Beaulieu, M., Y. Beaulieu, J. Melinard, S. Pandian and J. Goulet. (1995), Influence of ammonium salts and cane molasses on growth of *alcaligenes eutrophus* and production of polyhydroxybutyrate, *Applied and Environment Microbiology*, **61**, 1, pp. 165-169.
- Belfars, L., M. Perrier, B.R. Ramsay, M. Jolicoeur and C. Chavari. (1995), Multi inhibition kinetic model for the growth of *alcagenes eutrophus*, *Canadian Journal of Microbiology*, **41**, pp. 249-256.
- Benizria, E., S. Piuttia, S. Vergerb, L. Pagesc, G. Vercambrec, J.L. Poesseld and P. Michelote. (2005), Replant diseases: Bacterial community structure and diversity in peach rhizosphere as determined by metabolic and genetic fingerprinting, *Soil Biology and Biochemistry*, **37**, pp. 1738-1746.
- Beun, J.J, K Dircks, MCM. Van Loosdrecht and Heijnen JJ. (2002), Poly-3-hydroxybutyrate metabolism in dynamically fed mixed microbial cultures, *Water Research*, **36**, 5, pp. 1167-1180.
- Bormann, E.J and Roth URL M. (1999), The production of polyhydroxybutyrate by *methylobacterium rhodesianum* and *ralstonia eutropha* in media containing glycerol and casein hydrolysates, *Biotechnology Letters*, **21**, pp. 1059-1063.
- Borneman, J. and Triplett EW. (1997), Molecular microbial diversity in soils from eastern amazonia: Evidence for unusual microorganisms and microbial population shifts associated with deforestation, *Applied and Environment Microbiology*, **63**, pp. 2647-2653.
- Boustead, I. (1999). "Eco-profiles of plastics and related intermediates (about 55 products)". Available online: <http://www.apme.org>
- Bramer, C. O., P. Vandamme, L. F. da Silva, J. G. C. Gomez and A. URL Steinbuchel. (2001), *Burkholderia sacchari* sp nov., a polyhydroxyalkanoate-accumulating bacterium isolated from soil of a sugar-cane plantation in brazil, *International Journal of Systematic and Evolutionary Microbiology*, **51**, pp. 1709-1713.

- Braunegg, G, Lefebvre G and Genser KF URL. (1998), Polyhydroxyalkanoates, biopolyesters from renewable resources: Physiological and engineering aspects, *Journal of Biotechnology*, **65**, pp. 127-161.
- Brioudea, Michel de Meireles, Danilo Hansen Guimarães, Raigenis da Paz Fiúzaa, Luis Antônio Sanches de Almeida Pradob, Jaime Soares Boaventuraa and Nadia Mamede Joséa. (2007), Synthesis and characterization of aliphatic polyesters from glycerol, by-product of biodiesel production, and adipic acid, *Materials Research*, **10**, 4, pp. 335-339.
- Çalık, Pınar, Serpil Takacı, Güzide Çalık and Tunçer H Özdamar. (2000), Serine alkaline protease overproduction capacity of *bacillus licheniformis*, *Enzyme and Microbial Technology*, **26**, 1, pp. 45-60.
- Celik, G. Y., Y. Beyatli and B. Aslim. (2005), Determination of poly-β-hydroxybutyrate (phb) production in sugarbeet molasses by *pseudomonas cepacia* g13 strain, *Zuckerindustrie*, **130**, 3, pp. 201-203.
- Chaijamrus, S. and N. Udpuay. (2008), Production and characterization of polyhydroxybutyrate from molasses and corn steep liquor produced by *bacillus megaterium* atcc 6748, *Agricultural Engineering International: CIGR Journal*, **10**
- Chen, G. Q. and Q. URL Wu. (2005), The application of polyhydroxyalkanoates as tissue engineering materials, *Biomaterials*, **26**, 33, pp. 6565-6578.
- Choi and S.Y Lee. (1999), Factors affecting the economics of polyhydroxyalkanoate production by bacterial fermentation, *Applied Microbiology and Biotechnology*, **51**, 1, pp. 13-21.
- Choi and Lee SY. (1997), Process analysis and economic evaluation for poly (3-hydroxybutyrate) production by fermentation, *Bioprocess Engineering*, **17**, pp. 335-342.
- Choi, M.H. and S.C. Yoon. (1994), Polyester biosynthesis characteristics of *pseudomonas citronellosis* grown on various carbon sources, including 3-methyl-branched substrates, *Applied and Environment Microbiology*, **60**, pp. 3245–3254.
- Chua, A.S.M., H. Takabatake, H. Satoh and T. Mino. (2003), Production of polyhydroxyalkanoates (pha) by activated sludge treating municipal wastewater: Effect of ph, sludge retention time (srt), and acetate concentration in influent, *Water Research*, **37**, 15, pp. 3602–3611.

- Ciesielski, S, A. Cydzik-Kwiatkowska, T. Pokoj and E. Klimiuk. (2006), Molecular detection and diversity of medium-chain-length polyhydroxyalkanoates-producing bacteria enriched from activated sludge, *Journal of Applied Microbiology*, **101**, 1, pp. 190-199.
- Ciesielski, S, Ewa Klimiuk, Justyna Mozejko, Ewa Nowakowska and Tomasz Pokoj. (2009), Changes in microbial community structure during adaptation towards polyhydroxyalkanoates production, *Polish Journal of Microbiology*, **58**, 2, pp. 131-139.
- Ciesielski, S, T. Pokoj and E. Klimiuk. (2010), Cultivation-dependent and -independent characterization of microbial community producing polyhydroxyalkanoates from raw-glycerol, *Journal of Microbiology and Biotechnology*, **20**, 5, pp. 853-861.
- Comstock, K, Farrell D and Xi Y Godwin C. (2004), From hydrocarbons to carbohydrates: Food packaging of the future,
- Dawes, E.A. and P.J. Senior. (1973), The role and regulation of energy reserve polymers in micro-organisms, *Advances in Microbial Physiology*, **10**, pp. 135-266.
- Degli, I.F. (2003), *Biodegradability and compostability*, Kluwer Academic Plenum Publishers, New York.
- Dias, J. M., P. C. Lemos, L. S. Serafim, C. Oliveira, M. Eiroa, M. G. Albuquerque, A. M. Ramos, R. Oliveira and M. A. Reis. (2006), Recent advances in polyhydroxyalkanoate production by mixed aerobic cultures: From the substrate to the final product, *Macromolecular Bioscience*, **6**, 11, pp. 885-906.
- Dionisi, D., M. Majone, V. Papa and M. URL Beccari. (2004), Biodegradable polymers from organic acids by using activated sludge enriched by aerobic periodic feeding, *Biotechnology and Bioengineering*, **85**, 6, pp. 569-579.
- El-Sayed, A. Azhar, Hemmat M. Abdelhady, A.M. Abdel Hafez and T.A. Khodair. (2009), Batch production of polyhydroxybutyrate (phb) by *ralstonia eutropha* and *alcaligenes latus* using bioreactor different culture strategies, *Journal of Applied Sciences Research*, **5**, 5, pp. 556-564.
- Fei, B., C. Chen, S. Chen, S. W. Peng, Y. G. Zhuang, Y. X. An and L. S. Dong. (2004), Crosslinking of poly[(3-hydroxybutyrate)-co-(3-hydroxyvalerate)] using dicumyl peroxide as initiator, *Polymer International*, **53**, 7, pp. 937-943.
- Fiedler, S., A. Steinbuchel and B. H. A. URL Rehm. (2002), The role of the fatty acid b-oxidation multienzyme complex from *pseudomonas oleovorans* in

- polyhydroxyalkanoate biosynthesis: Molecular characterization of the fadba operon from *p. Oleovorans* and of the enoyl-coa hydratase genes phaj from *p. Oleovorans* and *pseudomonas putida*, *Archives of Microbiology*, **178**, 2, pp. 149-160.
- Fisher and E. W. Triplett. (1999), Automated approach for ribosomal intergenic spacer analysis of microbial diversity and its application to freshwater bacterial communities, *Applied and Environment Microbiology*, **65**, 10, pp. 4630-4636.
- Foster, L. J. R., V. Sanguanchaipaiwong, C. L. Gabelish, J. Hook and M. Stenzel. (2005), A natural-synthetic hybrid copolymer of polyhydroxyoctanoate-diethylene glycol: Biosynthesis and properties, *Polymer*, **46**, 17, pp. 6587-6594.
- Fox, P. and F.G. Pohland. (1994), Anaerobic treatment applications and fundamentals: Substrate specificity during phase separation, *Water Environment Research*, **66**, pp. 716-724.
- Fromin, N., J. Hamelin, S. Tarnawski, D. Roesti, K. Jourdain-Miserez, N. Forestier, S. Teyssier-Cuvelle, F. Gillet, M. Aragno and P. Rossi. (2002), Statistical analysis of denaturing gel electrophoresis (dge) fingerprinting patterns, *Environmental Microbiology*, **4**, pp. 634-643.
- Fuhs, G. W. and M. Chen. (1975), Microbiological basis of phosphate removal in the activated sludge process for the treatment of wastewater, *Microbial Ecology*, **2**, pp. 119-138.
- Fuller, R.C, Odonnell J.P, Saulnier J, Redlinger T.E, Foster J and Lenz R.W. URL. (1992), The supramolecular architecture of the polyhydroxyalkanoate inclusions in *pseudomonas oleovorans*, *FEMS Microbiology Review*, **103**, 2-4, pp. 279-288.
- Garcia-Ribera, R., M. Monteoliva-Sanchez and A. Ramos-Cormenzana. (2001), Production of polyhydroxyalkanoates by *pseudomonas putida* kt2442 harboring psk2665 in wastewater from olive oil mills, *Journal of Biotechnology*, **4**, pp. 116-119.
- García, Marcela Hernández. (2008), Novel s-triazine-degrading bacteria isolated from agricultural soils of central chile for herbicide bioremediation, *Journal of Plant Nutrition and Soil Science*, **8**, 2, pp. 19-30.
- Gerngross. (1999), Can biotechnology move us toward a sustainable society?, *Nature Biotechnology*, **1**, 6, pp. 541-544.
- Gerngross and S. Slater. (2000), How green are green plastics?, *Scientific American*, pp. 37-41.

- Ghatnekar, M. S., J. S. Pai and M. URL Ganesh. (2002), Production and recovery of poly-b-hydroxybutyrate from *methylobacterium* sp v49, *Journal of Chemical Technology and Biotechnology*, **77**, 4, pp. 444-448.
- Gilbride, KA, Lee DY and Beaudette LA. M. (2006), Molecular techniques in wastewater: Understanding microbial communities, detecting pathogens, and real-time process control, *Journal of Microbiological Methods*, **66**, pp. 1-20.
- Green, P. R., J. Kemper, L. Schechtman, L. Guo, M. Satkowski, S. Fiedler, A. Steinbuchel and B. H. Rehm. (2002), Formation of short chain length/medium chain length polyhydroxyalkanoate copolymers by fatty acid b-oxidation inhibited *ralstonia eutropha*, *Biomacromolecules*, **3**, 1, pp. 208-213.
- Gürtler, V. and V. A. Stanisich. (1996), New approaches to typing and identification of bacteria using the 16s-23s rdna spacer region, *Microbiology*, **142**, pp. 3-16.
- Han, J., Y. Z. Qiu, D. C. Liu and G. Q. Chen. (2004), Engineered *aeromonas hydrophila* for enhanced production of poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) with alterable monomers composition, *FEMS Microbiology Letters*, **239**, 1, pp. 195-201.
- Hauschild, MZ. (2005), Assessing environmental impacts in a life cycle perspective, *Environmental Science and Technology*, **39**, pp. 905-912.
- Hawkes, FR., Hussy I, Kyazze G, Dinsdale R and Hawkes DL. (2007), Continuous dark fermentative hydrogen production by *mesophilic microflora*: Principles and progress, *International Journal of Hydrogen Energy*, **32**, 2, pp. 172-184.
- Haywood, G. W., A. J. Anderson, D. R. Williams, E. A. Dawes and D. F. Ewing. (1991), Accumulation of a poly(hydroxyalkanoate) copolymer containing primarily 3-hydroxyvalerate from simple carbohydrate substrates by *rhodococcus* sp. Ncimb 40126, *International Journal of Biological Macromolecules*, **13**, 2, pp. 83-88.
- Heinzle, E. and R.M. Lafferty. (1980), A kinetic model for the growth and synthesis of poly-b-hydroxybutyric acid (phb) in *alcaligenes eutrophus* h16, *European Journal of Applied Microbial Biotechnology*, **11**, pp. 8-16.
- Helm and Naumann D. (1995), Identification of some bacterial cell components by ft-ir spectroscopy, *FEMS Microbiology Letters*, **126**, pp. 75-80.
- Helm, Labischinski H, Schallehn G and Naumann D. (1991), Classification and identification of bacteria by fourier-transform infrared spectroscopy, *Journal of General Microbiology*, **137**, pp. 69-79.

- Hesselmanna, Rolf P.X., Thomas Fleischmann, Roland Hanyb and Alexander J.B. Zehnder. (1999), Determination of polyhydroxyalkanoates in activated sludge by ion chromatographic and enzymatic methods, *Journal of Microbiological Methods*, **35**, pp. 111-119.
- Hoffmann, N. and B. H. Rehm. (2005), Nitrogen-dependent regulation of medium-chain length polyhydroxyalkanoate biosynthesis genes in *pseudomonads*, *Biotechnology Letters*, **27**, 4, pp. 279-282.
- Holmes, P.A. (1985), Applications of phb: A microbially produced biodegradable thermoplastic, *Physics in Technology*, **16**, pp. 32-36.
- Hori, K., M. Kaneko, Y. Tanji, X. H. Xing and H. Unno. (2002), Construction of self-disruptive *bacillus megaterium* in response to substrate exhaustion for polyhydroxybutyrate production, *Applied Microbiology and Biotechnology*, **59**, 2-3, pp. 211-216.
- Hu, W.F, H. Chua and P.H.F. Yu. (1997), Synthesis of poly(3-hydroxybutyrate-co-3-hydroxyvalerate) from activated sludge, *Biotechnology Letters*, **19**, 7, pp. 695–698.
- Huijberts, G.N.M, Eggink G, Dewaard P, Huisman GW and Witholt B. (1992), *Pseudomonas putida* kt2442 cultivated on glucose accumulates poly(3-hydroxyalkanoates) consisting of saturated and unsaturated monomers, *Applied and Environment Microbiology*, **58**, pp. 536-544.
- Hustedede, E and Steinbuchel A. (1993), Characterization of the polyhydroxyalkanoate synthase gene locus of *rhodobacter sphaeroides*, *Biotechnology Letters*, **15**, 7, pp. 709-714.
- Iwamoto, T., K. Tani, K. Nakamura, Y. Suzuki, N. Kitagawa, M. Eguchi and M. Nasu. (2000), Monitoring impact of in situ biostimulation treatment on groundwater bacterial community by dgge, *Fems Microbiology Ecology*, **32**, pp. 129– 141.
- Joynt, J., M. Bischoff, R. Turco, A. Konopka and C. H. Nakatsu. (2006), Microbial community analysis of soils contaminated with lead, chromium and petroleum hydrocarbons, *Microbial Ecology*, **51**, 2, pp. 209-219.
- Judge, B. M., R. E. Howells, N. B. Graham and M. E. McNeill. (1981), Sustained-release implants in the chemotherapy of experimental rodent malaria ii. The effects of sulphadiazine, pyrimethamine and cycloguanil in biodegradable polymer matrices, *Annals of Tropical Medicine and Parasitology*, **75**, 5, pp. 511-519.

- Kadouri, D., E. Jurkevitch, Y. Okon and S. Castro-Sowinski. (2005), Ecological and agricultural significance of bacterial polyhydroxyalkanoates, *Critical Reviews in Microbiology*, **31**, pp. 55-67.
- Kaewpipat, K. and C. P. L. Grady. (2002), Microbial population dynamics in laboratory-scale activated sludge reactors, *Water Science and Technology*, **46**, 1-2, pp. 19-27.
- Kahar, P., J. Agus, Y. Kikkawa, K. Taguchi, Y. Doi and T. Tsuge. (2005), Effective production and kinetic characterization of ultra-high-molecular-weight poly[(r)-3-hydroxybutyrate] in recombinant *escherichia coli*, *Polymer Degradation and Stability*, **87**, 1, pp. 161-169.
- Kansiz, M., A. Dominguez-Vidal, D. McNaughton and B. URL Lendl. (2007), Fourier-transform infrared (ftir) spectroscopy for monitoring and determining the degree of crystallisation of polyhydroxyalkanoates (phas), *Analytical and Bioanalytical Chemistry*, **388**, 5-6, pp. 1207-1213.
- Karahan, O., Derin Orhon and Mark C. M. van Loosdrecht. (2008), Simultaneous storage and utilization of polyhydroxyalkanoates and glycogen under aerobic conditions, *Water Science and Technology*, pp. 945-951.
- Katja, Johnson, Jorien van Geest, Robbert Kleerebezem and Mark C.M. van Loosdrecht. (2010), Short- and long-term temperature effects on aerobic polyhydroxybutyrate producing mixed cultures, *Water Research*, **44**, pp. 1689-1700.
- Kessler, B. and B. URL Witholt. (2001), Factors involved in the regulatory network of polyhydroxyalkanoate metabolism, *Journal of Biotechnology*, **86**, 2, pp. 97-104.
- Khanna, S. and A. K. URL Srivastava. (2005), A simple structured mathematical model for biopolymer (phb) production, *Biotechnology Progress*, **21**, 3, pp. 830-838.
- Kim, B.S., S.C. Lee, S.Y. Lee, H.N. Chang, Y.K. Chang and S.I. Woo. (1994), Production of poly(3-hydroxybutyric acid) by fed-batch culture of *alcaligenes eutrophus* with glucose concentration control, *Biotechnology and Bioengineering*, **43**, pp. 892-898.
- Kirk, J.L, L.A Beaudette, M Hart, P Moutoglis, J.N Klironomos and H Lee. (2004), Methods of studying soil microbial diversity, *Journal of Microbiological Methods*, **58**, pp. 169-188.
- Kleerebezem, R. and M. C. M. van Loosdrecht. (2007), Mixed culture biotechnology for bioenergy production, *Current Opinion in Biotechnology*, **18**, pp. 207-212.

- Koller, M., R. Bona, G. Braunegg, C. Hermann, P. Horvat, M. Kroutil, J. Martinz, J. Neto, L. Pereira and P. URL Varila. (2005), Production of polyhydroxyalkanoates from agricultural waste and surplus materials, *Biomacromolecules*, **6**, 2, pp. 561-565.
- Kumar, R. and R. P. URL Singh. (2001), Semi-solid-state fermentation of eicchornia crassipes biomass as lignocellulosic biopolymer for cellulase and b-glucosidase production by cocultivation of *aspergillus niger* rk3 and *trichoderma reesei* mtcc164, *Applied Biochemistry and Biotechnology*, **96**, 1-3, pp. 71-82.
- Kurdikar, D., M Paster, KJ Gruys, L Fournet, TU Gerngross, SC Slater and R Coulon. (2001), Greenhouse gas profile of a plastic material derived from a genetically modified plant, *Journal of Industrial Ecology*, **4**, 3, pp. 107–122.
- Labuzek, S. and I. URL Radecka. (2001), Biosynthesis of phb tercopolymer by *bacillus cereus* uw85, *Journal of Applied Microbiology*, **90**, 3, pp. 353-357.
- Lafferty, R. M., B. Korsatko and W. Korsatko. (1988), *Microbial production of poly- $\beta$ -hydroxybutyric acid*, VCH Publishers, New York.
- Lageveen, R. G., G. W. Huisman, H. Preusting, P. Ketelaar, G. Eggink and B. Witholt. (1988), Formation of polyesters by *pseudomonas oleovorans*: Effect of substrates on formation and composition of poly-(r)-3-hydroxyalkanoates and poly-(r)-3-hydroxyalkenoates, *Applied and Environment Microbiology*, **54**, pp. 2924-2932.
- Landis, AE., Miller SA and Theis TL. (2007), Life cycle of the corn-soybean agroecosystem for bio-based production, *Environmental Science and Technology*, **41**, pp. 1457–1464.
- Larsson, Gen and Mari Törnkvist. (1996), Rapid sampling, cell inactivation and evaluation of low extracellular glucose concentrations during fed-batch cultivation, *Journal of Biotechnology*, **49**, 1-3, pp. 69-82.
- Law, J. and R. A Slepecky. (1969), Assay of poly-b-hydroxybutyric acid, *Journal of Bacteriology*, **82**, pp. 52-55.
- Lee. (1996), Bacterial polyhydroxyalkanoates, *Biotechnology and Bioengineering*, **49**, 1, pp. 1-14.
- Lee. (1996), Plastic bacteria? Progress and prospects for polyhydroxyalkanoate production in bacteria, *Trends in Biotechnology*, **14**, 11, pp. 431-438.
- Lee, K. M. Lee, H. N. Chan and A. Steinbuchel. (1994), Comparison of recombinant *escherichia coli* strains for synthesis and accumulation of poly-(3-hydroxybutyric

- acid) and morphological changes, *Biotechnology and Bioengineering*, **44**, 11, pp. 1337-1347.
- Lee, S. W. Nam, Y. H. Rhee and J. Y. Kim. (1996), Cloning and functional expression in *escherichia coli* of the polyhydroxyalkanoate synthase (phac) gene from *alcaligenes* sp sh-69, *Journal of Microbiology and Biotechnology*, **6**, 5, pp. 309-314.
- Lee, H.D. Shin and Y.H. Lee. (2003), Metabolic engineering of pentose phosphate pathway in *ralstonia eutropha* for enhanced biosynthesis of poly-b-hydroxybutyrate, *Biotechnology Progress*, **19**, 5, pp. 1444-1449.
- Lee and Y.J. Yoo. (1994), High cell density culture of *alcaligenes eutrophus* and polyhydroxybutyrate production by optimization of medium compositions, *Journal of Applied Microbiology and Biotechnology*, **22**, pp. 401-406.
- Lee and Y. J. Yoo. (1991), Kinetics for the growth of *alcaligenes eutrophus* and the biosynthesis of poly-b-hydroxybutyrate, *Korean Journal of Applied Microbial Biotechnology*, **19**, pp. 186-192.
- Leeuwen, M.A. Van Aalst-van, Pot. MA., van Loosdrecht. MCM and Heijnen JJ. (1997), Kinetic modeling of poly ( $\beta$ -hydroxybutyrate): Production and consumption by *paraccus pantotrophus* under dynamic substrate supply, *Biotechnology and Bioengineering*, **55**, pp. 773-782.
- Lemoigne, M. (1926), Produits de deshydratation et the polymerisation de acide b-oxybutirique, *Bull. Soc. Chim. Biol*, **8**, 770,
- Lemos, P. C., L. S. Serafim and M. A. M. Reis. (2004), Polyhydroxyalkanoates production by activated sludge in a sbr using acetate and propionate as carbon sources, *Water Science and Technology*, **50**, 10, pp. 189-194.
- Lerman, L. S. and K. Silverstein. (1987), Computational simulation of DNA melting and its application to denaturing gradient gel electrophoresis, *Methods in Enzymology*, **155**, pp. 482-501.
- Lettinga, G. (1995), Anaerobic digestion and wastewater treatment systems, *Antonie Van Leeuwenhoek*, **67**, 1, pp. 3-28.
- Liu, W. T., T. Mino, T. Matsuo and K. URL Nakamura. (2000), Isolation, characterization and identification of polyhydroxyalkanoate-accumulating bacteria from activated sludge, *Journal of Bioscience and Bioengineering*, **90**, 5, pp. 494-500.

- Lopez-Cortes, A., A. Lanz-Landazuri and J. Q. Garcia-Maldonado. (2008), Screening and isolation of phb-producing bacteria in a polluted marine microbial mat, *Microbial Ecology*, **56**, 1, pp. 112-120.
- Lyons, J. (1994), Identification of mutant forms of g-protein a subunits in human neoplasia by polymerase chain reaction-based techniques, *Methods in Enzymology*, **237**, pp. 295-308.
- Ma, T. C., P. L. Chan, H. Lawford, H. Chua, W. H. Lo and P. H. URL Yu. (2005), Microbial synthesis and characterization of physiochemical properties of polyhydroxyalkanoates (phas) produced by bacteria isolated from activated sludge obtained from the municipal wastewater works in hong kong, *Applied Biochemistry and Biotechnology*, **121**, pp. 731-739.
- Madison and Huisman G.W. (1999), Metabolic engineering of poly(3-hydroxyalkanoates): From DNA to plastic, *Microbiological Molecular and Biological Review*, **63**, 1, pp. 21-53.
- Mahishi, L. H., G. Tripathi and S. K. Rawal. (2003), Poly(3-hydroxybutyrate) (phb) synthesis by recombinant *escherichia coli* harbouring *streptomyces aureofaciens* phb biosynthesis genes: Effect of various carbon and nitrogen sources, *Microbiological Research*, **158**, 1, pp. 19-27.
- Majone, M, Beccari M and Ramadori R. (Year). Biodegradable polymers from wastes by using activated sludges enriched by aerobic periodic feeding, *AIDIC Conference Series*, pp 163-170.
- Miskin, I.P., P. Farrimond and I.M. Head. (1999), Identification of novel bacterial lineages as active members of microbial populations in a freshwater sediment using a rapid rna extraction procedure and rt-pcr, *Microbiology*, **145**, pp. 1977-1987.
- Misra, S. K., S. P. Valappil, I. Roy and A. R. URL Boccaccini. (2006), Polyhydroxyalkanoate (pha)/inorganic phase composites for tissue engineering applications, *Biomacromolecules*, **7**, 8, pp. 2249-2258.
- Moeseneder, M. M., J. M. Arrieta, G. Muyzer, C. Winter and G. J. Herndl. (1999), Optimization of terminal-restriction fragment length polymorphism analysis for complex marine bacterioplankton communities and comparison with denaturing gradient gel electrophoresis, *Applied and Environment Microbiology*, **65**, 8, pp. 3518-3525.

- Mukhopadhyay, M., A. Patra and A. K. Paul. (2005), Production of poly(3-hydroxybutyrate) and poly(3-hydroxybutyrate-co-3-hydroxyvalerate) by *rhodospseudomonas palustris* sp5212, *World Journal of Microbiology and Biotechnology*, **21**, 5, pp. 765-769.
- Muyzer, G. and K. Smalla. (1998), Application of denaturing gradient gel electrophoresis (dgge) and temperature gradient gel electrophoresis (tgge) in microbial ecology, *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology*, **73**, pp. 127-141.
- Myers, R. M., S. G. Fischer, L. S. Lerman and T. Maniatis. (1985), Nearly all single base substitutions in DNA fragments joined to a gc-clamp can be detected by denaturing gradient gel electrophoresis, *Nucleic Acids Research*, **13**, 9, pp. 3131-3145.
- Nakatsu, C.H., V. Torsvik and L. Øvreås. (2000), Soil community analysis using dgge of 16s rDNA polymerase chain reaction products, *Soil Science Society of America Journal*, **64**, pp. 1382-1388.
- Naumann, D, Keller S, Helm D, Schultz NC and Schrader B. (1995), Ft-ir spectroscopy and ft-Raman spectroscopy are powerful analytical tools for the non-invasive characterization of intact microbial cells, *Journal of Molecular Structure*, **347**, pp. 399-406.
- Nicols, PD, Henson JM, Guckert JB, Nivens DE and White DC. (1984), Fourier transform-infrared spectroscopic methods for microbial ecology: Analysis of bacteria, bacterial polymer mixtures and biofilms, *Letters in Applied Microbiology*, **4**, pp. 79-94.
- Nikel, P. I., M. J. Pettinari, M. A. Galvagno and B. S. URL Mendez. (2006), Poly(3-hydroxybutyrate) synthesis by recombinant *Escherichia coli* arca mutants in microaerobiosis, *Applied and Environmental Microbiology*, **72**, 4, pp. 2614-2620.
- Noda, I., M. M. Satkowski, A. E. Dowrey and C. Marcott. (2004), Polymer alloys of nodax copolymers and poly(lactic acid), *Macromolecular Bioscience*, **4**, 3, pp. 269-275.
- Nübel, U., B. Engelen, A. Felske, J. Snaidr, A. Wieshuber, R.I. Amann, W. Ludwig and H. Backhaus. (1996), Sequence heterogeneities of genes encoding 16s rnas in *paenibacillus polymyxa* detected by temperature gradient gel electrophoresis, *Journal of Bacteriology*, **178**, pp. 5636-5643.
- Nubel, U., Garcia-Pichel F and Muyzer G. (1997), Pcr primers to amplify 16s rna genes from cyanobacteria, *Applied and Environment Microbiology*, **63**, pp. 3327-3332.

- Ojumu, T.V., J. Yu and B.O. Solomon. (2004), Production of polyhydroxyalkanoates, a bacterial biodegradable polymer, *African Journal of Biotechnology*, **3**, 1, pp. 18-24.
- Ouyang, S. P., Q. Liu, L. Fang and G. Q. Chen. (2007), Construction of pha-operon-defined knockout mutants of *pseudomonas putida* kt2442 and their applications in poly(hydroxyalkanoate) production, *Macromolecular Bioscience*, **7**, 2, pp. 227-233.
- Page, W.J. (1989), Production of poly-b-hydroxybutyrate by *azotobacter vinelandii* strain uwd during growth on molasses and other complex carbon sources, *Applied Microbiology and Biotechnology*, **31**, pp. 329-333.
- Page, W.J. (1995), Bacterial polyhydroxyalkanoates, natural biodegradable plastics with a great future, *Canadian Journal of Microbiology*, **41**, 1, pp. 1-3.
- Patel, J., K. Patel and R. Patel. (2009), Reactivity ratio of novel acrylic copolymer by nmr spectroscopy, *Colloid and Polymer Science*, **287**, 1, pp. 89-95.
- Peoples, O. P. and A. J. Sinskey. (1989), Poly-b-hydroxybutyric biosynthesis in *alcaligenes eutrophus* h16. Identification and characterization of the phb polymerase gene (phbc), *Journal of Biological Chemistry*, **264**, pp. 15298–15303.
- Poirier, Y., C. Somerville, L. A. Schechtman, M. M. Satkowski and I. Noda. (1995), Synthesis of high-molecular-weight poly([r]-(-)-3-hydroxybutyrate) in transgenic arabidopsis thaliana plant cells, *International Journal of Biological Macromolecules*, **17**, 1, pp. 7-12.
- Qi, Q. and B. H. Rehm. (2001), Polyhydroxybutyrate biosynthesis in *caulobacter crescentus*: Molecular characterization of the polyhydroxybutyrate synthase, *Microbiology*, **147**, Pt 12, pp. 3353-3358.
- Quillaguaman, J., O. Delgado, B. Mattiasson and R. URL Hatti-Kaul. (2006), Poly(b-hydroxybutyrate) production by a moderate halophile, *halomonas boliviensis* lc1, *Enzyme and Microbial Technology*, **38**, 1-2, pp. 148-154.
- Ramsay, B. A., I. Saracovan, J. A. Ramsay and R. H. Marchessault. (1992), Effect of nitrogen limitation on long-side-chain poly-b-hydroxyalkanoate synthesis by *pseudomonas resinovorans*, *Applied and Environment Microbiology*, **58**, 2, pp. 744-746.
- Rani, A., S. Porwal, R. Sharma, A. Kapley, H.J. Purohit and V.C. Kalia. (2008), Assessment of microbial diversity in effluent treatment plants by culture dependent and culture independent approaches, *Bioresource Technology*, **99**, pp. 7098–7107.

- Ranjard, L., F. Poly, J. C. Lata, C. Mougel, J. Thioulouse and S. Nazaret. (2001), Characterization of bacterial and fungal soil communities by automated ribosomal intergenic spacer analysis fingerprints: Biological and methodological variability, *Applied and Environmental Microbiology*, **67**, 10, pp. 4479-4487.
- Reddy, C.S.K., R. Ghai, Rashmi and V.C. Kalia. (2003), Polyhydroxyalkanoates: An overview, *Bioresource Technology*, **87**, pp. 137-146.
- Rehman, S., N. Jamil and S. Husnain. (2007), Screening of different contaminated environments for polyhydroxyalkanoates-producing bacterial strains, *Biologia*, **62**, 6, pp. 650-656.
- Reis, M. A. M., L. S. Serafim, P. C. Lemos, A. M. Ramos, F. R. Aguiar and M. C. M. Van Loosdrecht. (2003), Production of polyhydroxyalkanoates by mixed microbial cultures, *Bioprocess and Biosystems Engineering*, **25**, 6, pp. 377-385.
- Rittmann, B. E., M. Hausner, F. Löffler, N.G. Love, G. Muyzer, S. Oerther S. Okabe, D.B., J. Peccia, L. Raskin and M. M. Wagner. (2006), A vista for microbial ecology and environmental biotechnology, *Environmental Science and Technology*, **40**, pp. 1096-1103.
- Riviere, D, V Desvignes, E Pelletier, S Chaussonnerie, S Guermazi, J Weissenbach, T Li, P Camacho and A Sghir. (2009), Toward the definition of a core of microorganisms involved in anaerobic digestion of sludge, *Journal of the International Society for Microbial Ecology*, **3**, pp. 700-714.
- Sabirova, J. S., M. Ferrer, H. Lunsdorf, V. Wray, R. Kalscheuer, A. Steinbuchel, K. N. Timmis and P. N. URL Golyshin. (2006), Mutation in a "Tesb-like" Hydroxyacyl-coenzyme a-specific thioesterase gene causes hyperproduction of extracellular polyhydroxyalkanoates by *alcanivorax borkumensis* sk2, *Journal of Bacteriology*, **188**, 24, pp. 8452-8459.
- Sanz, J. L. and T. Köchling. (2007), Molecular biology techniques used in wastewater treatment: An overview, *Process Biochemistry*, **42**, pp. 119-133.
- Satkofsky, A. (2002), The status of degradable plastics for composting, *Biological Cycle*, **43**, 3, pp. 60-67.
- Satoh, Iwamoto Y, Mino T and Matsuo T. (1998), Activated sludge as a possible source of biodegradable plastics, *Water Science and Technology*, **38**, pp. 103-109.
- Satokari, R. M., E. E. Vaughan, A. D. L. Akkermans, M. Saarela and W. M. de Vos. (2001), Bifidobacterial diversity in human feces detected by genus-specific pcr and

- denaturing gradient gel electrophoresis, *Applied and Environmental Microbiology*, **67**, 2, pp. 504-513.
- Schubert, P., A. Steinbüchel and H. G. Schlegel. (1988), Cloning of *alcaligenes eutrophus* genes for synthesis of poly-b-hydroxybutyric acid (phb) and synthesis of phb in *escherichia coli*, *Journal of Bacteriology*, **170**, pp. 5837–5847.
- Serafim, L. S., P. C. Lemos, R. Oliveira and M. A. M. Reis. (2004), Optimization of polyhydroxybutyrate production by mixed cultures submitted to aerobic dynamic feeding conditions, *Biotechnology and Bioengineering*, **87**, 2, pp. 145-160.
- Shamala, T. R., A. Chandrashekar, S. V. N. Vijayendra and L. URL Kshama. (2003), Identification of polyhydroxyalkanoate (pha)-producing *bacillus* spp. Using the polymerase chain reaction (pcr), *Journal of Applied Microbiology*, **94**, 3, pp. 369-374.
- Shang, L., M. Jiang and H. N. Chang. (2003), Poly(3-hydroxybutyrate) synthesis in fed-batch culture of *ralstonia eutropha* with phosphate limitation under different glucose concentrations, *Biotechnology Letters*, **25**, 17, pp. 1415-1419.
- Simon, R., Priefer U and Pühler A. (1983), A broad host range mobilization system for in vivo genetic engineering: Transposon mutagenesis in gram negative bacteria, *Biotechnology*, **1**, pp. 784-791.
- Slater, S. C., W. H. Voige and D. E. Dennis. (1988), Cloning and expression in *escherichia coli* of the *alcaligenes eutrophus* h16 poly-b-hydroxybutyrate biosynthetic pathway, *Journal of Bacteriology*, **170**, pp. 4431–4436.
- Solaiman, D. K. Y., R. D. Ashby, T. A. Foglia and W. N. URL Marmer. (2006), Conversion of agricultural feedstock and coproducts into poly(hydroxyalkanoates), *Applied Microbiology and Biotechnology*, **71**, 6, pp. 783-789.
- Solaiman, D. K. Y., R. D. Ashby, A. T. Hotchkiss and T. A. URL Foglia. (2006), Biosynthesis of medium-chain-length poly(hydroxyalkanoates) from soy molasses, *Biotechnology Letters*, **28**, 3, pp. 157-162.
- Spiegelman, D., Whissell G and Greer CW. (2005), A survey of the methods for the characterization of microbial consortia and communities, *Canadian Journal of Microbiology*, **51**, pp. 355-386.
- Steinbüchel, A. (1992), Biodegradable plastics, *Current Opinion in Biotechnology*, **3**, pp. 291-297.

- Steinbüchel, A. and Schlegel H.G. (1989), Excretion of pyruvate by mutants of *alcaligenes eutrophus*, which are impaired in the accumulation of poly(hydroxybutyric acid) (phb), under conditions permitting synthesis of phb, *Applied Microbiology and Biotechnology*, **31**, pp. 168–175.
- Steinbüchel, A. and Valentin HE. (1995), Diversity of bacterial polyhydroxyalkanoic acids, *FEMS Microbiology Letters*, **128**, pp. 219-228.
- Steinbüchel, A. and T. URL Lutke-Eversloh. (2003), Metabolic engineering and pathway construction for biotechnological production of relevant polyhydroxyalkanoates in microorganisms, *Biochemical Engineering Journal*, **16**, 2, pp. 81-96.
- Steinbüchel, A. and H.G. Schlegel. (1991), Physiology and molecular genetics of poly(b-hydroxyalkanoic acid) synthesis in *alcaligenes eutrophus*, *Molecular Microbiology*, **5**, pp. 535–542.
- Stephanopoulos, Gregory N., Aristos A. Aristidou and Jens Nielsen. (1998), *Review of cellular metabolism*, Academic Press, San Diego.
- Stephen, J. R., G. A. Kowalchuk, M. A. V. Bruns, A. E. McCaig, C. J. Phillips, T. M. Embley and J. I. Prosser. (1998), Analysis of b-subgroup proteobacterial ammonia oxidizer populations in soil by denaturing gradient gel electrophoresis analysis and hierarchical phylogenetic probing, *Applied and Environment Microbiology*, **64**, 8, pp. 2958-2965.
- Sudesh, K, Fukui T, Iwata T and Doi Y URL. (2000), Factors affecting the freeze-fracture morphology of in vivo polyhydroxyalkanoate granules, *Canadian Journal of Microbiology*, **46**, 4, pp. 304-311.
- Sujatha, K.V.V.S.N. Bapi Raju and T. Ramana. (2005), Studies on a new marine streptomycete bt408 producing polyketide antibiotic sbr 22 effective against methicillin resistant *stephylococcus aureus*, *Research in Microbiology*, **160**, pp. 119-126.
- Sujatha, K. and R. Shenbagarathai. (2006), A study on medium chain length-polyhydroxyalkanoate accumulation in *escherichia coli* harbouring phac1 gene of indigenous *pseudomonas* sp. Ldc-5, *Letters in Applied Microbiology*, **43**, 6, pp. 607-614.
- Suzuki, T, T Yamane and S. Shimizu. (1986), Mass production of poly- $\beta$ -hydroxybutyric acid by fully automatic fed-batch culture of methylotroph, *Applied Microbiology and Biotechnology*, **23**, pp. 322-329.

- Tajima, K., T. Igari, D. Nishimura, M. Nakamura, Y. Satoh and M. URL Munekata. (2003), Isolation and characterization of *bacillus* sp int005 accumulating polyhydroxyalkanoate (pha) from gas field soil, *Journal of Bioscience and Bioengineering*, **95**, 1, pp. 77-81.
- Tajima, K., Y. Satoh, K. Nakazawa, H. Tannai, T. Erata, M. Munekata, M. Kamachi and R. W. Lenz. (2004), Chemoenzymatic synthesis of poly(3-hydroxybutyrate) in a water-organic solvent two-phase system, *Macromolecules*, **37**, 12, pp. 4544-4546.
- Takagi, Y., R. Yasuda, A. Maehara and T. URL Yamane. (2004), Microbial synthesis and characterization of polyhydroxyalkanoates with fluorinated phenoxy side groups from *pseudomonas putida*, *European Polymer Journal*, **40**, 7, pp. 1551-1557.
- Temudo, M. F., R. Kleerebezem and M. URL van Loosdrecht. (2007), Influence of the ph on (open) mixed culture fermentation of glucose: A chemostat study, *Biotechnology and Bioengineering*, **98**, 1, pp. 69-79.
- Thompson, J. D., D. G. Higgins and T. J. Gibson. (1994), Clustal w: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positionsspecific gap penalties and weight matrix choice, *Nucleic Acids Research*, **22**, pp. 4673-4680.
- Tobin, K. M. and K. E. O'Connor. (2005), Polyhydroxyalkanoate accumulating diversity of *pseudomonas* species utilising aromatic hydrocarbons, *FEMS Microbiology Letters*, **253**, 1, pp. 111-118.
- Tohyama, M, T Patarinska, Z Qiang and K Shimizu. (2002), Modeling of the mixed culture and periodic control for phb production, *Biochemical Engineering Journal*, **10**, pp. 157-173.
- Tsuge, T., T. Fukui, H. Matsusaki, S. Taguchi, G. Kobayashi, A. Ishizaki and Y. URL Doi. (2000), Molecular cloning of two (r)-specific enoyl-coa hydratase genes from *pseudomonas aeruginosa* and their use for polyhydroxyalkanoate synthesis, *FEMS Microbiology Letters*, **184**, 2, pp. 193-198.
- Tsunemasa, N. (1998), Utilization of excess sludge for the production of biodegradable plastics, *Water Research*, **40**, pp. 981-986.
- Ueno, Y., S. Haruta, M. Ishii and Y. Igarashi. (2001), Microbial community in anaerobic hydrogen-producing microflora enriched from sludge compost, *Applied Microbiology and Biotechnology*, **57**, 4, pp. 555-562.

- Valappil, S. P., D. Peiris, G. J. Langley, J. M. Hemiman, A. R. Boccaccini, C. Bucke and I. URL Roy. (2007), Polyhydroxyalkanoate (pha) biosynthesis from structurally unrelated carbon sources by a newly characterized *bacillus* spp., *Journal of Biotechnology*, **127**, 3, pp. 475-487.
- Valentin, H. E., Pierre A. Berger, Kenneth J. Gruys, Maria Filomena de Andrade Rodrigues, Alexander Steinbuchel, Minhtien Tran and Jawed Asrar. (1999), Biosynthesis and characterization of poly(3-hydroxy-4-pentenoic acid), *Macromolecules*, **32**, pp. 7389-7395.
- Wang and S. Y. Lee. (1997), Production of poly(3-hydroxybutyrate) by fed-batch culture of filamentation-suppressed recombinant *escherichia coli*, *Applied and Environment Microbiology*, **63**, 12, pp. 4765-4769.
- Wang and S.Y Lee. (1997), Poly(3-hydroxybutyrate) production with high productivity and high polymer content by a fed-batch culture of *alcaligenes latus* under nitrogen limitation, *Applied and Environment Microbiology*, **63**, 9, pp. 3703-3706.
- Wang, X Y. Sheng, R. C. Equi, M. A. Trainer, T. C. Charles and B. W. S. URL Sobral. (2007), Influence of the poly-3-hydroxybutyrate (phb) granule-associated proteins (phap1 and phap2) on phb accumulation and symbiotic nitrogen fixation in *sinorhizobium meliloti* rm1021, *Journal of Bacteriology*, **189**, 24, pp. 9050-9056.
- Ward, P. G. and K. E. URL O'Connor. (2005), Bacterial synthesis of polyhydroxyalkanoates containing aromatic and aliphatic monomers by *pseudomonas putida* ca-3, *International Journal of Biological Macromolecules*, **35**, 3-4, pp. 127-133.
- Williamson, D.H and Wilkinson J.F. (1958), The isolation and estimation of the poly-b-hydroxybutyrate inclusions of *bacillus* species, *General Microbiology*, **19**, pp. 198-209.
- Wong, P. A. L., M. K. Cheung, W. L. Lo, H. Chua and P. H. F. Yu. (2005), Effects of types of food waste as carbon source on the molecular weight distributions and thermal properties of the biopolymer (polyhydroxybutyrate) produced by two strains of microorganisms, *Materials Research Innovations*, **9**, 1, pp. 4-5.
- Xu, J., B. H. Guo, Z. M. Zhang, Q. Wu, Q. Zhou, J. C. Chen, G. Q. Chen and G. D. URL Li. (2005), A mathematical model for regulating monomer composition of the microbially synthesized polyhydroxyalkanoate copolymers, *Biotechnology and Bioengineering*, **90**, 7, pp. 821-829.

- Yamane. (1993), Yield of poly-d(-)-3-hydroxybutyrate from various carbon sources: A theoretical study, *Biotechnology and Bioengineering*, **41**, pp. 165-170.
- Yang, C.H. and D.E. Crowley. (2000), Rhizosphere microbial community structure in relation to root location and plant iron nutritional status, *Applied and Environment Microbiology*, **66**, pp. 345-351.
- Yaoping, LÜ. (2007), Advance on the production of polyhydroxyalkanoates by mixed cultures, *Frontiers of Biology in China*, **2**, 1, pp. 21-25.
- Yazdani, S. S. and R. Gonzalez. (2007), Anaerobic fermentation of glycerol: A path to economic viability for the biofuels industry, *Current Opinion in Biotechnology*, **18**, 3, pp. 213-219.
- Yilmaz, M., H. Soran and Y. Beyatli. (2005), Determination of poly-b-hydroxybutyrate (phb) production by some *bacillus* spp., *World Journal of Microbiology and Biotechnology*, **21**, 4, pp. 565-566.
- Yin, B., Crowley DE, Sparovek G, De Melo WJ and Borneman J. (2000), Bacterial functional redundancy along a soil reclamation gradient, *Applied and Environment Microbiology*, **66**, pp. 4361-4365.
- Yu, H Chua, A.L Huang and K.P. Ho. (1999), Conversion of industrial food wastes by *alcaligenes latus* into polyhydroxyalkanoates, *Applied Biochemistry and Biotechnology*, **77-79**, pp. 445-454.
- Yu, C. C. Lin and J. R. Too. (2005), Phbv production by *ralstonia eutropha* in a continuous stirred tank reactor, *Process Biochemistry*, **40**, 8, pp. 2729-2734.
- Zakaria, M.R., Hidayah Ariffin Suraini Abd-Aziz, NorAini Abdul Rahman, Phang Lai Yee and Mohd Ali Hassan. (2008), *Comamonas* sp. Eb172 isolated from digester treating palm oil mill effluent as potential polyhydroxyalkanoate (pha) producer, *African Journal of Biotechnology*, **7**, 22, pp. 4118-4121.
- Zevenhuizen, L.P.T.M and Ebbink A.G. (1974), Interrelations between glycogen, polyhydroxybutyric acid and lipids during accumulation and subsequent utilization in a *pseudomonas*, *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology*, **40**, 1, pp. 103-120.
- Zhang, G, Hang X, Green P, Ho K.P and Chen G.Q URL. (2001), Pcr cloning of type ii polyhydroxyalkanoate biosynthesis genes from two *pseudomonas* strains, *FEMS Microbiology Letters*, **198**, 2, pp. 165-170.

- Zhang, H., V. Obias, K. Gonyer and D. Dennis. (1994), Production of polyhydroxyalkanoates in sucrose-utilizing recombinant *escherichia coli* and *klebsiella* strains, *Applied and Environment Microbiology*, **60**, 4, pp. 1198-1205.
- Zoetemeyer, R.J., Arnoldy P, Cohen A and Boelhouwer C. (1982), Influence of temperature on the anaerobic acidification of glucose sin a mixed culture forming part of a 2-stage digestion process, *Water Research*, **16**, 3, pp. 313–321.