



REFERENCES

- Aherne, S.A. and O'Brien, N.M. 2002. Dietary flavonols: chemistry, food content, and metabolism. *Nutrition*, 18 (1):75– 81.
- Anwar-Mohamed, A. and El-Kadi, A.O. 2009. Down-regulation of the detoxifying enzyme NAD(P)H:quinone oxidoreductase 1 by vanadium in Hepa 1c1c7 cells. *Toxicol Appl. Pharmacol.*, 236(3): 261-269.
- Asano, N., Katsuma, Y., Tamura, H., Higashikuni, N. and Hayashi, M. 1998. An automated new technique for scoring the rodent micronucleus assay: computerized image analysis of acridine orange supravivally stained peripheral blood cells. *Mutat. Res.*, 404 (1-2): 149-154.
- Attasara, P., Buasom, R. Hospital-based cancer registry. National Cancer Institute, Bangkok 2008.
- Bannasch, P. 1986. Preneoplastic lesions as end points in carcinogenicity testing. I. Hepatic preneoplasia. *Carcinogenesis*, 7(5): 689-695.
- Benson, A.M., Hunkeler, M.J. and Talalay, P. 1980. Increase of NAD(P)H:quinone reductase by dietary antioxidants: Possible role in protection against carcinogenesis and toxicity. *Proc. Nat. Acad. Sci.*, 77(9): 5216–5220.
- Bernhardt, R. 2006. Cytochromes P450 as versatile biocatalysts. *J. biotechnol.*, 124: 128–145.
- Bianco, N.R., Chaplin, L.J. and Montano, M.M. 2005. Differential induction of quinone reductase by phytoestrogens and protection against oestrogen-induced DNA damage. *Biochem. J.*, 385(Pt 1): 279-287.
- Bock, K.W., Josting, D., Lilienblum, W. and Pfeil, H. 1979. Purification of rat-liver microsomal UDP-glucuronyltransferase. Separation of two enzyme forms inducible by 3-methylcholanthrene or phenobarbital. *Eur. J. Biochem.*, 98(1): 19-26.

- Bonassi, S., Znaor, A., Ceppi, M., Lando, C., Chang, W.P., Holland, N., Kirch-Volders, M., Zeiger, E., Ban, S., Barale, R., Bigatti, M.P., Bolognesi, C., Cebulska-Wasilewska, A., Fabianiva, E., Fucic, A., Hagmar, L., Jokesic, G., Martelli, A., Migliore, L., Mirkova, E., Scarfi, M.R., Zijno, A., Norppa, H. and Fenech, M. 2007. An increased micronucleus frequency in peripheral blood lymphocytes predicts the risk of cancer in humans. *Carcinogenesis*, 28(3): 625-631.
- Brambilla, G. and Martelli, A. 2004. Failure of the standard batter of short-term tests in detecting some rodent and human genotoxic carcinogens. *Toxicology*, 196: 1-19.
- Brankova, L., Ivanov, S., and Alexieva, V. 2007. The induction of microsomal NADPH: cytochrome P450 and NADH:cytochrome b_5 reductases by long-term salt treatment of cotton (*Gossypium hirsutum* L.) and bean (*Phaseolus vulgaris* L.) plants. *Plant Physiol. Bioch.*, 45 (9): 691-695.
- Charoensin, S., Punvittayagul, C., Pompimon, W. Mevatee, U. and Wongpoomchai, R. Toxicological and clastogenic evaluation of some flavanones isolated from *Boesenbergia pandurata* (Roxb.) in Wistar rats. *Thai. J. Toxicol.* (inpress).
- Charoensin, S. Inhibitory mechanism of pinostrobin isolated from *Boesenbergia pandurata* on diethylnitrosamine-induced initiation stage of rat hepatocarcinogenesis. Master thesis, Department of Biochemistry, Graduate School, Chiang Mai University, 2008.
- Chen, C, and Kong, A.N. 2004. Dietary chemopreventive compounds and ARE/EpRE signaling. *Free Radic. Biol. Med.*, 36(12): 1505-1516.
- Chen, C. and Kong, A.N. 2005. Dietary cancer-chemopreventive compounds: from signaling and gene expression to pharmacological effects. *Trends Pharmacol. Sci.*, 26(6): 318-326.
- Chen, P.N., Hsieh, Y.S., Chiou, H.L. and Chu, S.C. 2005. Silibinin inhibits cell invasion through inactivation of both PI3K-Akt and MAPK signaling pathways. *Chem. Biol. Interact.*, 156: 141-150.
- Chen, P.N., Hsieh, Y.S., Chiang, C.L., Chiou, H.L., Yang, S.F. and Chu, S.C. 2006. Silibinin inhibits invasion of oral cancer cells by suppressing the MAPK pathway, *J. Den. Res.*, 85: 220-225.

- Combes, R.D. 2000. The use of structure-activity relationships and markers of cell toxicity to detect non-genotoxic carcinogens. *Toxicol. In Vitro*, 14(4): 387-99.
- Cook, N.C., Samman, S. 1996. Flavonoids-Chemistry, metabolism, cardioprotective effects, and dietary sources. *Nutr. Biochem.*, 7: 66-76.
- Cullen, J.J., Hinkhouse, M.M., Grady, M., Gaut, A.W., Liu, J., Zhang, Y.P., Barby Weydert, C.J., Domann, F.E. and Oberley, L.W. 2003. Dicumarol inhibition of NADPH:quinone oxidoreductase induces growth inhibition of pancreatic cancer via a superoxide-mediated mechanism. *Cancer Res.*, 63(17): 5513-5520.
- Dasgupta T, Rao RA, Yadava KP. Chemomodulatory action of curry leaf (*Murraya koenigii*) extract on hepatic and extrahepatic xenobiotic metabolising enzymes, antioxidant levels, lipid peroxidation, skin and forestomach papillomagenesis. *Nutr Res* 2003; 23: 1427–1446.
- Debersac, P., Vernevaut, M.F., Amiot, M.J., Suschetet, M., Siess, M.H. 2001. Effects of a water-soluble extract of rosemary and its purified component rosmarinic acid on xenobiotic-metabolizing enzymes in rat liver. *Food Chem. Toxicol.*, 39(2): 109-117.
- Del Rayo Camacho, M., Sanchez, B., Quiroz, H., Contreras, J.L. and Mata, R. 1991. Pinoцембрin: a bioactive flavanone from *Teloxys graveolens*. *J. Ethnopharmacol.*, 31: 383–389.
- Doi, K., Wei, M., Kitano, M., Uematsu, N., Inoue, M. and Wanibuchi, H. 2009. Enhancement of preneoplastic lesion yield by Chios Mastic Gum in a rat liver medium-term carcinogenesis bioassay. *Toxicol. Appl. Pharmacol.*, 234(1): 135-142.
- Doostdar, H., Burke, M.D. and Mayer, R.T. 2000. Bioflavonoids: selective substrates and inhibitors for cytochrome P450 CYP1A and CYP1B1. *Toxicology*, 144(1-3): 31-38.
- Edwards, D.J. and Bernier, S.M. 1996. Inhibitory effect of grapefruit juice and its bitter principal, naringenin, on CYP1A2 dependent metabolism of caffeine in man. *Life Sciences*, 59 (13): 1025–1030.

- Ellinger-Ziegelbauer, H., Stuart, B., Wahle, B., Bomann, W., Ahr, H.J. 2005. Comparison of the expression profiles induced by genotoxic and nongenotoxic carcinogens in rat liver. *Mutat. Res.*, 575(1-2): 61-84.
- Erlund, I. 2004. Review of the flavonoids quercetin, hesperetin, and naringenin. Dietary sources, bioactivities, bioavailability, and epidemiology. *Nutr. Res.*, 24: 851-874.
- Farombi, E.O. 2004. Diet-related cancer and prevention using anticarcinogens. *Afr. J. Biotechnol.*, 3 (12): 651-661.
- Fatemi, F., Allameh, A., Dadkhah, A., Forouzandeh, M., Kazemnejad, S. and Sharifi, R. 2006. Changes in hepatic cytosolic glutathione -S-transferase activity and expression of its class-P during prenatal and postnatal period in rats treated with aflatoxin B1. *Arch. Toxicol.*, 80(9): 572-579.
- Fukushima, S., Morimura, K., Wanibuchi, H., Kinoshita, A. and Salim, E.I. 2005. Current and emerging challenges in toxicopathology: carcinogenic threshold of phenobarbital and proof of arsenic carcinogenicity using rat medium-term bioassays for carcinogens. *Toxicol. Appl. Pharmacol.*, 207(2 Suppl): 225-229.
- Galati, G. and O'Brien, P.J. 2004. Potential toxicity of flavonoids and other dietary phenolics: significance for their chemopreventive and anticancer properties. *Free Radic. Biol. Med.*, 37 (3): 287 - 303.
- Ghia, M., Mattioli, F. and Mereto, E. 1996. A possible medium-term assay for detecting the effects of liver and colon carcinogens in rats. *Cancer Lett.* 105(1): 71-5.
- Gish, R.G. 2006. Hepatocellular Carcinoma: Overcoming Challenges in Disease Management. *Clin. Gastroenterol. H.*, 4: 252-261.
- Gordon, G.G.; Skett, P. *Introduction to drug metabolism*, 2nd ed.; Stanley Thornes Publishers, Ltd: English, 1994, pp. 236-237.
- Grancharov, K., Naydenova, Z., Lozeva, S. and Golovinsky, E. 2001. Natural and synthetic inhibitors of UDP-glucuronosyltransferase. *Pharmacol. Ther.*, 89(2): 171-186.

- Guyton, K.Z., Kyle, A.D., Aubrecht, J., Cogliano, V.J., Eastmond, D.A., Jackson, M., Keshava, N., Sandy, M.S., Sonawane, B., Zhang, L., Waters, M.D. and Smith, M.T. 2009. Improving prediction of chemical carcinogenicity by considering multiple mechanisms and applying toxicogenomic approaches. *Mutat. Res.*, 681(2-3): 230-240.
- Habig, W. H. and Jakoby, W. B., 1981. Assays for differentiation of glutathione-S-Transferases. *Methods Enzymol.*, 77: 398-405.
- Hail, N.Jr., Cortes, M., Drake, E.N., Spallholz, J.E. 2008. Cancer chemoprevention: a radical perspective. *Free Radic. Biol. Med.*, 45(2): 97-110.
- Hakura, A., Kadoi, M., Suzuki, T. and Saeki, K. 2007. Clastogenicity of quinoline derivatives in the liver micronucleus assay using rats and mice. *J. Health. Sci.*, 53(4): 470-474.
- Hannemann, F., Bichet, A., Ewen, K.M., Bernhardt, R. 2007. Cytochrome P450 systems—biological variations of electron transport chains. *Biochim. Biophys. Acta.*, 1770: 330–344.
- Havsteen, B.H. 2002. The biochemistry and medical significance of the flavonoids. *Pharmacol. Ther.*, 96(2-3): 67-202.
- Hawighorst, T., Velasco, P., Streit, M., Hong, Y.K., Kyriakides, T.R., Brown, L.F., Bornstein, P. and Detmar, M. 2001. Thrombospondin-2 plays a protective role in multistep carcinogenesis: a novel host anti-tumor defense mechanism. *EMBO J.*, 20(11): 2631-2640.
- Henderson, C.J., Sahraoui, A. and Wolf, C.R. 2000. Cytochrome P450s and chemoprevention. *Biochem. Soc. Trans.*, 28(2): 42-46.
- Hernandez, L.G., Steeg, H., Luijten, M. and Benthem, J. 2009. Mechanisms of non-genotoxic carcinogens and importance of a weight of evidence approach. *Mutat. Res.*, 682(2-3): 94-109.
- Higashi, K., Hiai, H., Higashi, T. and Muramatsu, M. 2004. Regulatory mechanism of glutathione S-transferase P-form during chemical hepatocarcinogenesis: old wine in a new bottle. *Cancer Lett.*, 209: 155–163.

- Higgin, G.M. and Anderson, R.M., 1931. Experimental pathology of the liver, Restoration of the liver of white rat following partial surgical removal. *Arch. Pathol.*, 12: 186-202.
- Hill, M., Pereira, V., Chauveau, C., Zagani, R., Remy, S., Tesson, L., Mazal, D., Ubillos, L., Brion, R., Ashgar, K., Mashreghi, M.F., Kotsch, K., Moffett, J., Doebis, C., Seifert, M., Boczkowski, J., Osinaga, E. and Anegon, I. 2005. Heme oxygenase-1 inhibits rat and human breast cancer cell proliferation: mutual cross inhibition with indoleamine 2, 3-dioxygenase. *FASEB J.*, 19(14): 1957-1968.
- Hodek, P., Trefil, P. and Stiborova, M. 2002. Flavonoids-potent and versatile biologically active compounds interacting with cytochromes P450. *Chem-Biol. Interact.*, 139: 1-21.
- Hudson, C.E., Kelly, M.M., Schwartz, D.A., Schofield, D.A., Dehaven, J.E., Schulte, B. A. and Norris, J.S. 1998. Glutathione S-transferase in hormonal carcinogenesis. *Chem-Biol. Interact.*, 111-112: 343-50.
- Hwang, E.I., Kaneko, M., Ohnishi, Y. and Horinouchi, S. 2003. Production of plant-specific flavanones by *Escherichia coli* containing an artificial gene cluster. *Appl. Environ. Microbiol.*, 69: 2699-2706.
- Iarmarcovai, G., Bonassi, S. and Botta, A., Baan, R.A. and Orsiere, T. 2008. Genetic polymorphisms and micronucleus formation: a review of the literature. *Mutat. Res.*, 658(3): 215-233.
- Ichihara, T., Wanibuchi, H., Lee, C.C., Nakajima, K., Yano, Y., Taniyama, T., Otani, S., Shimizu, Y. and Fukushima, S. 1999. Lack of inhibitory effects of the Ju-myo protein on development of glutathione-S-transferase placental form-positive foci in the male F344 rat liver. *J Toxicol. Sci.*, 24: 27-31.
- Igarashi, M. and Shimada, H. 1997. An improved method for the mouse liver micronucleus test. *Mutat. Res.*, 391(1-2): 49-55.
- Imaida, K. and Fukushima, S. 1996. Initiation-promotion model for assessment of carcinogenicity: medium-term liver bioassay in rats for rapid detection of carcinogenic agents. *J. Toxicol. Sci.*, 21: 483-487.

- Iqbal, M. and Okada, S. 2003. Induction of NAD(P)H:quinone reductase by probucol: A possible mechanism for protection against chemical carcinogenesis and toxicity. *Pharmacol. Toxicol.*, 93: 259–263.
- Ito, N., Hasegawa, R., Imaida, K., Hirose, M., Asamoto, M. and Shirai, T. 1995. Concepts in multistage carcinogenesis. *Crit. Rev. Oncol. Hematol.*, 21(1-3): 105-133.
- Ito, N., Hasegawa, R., Imaida, K., Hirose, M. and Shirai, T. 1996. Medium-term liver and multi-organ carcinogenesis and chemopreventive agents. *Exp. Toxicol. Pathol.*, 48: 113–119.
- Ito, N., Tamano, S. and Shirai, T. 2003. A medium-term rat liver bioassay for rapid in vivo detection of carcinogenic potential of chemicals. *Cancer Sci.*, 94(1): 3-8.
- Iyanagi, T. 2005. Structure and function of NADPH-cytochrome P450 reductase and nitric oxide synthase reductase domain. *Biochem. Biophys. Res. Commun.*, 338(1): 520-528.
- Jagan, S., Ramakrishnan, G., Anandakumar, P., Kamaraj, S. and Devaki, T. 2008. Antiproliferative potential of gallic acid against diethylnitrosamine-induced rat hepatocellular carcinoma. *Mol. Cell Biochem.*, 319(1-2): 51-59.
- Jaipetch, T., Kanghae, S., Pancharoen, O., Patrick, V.A., Reutrakul, V., Tuntiwachwuttikul, P., and White, A.H. 1982. Constituents of *Boesenbergia pandurata* (syn. *Kaempferia pandurata*): Isolation, crystal structure and synthesis of (±)-Boesenbergin A. *Aust. J. Chem.*, 35: 351-361.
- Jena, B.G., Kaul, L.C., and Ramarao, P. 2005. Regulatory requirements and ICH guidelines on carcinogenicity testing of pharmaceuticals: A review on current status. *Indian J. Pharmacol.*, 37(4): 209-222.
- Kaferstein, H. 2009. Forensic relevance of glucuronidation in phase-II-metabolism of alcohols and drugs. *Leg. Med. (Tokyo)*, 11 Suppl 1: S22-6.
- Kikuchi, G., Yoshida, T. and Noguchi, M. 2005. Heme oxygenase and heme degradation. *Biochem. Biophys. Res. Commun.*, 338(1): 558-567.

- Kirkland, D., Aardema, M., Muller, L. and Hayashi, M. 2006. Evaluation of the ability of a battery of three in vitro genotoxicity tests to discriminate rodent carcinogens and non-carcinogens II. Further analysis of mammalian cell results, relative predictivity and tumour profiles. *Mutat. Res.*, 608(1): 29-42.
- Klaassen, C. D., *Casarett and Doull's: Toxicology - The Basic Science of Poisons*, 7th ed; The McGraw-Hill Companies, Inc.: USA, 2008, pp. 131-159.
- Klaunig, J.E. and Kamendulis, L.M. 2004. The role of oxidative stress in carcinogenesis. *Annu. Rev. Pharmacol. Toxicol.*, 44: 239-267.
- Klaunig, J.E., Xu, Y., Isenberg J.S., Bachowski, S., Kolaja, K.L., Jiang, J., Stevenson, D. E. and Walborg Jr, E.F. 1998. The role of oxidative stress in chemical carcinogenesis. *Environ. Health Perspect.*, 106 Suppl 1: 289-295.
- Klein, C.B. and Costa, M. 1997. DNA methylation, heterochromatin and epigenetic carcinogens. *Mutat. Res.*, 386(2): 163-180.
- Ko, C.H., Shen, S.C., Lin, H.Y., Hou, W.C., Lee, W.R., Yang, L.L. and Chen, Y.C. 2002. Flavanones structure-related inhibition on TPA-induced tumor promotion through suppression of extracellular signal-regulated protein kinases: involvement of prostaglandin E2 in anti-promotive process, *J. Cell Physiol.*, 193: 93-102.
- Kohno, H., Taima, M., Sumida, T., Azuma, Y., Ogawa, H. and Tanaka, T. 2001. Inhibitory effect of mandarin juice rich in beta-cryptoxanthin and hesperidin on 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced pulmonary tumorigenesis in mice. *Cancer Lett.*, 174: 141-150.
- Kushida, M., Sukata, T., Uwagawa, S., Ozaki, K., Kinoshita, A., Wanibuchi, H., Morimura, K., Okuno, Y. and Fukushima, S. 2005. Low dose DDT inhibition of hepatocarcinogenesis initiated by diethylnitrosamine in male rats: possible mechanisms. *Toxicol. Appl. Pharmacol.*, 208(3): 285-294.
- Kwon, K.H., Barve, A., Yu, S., Huang, M. and Kong, A.N. 2007. Cancer chemoprevention by phytochemicals: potential molecular targets, biomarkers and animal models. *Acta Pharmacol. Sin.*, 28(9): 1409-1421.
- Lee, B.M. and Park, K.K. 2003. Beneficial and adverse effects of chemopreventive agents. *Mutat. Res.*, 523-524: 265-278.

- Letelier, M.E., Pimentel, A., Pino, P., Lepe, A.M., Faundez, M., Aracena, P. and Speisky, H. 2005. Microsomal UDP-glucuronyltransferase in rat liver: oxidative activation. *Basic Clin. Pharmacol. Toxicol.*, 96(6): 480-486.
- Li Ching, A. Y., Sook Wah, T. , Sukari, M. A., Cheng Lian G. E., Rahmani M., and Khalid, K., 2007. Characterization of flavonoid derivatives from *Boesenbergia rotunda* (L.). *The Malaysian Journal of Analytical Sciences*, 11(1): 154-159.
- Liska, D.J. 1998. The detoxification enzyme systems. *Altern. Med. Rev.*, 3(3): 187-98.
- Liu, R., Gao, M., Yang, Z.H., Du, G-H. 2008. Pinocembrin protects rat brain against oxidation and apoptosis induced by ischemia-reperfusion both in vivo and in vitro. *Brain Res.*, 1216: 104-115.
- Liu, C., Shang, Y.F., Zhang, X.F., Zhang, X.G., Wang, B., Wu, Z., Liu, X.M., Yu, L., Ma, F. and Lv, Y. 2009. Co-administration of grapefruit juice increases bioavailability of tacrolimus in liver transplant patients: a prospective study. *Eur. J. Clin. Pharmacol.*, 65(9):881-885.
- Lowry, O. H., Rosebrough, N. J., Farr, A.L. and Randall, R. J., 1951. Protein measurement with the Folin-Phenol reagents. *J. Biol. Chem.*, 193: 265-275.
- Luch, A. 2005. Nature and nurture - lessons from chemical carcinogenesis. *Nat. Rev. Cancer*, 5(2): 113-125.
- Maines, M.D. and Gibbs, P.E. 2005. 30 some years of heme oxygenase: from a "molecular wrecking ball" to a "mesmerizing" trigger of cellular events. *Biochem. Biophys. Res. Commun.*, 338(1): 568-577.
- Majewska, M., Zajac, K., Dulak, J. and Szczepanik, M. 2008. Heme oxygenase (HO-1) is involved in the negative regulation of contact sensitivity reaction. *Pharmacol. Rep.*, 60(6): 933-940.
- Mally, A. and Chipman, J.K. 2002. Non-genotoxic carcinogens: early effects on gap junctions, cell proliferation and apoptosis in the rat. *Toxicology*, 180(3): 233-248.
- Manson, M.M., Gescher, A., Hudson E.A., Plummer, S.M., Squires, M.S. and Prigent, S. A. 2000. Blocking and suppressing mechanisms of chemoprevention by dietary constituents. *Toxicol. Lett.*, 112-113: 499-505.

- Manthey, J.A., Grohmann, K. and Guthrie, N. 2001. Biological properties of citrus flavonoids pertaining to cancer and inflammation. *Curr. Med. Chem.*, 8: 135–153.
- Matsuda, M., Nakamoto, Y., Suzuki, S., Kurata, T. and Kaneko, S. 2005. Interferon-gamma-mediated hepatocarcinogenesis in mice treated with diethylnitrosamine. *Lab Invest.*, 85(5): 655-663.
- Mazzanti, R., Gramantieri, L. and Bolondi, L. 2008. Hepatocellular carcinoma: epidemiology and clinical aspects. *Mol. Aspects. Med.*, 29(1-2): 130-143.
- McKinney, M. A., Arukwe, A., Guise, S., Martineau, D., Beland, P., Dallaire, A., Lebeuf, M. and Letcher, R.J. 2004. Characterization and profiling of hepatic cytochromes P450 and phase II xenobiotic-metabolizing enzymes in beluga whales (*Delphinapterus leucas*) from the St. Lawrence River Estuary and the Canadian Arctic. *Aquat. Toxicol.*, 69(1): 35-49.
- McNally, S. J., Ross, J. A., Garden, O. J., and Wigmore, S. J., 2004. Optimization of the paired enzyme assay for heme oxygenase activity. *Anal. Biochem.*, 332, 398–400.
- Misra, V., Grondin, A., Klamut, H.J. and Rauth, A.M. 2000. Assessment of the relationship between genotypic status of a DT-diaphorase point mutation and enzymatic activity. *Br. J. Cancer*, 83(8): 998-1002.
- Moon, Y.J., Wang, X., Morris, M.E., 2006. Dietary flavonoids: effects on xenobiotic and carcinogen metabolism. *Toxicol. In Vitro*, 20(2): 187-210.
- Morse, M.A. and Stoner, G.D. 1993. Cancer chemoprevention: principles and prospects. *Carcinogenesis*, 14(9): 1737-1746.
- Moulari, B., Pellequer, Y., Lboutounne, H., Girard, C., Chaumont, J.P., Millet, J. and Muyard, F. 2006. Isolation and in vitro antibacterial activity of astilbin, the bioactive flavanone from the leaves of *Harungana madagascariensis* Lam. ex Poir. (Hypericaceae). *J. Ethnopharmacol.*, 106: 272–278.
- Muller-Tegethoff, K. Kasper, P. and Muller, L. 1995. Evaluation studies on the in vitro rat hepatocyte micronucleus assay. *Mutat. Res.*, 335: 293-307.

- Nair, S., Li, W., Kong, A.N. 2007. Natural dietary anti-cancer chemopreventive compounds: redox-mediated differential signaling mechanisms in cytoprotection of normal cells versus cytotoxicity in tumor cells. *Acta Pharmacol. Sin.*, 28(4): 459-472.
- Nebbia, C. 2001. Biotransformation enzymes as determinants of xenobiotic toxicity in domestic animals. *Vet. J.*, 161(3): 238-252.
- Nikolic, D. and Breemen, R.B. 2003. New metabolic pathways for flavanones catalyzed by rat liver microsomes. *Drug. Metab. Dispos.*, 32(4): 387-397.
- Nioi, P. and Hayes, J.D. 2004. Contribution of NAD(P)H:quinone oxidoreductase 1 to protection against carcinogenesis, and regulation of its gene by the Nrf2 basic-region leucine zipper and the arylhydrocarbon receptor basic helix-loop-helix transcription factors. *Mutat. Res.*, 555(1-2): 149-171.
- Ogiso, T., Tatematsu, M., Tamano, S., Hasegawa, R. and Ito, N. 1990. Correlation between medium-term liver bioassay system data and results of long-term testing in rats. *Carcinogenesis*, 11(4): 561-566.
- Ohkawa, H., Ohishi, N. and Yagi, K., 1979. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.*, 95(2): 351-358.
- Oliveira, P.A., Colaco, A., Chaves, R., Guedes-Pinto, H., De-La-Cruz, Luis F. and Lopes, C. 2007. Chemical carcinogenesis. *An. Acad. Bras. Cienc.*, 79(4): 593-616.
- Ortiz de Montellano, P.R., 2000. The mechanism of heme oxygenase. *Curr. Opin. Chem. Biol.*, 4: 221-227
- Pepeljnjak, S., Jalsenjak, I. and Maysinger, D. 1985. Flavonoid content in propolis extracts and growth inhibition of *Bacillus subtilis*. *Pharmazie*, 40: 122-123.
- Prawan, A., Kundu, J.K. and Surh, Y.J., 2005. Molecular basis of heme oxygenase-1 induction: implications for chemoprevention and chemoprotection. *Antioxid. Redox. Signal.*, 7: 1688-1703.
- Puatanachokchai, R., Noguchi, T., Vinitketkumnuen, U. and Matsushima, T. Rat liver micronucleus assay. Proceeding of the 4th Southeast Asian Workshop on Short-term assays for detection of environmental mutagens, carcinogens and teratogens. Naresuan University, Phitsanulok, Thailand, December 16th- 20th, 1996: 14-23.

- Puatanachokchai, R., Morimura, K., Wanibuchi, H., Oka, M., Kinoshita, A., Mitsuru, F., Yamaguchi, S., Funae, Y., and Fukushima, S., 2006. Alpha-benzene hexachloride exerts hormesis in preneoplastic lesion formation of rat hepatocarcinogenesis with the possible role for hepatic detoxifying enzymes. *Cancer Lett.*, 18(240;1): 102-113.
- Rundle, A. 2006. Carcinogen-DNA adducts as a biomarker for cancer risk. *Mutat. Res.*, 600(1-2): 23-36.
- Sadik, N., El-Maraghy, S.A. and Ismail, M.F. 2008. Diethylnitrosamine-induced hepatocarcinogenesis in rats: possible chemoprevention by blueberries. *Afr. J. Biochem. Res.*, 2(3): 81-87.
- Sakai, M. and Muramatsu, M. 2007. Regulation of glutathione transferase P: A tumor marker of hepatocarcinogenesis. *Biochem. Bioph. Res. Co.*, 357: 575-578.
- Sala, A., Recio, M.C., Schinella, G.R., Máñez, S., Giner, R.M., Cerdá-Nicolás, M. and Rosí, J.L. 2003. Assessment of the anti-inflammatory activity and free radical scavenger activity of tiliroside. *Eur. J. Pharmacol.*, 461: 53-61.
- Saracino, M.R. and Lampe, J.W. 2007. Phytochemical regulation of UDP-glucuronosyltransferases: implications for cancer prevention. *Nutr. Cancer*, 59(2): 121-41.
- Sasagawa, T., Kosai, K., Ota, Y., Mori, M. and Okita, M. 2002. Influences of a dietary fatty acid composition on the emergence of glutathione S-transferase-P (GST-P) positive foci in the liver of carcinogen-treated rats. *Prostag. Leukotr. Ess.*, 67(5): 327-332.
- Santos, A.C., Uyemura, S.A., Lopes, J.L., Bazon, J.N., Mingatto, F.E. and Curti, C. 1998. Effect of naturally occurring flavonoids on lipid peroxidation and membrane permeability transition in mitochondria. *Free Radic. Biol. Med.*, 24: 1455-1461.
- Schafer, D.F. and Sorrell, M.F. 1999. Hepatocellular carcinoma. *Lancet*, 353: 1253-57.
- Shindo, K., Kato, M., Kinoshita, A., Kobayashi, A. and Koike, Y. 2006. Analysis of antioxidant activities contained in the *Boesenbergia pandurata* Schult. rhizome. *Biosci. Biotechnol. and Biochem.*, 70: 2281-2284.

- Shirai, T. 1997. A medium-term rat liver bioassay as a rapid in vivo test for carcinogenic potential: A historical review of model development and summary of results from 291 tests. *Toxicol. Pathol.*, 25(5): 453-460.
- Siess, H.M., Leclerc, J., Canivenc-Lavire, M., Rat, P. and Suschetet, M. 1995. Heterogenous effect of natural flavonoids on monooxygenase activities in human and rat liver microsomes. *Toxicol. Appl. Pharmacol.*, 130: 73-78.
- Silva, R.N., Ribeiro, D.A., Salvadori, D.M. and Marques, M.E. 2007. Placental glutathione S-transferase correlates with cellular proliferation during rat tongue carcinogenesis induced by 4-nitroquinoline 1-oxide. *Exp. Toxicol. Pathol.*, 59(1): 61-68.
- Srivatanakul, P. 2001. Epidemiology of liver cancer in Thailand. *Asian Pac. J. Cancer Prev.*, 2(2): 117-121.
- Surh, Y-J., Kundu, J. K. and Na, H-K. 2008. Nrf2 as a master redox switch in turning on the cellular signaling involved in the induction of cytoprotective genes by some chemopreventive phytochemicals. *Planta Med.*, 74: 1526-1539.
- Suzuki, H., Shirotori, T. and Hayashi, M. 2004. A liver micronucleus assay using young rats exposed to diethylnitrosamine: methodological establishment and evaluation. *Cytogenet. Genome Res.*, 104(1-4): 299-303.
- Tan, X.L. and Spivack, S.D. 2009. Dietary chemoprevention strategies for induction of phase II xenobiotic-metabolizing enzymes in lung carcinogenesis: A review. *Lung Cancer*, 65(2): 129-137.
- Tanaka, T., Makita, H., Kawabata, K., Mori, H., Kakumoto, M., Satoh, K., Hara, A., Sumida, T., Fukutani, K. and Ogawa, H. 1997. Modulation of N-methyl-N-amylnitrosamine-induced rat oesophageal tumourigenesis by dietary feeding of diosmin and hesperidin, both alone and in combination. *Carcinogenesis*, 18: 761-769.
- Townsend, D.M. and Tew, K.D. 2003. The role of glutathione-S-transferase in anti-cancer drug resistance. *Oncogene*, 22(47): 7369-7375.

- Trakoontivakorn, G., Nakahara, K., Shinmoto, H., Takenaka, M., Onishi-Kameyama, M., Ono, H., Yoshida, M., Nagata, T. and Tsushida, T. 2001. Structural analysis of a novel antimutagenic compound, 4-hydroxypanduratin A, and the antimutagenic activity of flavonoids in a Thai spice, fingerroot (*Boesenbergia pandurata* Schult.) against mutagenic heterocyclic amines. *J. Agric. Food Chem.*, 49: 3046-3050.
- Tripoli, E. La Guardia, M., Giammanco, S., Di Majo, D. and Giammanco M. 2007. Citrus flavonoids: Molecular structure, biological activity and nutritional properties: A review. *Food Chem.*, 104: 466–479.
- Tsuda, H., Fukushima, S., Wanibuchi, H., Morimura, K., Nakae, D., Imaida, K., Tatematsu, M., Hirose, M., Wakabayashi, K. and Moore, M.A. 2003. Value of GST-P positive preneoplastic hepatic foci in dose-response studies of hepatocarcinogenesis: evidence for practical thresholds with both genotoxic and nongenotoxic carcinogens. A review of recent work. *Toxicol. Pathol.*, 31(1): 80-86.
- Tsuda, H., Ohshima, Y., Nomoto, H., Fujita, K., Matsuda, E., Iigo, M., Takasuka, N. and Moore, M.A. 2004. Cancer prevention by natural compounds. *Drug Metab. Pharmacokinet.*, 19(4): 245-263.
- Tuchinda, P., Reutrakul, V., Claesona, P., Pongprayoon, U., Sematong, T., Santisuk, T. and Taylor, W.C. 2002. Anti-inflammatory cyclohexenyl chalcone derivatives in *Boesenbergia pandurata*. *Phytochemistry*, 59: 169–173.
- Verna, L., Whysner, J. and Williams, G. 1996. N-nitrosodiethylamine mechanistic data and risk assessment: bioactivation, DNA-adduct formation, mutagenicity, and tumor initiation. *Pharmacol. Ther.*, 71(1-2): 57-81.
- Walle, T., Ta, N., Kawamori, T., Wen, X., Tsuji, P. A. and Walle U. K., 2007. Cancer chemopreventive properties of orally bioavailable flavonoids—Methylated versus unmethylated flavones. *Biochem. Pharmacol.*, 73: 1288-1296.
- Wen, X. and Walle, T. 2006. Methylated flavonoids have greatly improved intestinal absorption and metabolic stability. *Drug. Metab. Dispos.*, 34(1):1786–1792.
- Wilce, M.C. and Parker, M.W. 1994. Structure and function of glutathione S-transferases. *Biochim. Biophys. Acta.*, 1205(1): 1-18.

- Williams, G.M. 2001. Mechanisms of chemical carcinogenesis and application to human cancer risk assessment. *Toxicology*, 166(1-2): 3-10.
- Wu, L., Tang, Z.Y. and Li, Y. 2009. Experimental models of hepatocellular carcinoma: developments and evolution. *J. Cancer Res. Clin. Oncol.*, 135(8): 969-981.
- Xie, W., Uppal, H., Saini, S.P., Mu, Y., Little, J.M., Radominska-Pandya, A. and Zemaitis, M.A. 2004. Orphan nuclear receptor-mediated xenobiotic regulation in drug metabolism. *Drug Discov. Today*, 9(10): 442-449.
- Yang, M., Tanaka, T., Hirose, Y., Deguchi, T., Mori, H. and Kawada, Y. 1997. Chemopreventive effects of diosmin and hesperidin on *N*-butyl-*N*-(4-hydroxybutyl) nitrosamine-induced urinary-bladder carcinogenesis in male ICR mice. *Int. J. Cancer*, 73: 719-724.
- Yang, Z., Liu, R., Li, X., Tian, S., Liu, Q. and Du, G. 2009. Development and validation of a high-performance liquid chromatographic method for determination of pinocembrin in rat plasma: Application to pharmacokinetic study. *J. Pharmaceut. Biomed.*, 49: 1277-1281.
- Yates, M.S. and Kensler, T.W. 2007. Keap1 eye on the target: chemoprevention of liver cancer. *Acta Pharmacol. Sin.*, 28(9): 1331-1342.
- Yim, S.K., Yun, C.H., Ahn, T., Jung, H.C., and Pan, J.G. 2005. A continuous spectrophotometric assay for NADPH-cytochrome P450 reductase activity using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide. *J. Biochem. Mol. Biol.*, 38(3): 366-369.

APPENDICES

APPENDIX A

List of the chemicals and materials used in the study

Chemicals/Materials	Source
β - NADPH	Oriental Yeast, Japan
1-Chloro-2, 4-dinitrobenzene	Fluka A.G., Switzerland
2-aminophenol	Sigma-Aldrich, USA
4', 6-diamidino-2-phenylindole dihydrochloride	Invitrogen, USA
5, 5'-Dithio-bis (2-Nitrobenzoic acid)	Sigma-Aldrich, USA
Absolute ethanol	BDH, England
Acrylamide	Biorad, USA
Ammonium persulphate	Carlo-Erba, Italy
Ammonium sulfamate	Sigma-Aldrich, USA
2-Aniline	BDH, England
Ascorbic acid	Merck A.G., Germany
Bis-acrylamide	Biorad, USA
Bovine serum albumin	Sigma-Aldrich, USA
Butanol	BDH, England
Calcium chloride dihydrate	Merck A.G., Germany
Chloroform	LAB-SCAN, Thailand
Collagenase type IV	Invitrogen, USA
Copper sulphate	Merck A.G., Germany
Cytochrome c Type VI	Sigma-Aldrich, USA
Diethyl ether	BDH, England
Disodium hydrogen orthophosphate	BDH, England
Ethylene diamine tetraacetic acid	Sigma-Aldrich, USA

Ethylene glycol tetraacetic acid	Sigma-Aldrich, USA
Folin & Ciocalteu's phenol reagent	Fluka A.G., Switzerland
Formalin	BDH, England
Glucose-6-phosphate	Sigma-Aldrich, USA
Glucose-6-phosphate dehydrogenase	Sigma-Aldrich, USA
Glutathione (Reduced form)	Wako, Japan
Glycerol	Sigma-Aldrich, USA
Glycine	Vivantis, Malaysia
Hemin	Sigma-Aldrich, USA
HEPES, free acid	Amresco, USA
Hydrochloric	BDH, England
Isopropanol	Sigma-Aldrich, USA
Magnesium chloride	APS Finechem, Australia
Malondialdehyde bis(dimethyl acetal)	Sigma-Aldrich, USA
Mercaptoethanol	Sigma-Aldrich, USA
Methanol	BDH, England
N-naphthylethylene diamine	BDH, England
Phenol red sodium salt	Amresco, USA
Phenylmethylsulphonyl fluoride	Sigma-Aldrich, USA
Potassium chloride	Carlo-Erba, Italy
Potassium cyanide	Merck A.G., Germany
Potassium Dihydrogen Phosphate	May and Baker, England
Potassium sodium ttrate	Mallinckrodt chemical work, USA
Skimmed milk	Merck A.G., Germany
Sodium carbonate	BDH, England
Sodium chloride	BDH, England
Sodium dihydrogenphosphate	BDH, England
Sodium dodecyl sulfate	Biorad, USA
Sodium hydrogen carbonate	BDH, England
Sodium hydroxide	BDH, England
Sodium nitrite	AJAX Chemicals, Australia

TEMED	USB, USA
Thiobarbituric acid	Sigma-Aldrich, USA
Trichloroacetic acid	BDH, England
Tris base	Vivantis, Malaysia
Triton x-100	Sigma-Aldrich, USA
Tween-20	USB, USA
Tween-80	BDH, England
UDP-glucuronic acid	Sigma-Aldrich, USA

APPENDIX B

List of the instruments used in the study

Instrument	Model	Source
Analytical balance	300A	Precisa, USA
Blotting apparatus	Trans-blot SD cell	Biorad, USA
Centrifugator	PMC-060	Tomy Seiko, Japan
	22R D-78532	Mikro, Germany
Electrophoretic apparatus	AE-6500	Atto corp., Japan
Film cassette	RPN 11649	Amersham, England
Fluorescent microscope	AX-70	Olympus, Japan
Freezer (-86°C)	0838	Forma Scientific, USA
Homogenizer	HS-30E	Daihan, Korea
Hot plate/stirrer	HPMS	Whatman, USA
Light microscope		Olympus, Japan
Microplate reader	MCC/340	ICN, Flow, USA
Mini rocker		Biosan, Taiwan
pH meter	320	Mettler Toledo, USA
Peristaltic pump	MP-100	Tokyo Rikakikai, Japan

Refrigerator	SR-F511	Sanyo, Thailand
Ultracentrifuge	L-100 XP	Beckman Coulter
UV-Spectrophotometer	UV-1700	Shimazu, Japan
Vortex	G-560E	Scientific industries, USA
Water bath	W-350	Mammert, Germany

APPENDIX C

Reagents and buffers preparation

1. Preparation of mediums for liver-cell suspension

1.1 Preperfusion medium

NaCl	8	g
KCl	0.4	g
KH ₂ PO ₄	0.06	g
Na ₂ HPO ₄	0.09	g
EGTA	0.195	g
HEPES	2.39	g
NaHCO ₃	0.35	g

After dissolve in distilled water, adjust pH to 7.4 with 1 N NaOH and bring to 1L with distilled water.

1.2 Collagenase medium

NaCl	8	g
KCl	0.4	g
KH ₂ PO ₄	0.06	g
Na ₂ HPO ₄	0.09	g
Phenol red	0.01	g
CaCl ₂ ·2H ₂ O	0.56	g
HEPES	2.39	g
NaHCO ₃	0.35	g
Collagenase type IV	0.5	g

After completely dissolve in 1 liter of distilled water, adjust pH to 7.4 with 1 N NaOH

1.3 Phosphate buffer saline (Mg²⁺, Ca²⁺, free)

NaCl	8	g
KCl	0.2	g
KH ₂ PO ₄	0.2	g
Na ₂ HPO ₄	1.15	g

After dissolve in distilled water, adjust pH to 7.4 with 1 N NaOH and fill to 1 liter.

2. Preparation of buffers for microsome fraction*2.1 Homogenizing buffer*

KCl	11.5	g
EDTA	0.37	g

After completely dissolved in deionized water, add 1 ml of 0.25 M PMSF and fill to 1 liter

2.2 Microsome suspension buffer

KH ₂ PO ₄	0.14	g
EDTA	0.004	g
DTT	0.002	g
Glycerol	3	ml

After completely dissolved in deionized water, adjust pH to 7.4 with conc. KOH and fill to 10 mL.

3. Preparation of SDS-PAGE reagent and buffer*3.1 30% Acrylamide*

Acryamide	300	g
Bisacrylamide	8	g

After dissolve in deionized water and fill to 100 mL, store at 4 °C.

3.2 Separating gel buffer pH 8.8

Tris base	19.71	g
SDS	4	g

After completely dissolved in deionized water, adjust pH to 8.8 with conc. HCl and fill to 1 liter.

3.3 Stacking gel buffer pH 6.8

Tris base	60.57	g
SDS	8	g

After completely dissolved in deionized water, adjust pH to 6.8 with 1 N NaOH and fill to 1 liter.

3.4 10% Ammonium persulphate

Dissolve 10 g of ammonium persulphate in deionized water and fill to 100 mL.

3.5 Sample buffer pH 6.8

Tris base	0.15	g
SDS	0.40	g
Glycerol	2	ml
2-mercaptoethanol	1	ml
0.002% bromophenol blue	5	μl

Dissolve in deionized water, adjust pH to 6.8 with conc. HCl and fill to 1 liter.

3.6 Electrode buffer

Tris base	3.03	g
Glycine	4.41	g
SDS	1	g

Dissolve in deionized water and fill to 1 liter.

3.7 Blotting buffer

Tris Base	12.11	g
Glycine	14.4	g

After dissolve in deionized water, mix 200 mL of methanol and bring to 1 liter.

4. Preparation of immunostaining buffer*4.1 Phosphate buffer saline*

NaH ₂ PO ₄ ·H ₂ O	1.56	g
NaCl	9.00	g

Dissolve in deionized water, adjust pH to 7.5 with 1 N NaOH and fill to 1 liter.

4.2 Tween-phosphate buffer saline

NaH ₂ PO ₄ ·H ₂ O	1.56	g
NaCl	9.00	g

Dissolve in deionized water, after completely dissolved, add 0.5 ml tween-20 and adjust pH to 7.5 with 1N NaOH and fill to 1 liter.

4.3 5% skimmed milk

Dissolve 5 g of non-fat dried milk in TPBS and fill to 100 ml.

4.4 10% BSA

Dissolve 10 g of BSA TPBS and fill to 100 ml.

4.5 TPBS-0.2% BSA

Mix 100 ml of TPBS with 2 ml of 10% BSA.

5. Preparation of reagents and buffer for lipid peroxidation assay (Thiobarbituric acid method)

5.1 0.1 M Sodium phosphate buffer pH 7.0

Dissolve 6.2404 g of Na₂HPO₄ in 200 ml of distilled water.

Dissolve 14.3256 g of NaH₂PO₄ in 200 ml of distilled water.

Add NaH₂PO₄ solution into the Na₂HPO₄ solution until pH value reaches 7.0.

5.2 Malondialdehyde dimethyl acetal (MDA) stock solution

MDA	5	μl
Methanol	5	ml
0.2 M HCl	5	ml

All chemicals mix and store the reagent at room temperature (freshly prepare)

5.3 50% Trichloroacetic acid (TCA)

Dissolve 50 g of TCA in 70 ml of distilled water and made up to 100 ml.

5.4 0.67% Thiobarbituric acid (TBA)

Dissolve 0.67 g of TBA in 90 ml of distilled water and made up to 100 ml.

6. Preparation of reagents and buffer for NADPH-P450 reductase activity assay

6.1 0.3M potassium phosphate buffer (pH 7.5)

Dissolve 4.08 g of KH₂PO₄ in deionized water, adjust pH to 7.5 with conc. KOH and made up to 100 ml.

6.2 10 mM NADPH

Dissolve 9.1 mg of NADPH in 1 ml of deionized water.

6.3 1mM Cytochrome c Type VI (Horse Heart Type VI)

Dissolve 12.384 mg of Cytochrome *c* in 1 ml of deionized water.

6.4 50 mM KCN

Dissolve 32.56 mg of KCN in deionized water and made up to 10 ml.

7. Preparation of reagents and buffer for heme oxygenase activity assay

7.1 0.1M potassium phosphate buffer containing MgCl₂, pH 7.4

Dissolve 13.61 g of KH₂PO₄ in deionized water, adjust pH to 7.4 with conc. KOH. Add 0.406 g of MgCl₂ and made up to 1 liter.

7.2 400 μM Hemin

Dissolve 2.6 mg of hemin in dichloromethane 10 ml

7.3 8mM Gucllose-6-phosephate

Dissolve 22.57 mg of Gucllose-6-phosephate in deionized water 10 ml

7.4 6.4 mM β-NADPH

Dissolve 5.33 mg of β-NADPH in deionized water 1 ml

8. Preparation of reagents and buffer for NADPH quinone reductase activity assay

8.1 37.5mM Tris-Hcl buffer, pH 7.4

Dissolve 4.54 g of Tris in deionized water, adjust pH to 7.4 with conc. HCl and made up to 1 liter.

8.2 30 mg/ml BSA

Dissolve 30 mg of BSA in 1 ml of deionized water.

8.3 36 mM DCPIP

Dissolve 104.4 mg of DCPIP in 10 ml of deionized water.

8.4 300 mM β-NADPH

Dissolve 1 g of NADPH in 4 ml of deionized water.

8.5 4.5 mM FAD

Dissolve 37.32 mg of FAD in 10 ml of deionized water.

9. Preparation of reagents and buffer for UDP-glucuronyl transferase activity assay

9.1 0.1 M Tris buffer pH 8.0

Dissolve 12.1 g of Tris in deionized water, adjust pH to 8 with conc. HCl and made up to 100 ml.

9.2 1mM 2-aminophenol

Dissolve 11 mg of 2-aminophenol in 90 ml of distilled water and made up to 100 ml.

9.3 20%(w/v) TCA

Dissolve 200 g of TCA in distilled water and made up to 1 liter.

9.4 6%(w/v) TCA

Dissolve 60 g of TCA in distilled water and made up to 1 liter.

9.5 0.15 M $MgCl_2$

Dissolve 15.25 g of $MgCl_2$ in distilled water and made up to 500 ml.

9.6 1% (w/v) Triton x-100

Dissolve 100 mg of triton x-100 in distilled water and made up to 50 ml.

9.7 0.02 M ascorbic acid

Dissolve 180 mg of ascorbic acid in distilled water and made up to 50 ml.

9.8 20%(w/v) TCA in 0.1 M phosphate buffer pH 2.7

Dissolve 200 g of TCA in 0.1 M phosphate buffer pH 2.7 made up to 1 liter.

9.9 6%(w/v) TCA in 0.1 M phosphate buffer pH 2.7

Dissolve 60 g of TCA in 0.1 M phosphate buffer pH 2.7 made up to 1 liter.

9.10 0.1% (w/v) sodium nitrite

Dissolve 100 mg of sodium nitrite in distilled water and made up to 100 ml (freshly prepare).

9.11 0.5 % (w/v) ammonium sulfamate

Dissolve 500 mg of ammonium sulfamate in distilled water and made up to 100 ml.

9.12 0.1% (w/v) N-naphthlethylene diamine

Dissolve 100 mg of N-naphthlethylene diamine in distilled water and made up to 100 ml.

9.13 0.1 mM aniline in 6%(w/v) TCA

Aniline (HCl salt)	13	mg
TCA	6	g

Dissolve in distilled water and fill to 100 ml.

10. Preparation of reagents and buffer for glutathione-S-transferase activity assay

10.1 0.2 M potassium phosphate buffer pH 6.5

Dissolve 2.7 g of KH_2PO_4 in 80 ml of deionized water, adjust pH to 6.5 with conc. KOH and made up to 100 ml.

10.2 10 mM 1-chloro-2,4-dinitrobenzene

Dissolve 102 mg of 1-chloro-2,4-dinitrobenzene in 20 ml of absolute ethanol. After completely dissolved, total volume was adjusted to 50 ml by deionized water.

10.3 10 mM reduced glutathione

Dissolve 307 mg of reduced glutathione in 10 ml of deionized water.

11. Preparation of reagents and buffers for total protein assay by Lowry method

11.1 2% Na_2CO_3 in 0.1 N NaOH (Reagent A)

Dissolve 10 g of Na_2CO_3 in 400 ml of deionized water. After completely dissolved, add 2 g of NaOH and adjust volume to 500 ml.

11.2 0.5% $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (Reagent B)

Dissolve 1 g of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ in deionized water and made up to 200 ml.

11.3 1% potassium sodium tartrate (Reagent C)

Dissolve 1 g of potassium sodium tartrate in deionized water and made up to 100 ml.

11.4 Carbonate-copper solution (Reagent D)

Mix reagent A, B, and C at ratio 50:1:1.

11.5 Diluted Folin reagent

Dilute Folin-Ciocalteu's phenol reagent in deionized water at equal volume.

APPENDIX D

Extraction and isolation of pinocembrin from fingerroot

1. Extraction and isolation

Figure S-1 presents the procedure of pinocembrin isolated from *Boesenbergia pandurata*. Briefly, chopped-dried rhizomes (1 kg) of *Boesenbergia pandurata* were three times extracted with n-hexane, ethyl acetate at room temperature, followed by filtration. The solvent was evaporated under reduced pressure to afford the n-hexane (29.09 g), ethyl acetate (79.99 g), respectively.

A part of the ethyl acetate extract (150 g) was further subjected to column chromatography on silica gel (400 g) eluting with hexane/ethyl acetate/methanol to yield 6 fractions (F1–F6). Fraction F5 was chromatographed by on silica gel (400 g) eluting with hexane/ methanol to give 5 subfractions (F1–F5). Subfraction F5 (9g) was recrystallized from ethanol to give yellow crystal of pinocembrin (130 mg), Figure S-2.




Figure S-1 Extraction scheme of *Boesenbergia pandurata*




Figure S-2 Isolation scheme of pinocembrin

Structure elucidation of 5, 7 dihydroxyflavanone (Pinocembrin)

UV $\lambda_{\text{max}}^{\text{MeOH}}$ **nm (log ϵ):** 288 (4.10), 323 (3.83).

FTIR $\nu_{\text{max}}^{\text{KBr}}$ **cm⁻¹:** 3415 (O-H stretching), 1630 (C=O stretching of ketone), 1583, 1487, 1467, 1357, 1302, 1089.

¹H-NMR (400 MHz, CDCl₃): δ 12.13 (*s*, 1H, 5-OH), 10.85 (*br s*, 1H, 7-OH), 7.41 (1H, *m*, 2',3',4',5',6'), 5.93 (*d*, 1H, *J*= 2.1 Hz, H-8), 5.90 (*d*, 1H, *J*= 2.1 Hz, H-6) 5.58 (*dd*, 1H, *J*= 13, 3 Hz, H-2), 3.23(*dd*, 1H, *J*= 18, 13 Hz, H-3a), 2.72(*dd*, 1H, *J*= 18, 3 Hz, H-3a)

¹³C-NMR (100 MHz, CDCl₃): δ 195.81 (C-4), 166.63 (C-7), 163.42 (C-5), 162.63 (C-9), 138.61 (C-1'), 128.44(C-3'-5'), 126.49 (C-2',6'), 101.71 (C-10), 95.85 (C-6), 78.29 (C-2), 42.07(C-3).

EIMS m/z (% relative intensity): 256 (M⁺, 64.73), 255 (100), 238 (29.91), 213 (4.60), 179 (73.29), 152 (16.92), 124 (25.01), 103(8.14), 96 (11.27), 78(13.08), 69(11.87), 50(8.83).

Crystals: yellow needles from EtOH m.p. 201.7-202.2 °C (Lit. m.p. 200-201°C).

CURRICULUM VITAE

Name Miss Charatda Punvittayagul

Date of birth Febuary 12, 1985

Education 2003, Certificate of high school, Wangkhamvittayakhom school, Kamphang-Phet, Thailand

2007, Bachelor of Science (Biology; 2nd Class honor), Naresuan University, Phitsanulok, Thailand

Publications

1. Charoensin S, Punvittayagul C, Pompimon W, Mevatee U and Wongpoomchai R. Toxicological and clastogenic evaluation of pinocembrin and pinostrobin isolated from *Boesenbergia pandurata* in Wistar rats. *Thai J Toxicol.* (inpress).
2. Punvittayagul C., Charoensin S., Taya S., Pompimon W., Wongpoomchai R. Effect of pinocembrin on xenobiotic-metabolizing enzymes in rat liver. Poster presentation at the 3rd Asian Pacific Regional ISSX Meeting, Bangkok, Thailand. May 10-12, 2009. *Drug Met. Rev.* 2009, 41(2): 72.
3. Taya S., Charoensin S., Punvittayagul C., and Wongpoomchai R. Effect of *Cleistocalyx nervosum* on antioxidant and detoxifying enzymes in Wistar rat. Poster presentation at the 3rd Asian Pacific Regional ISSX Meeting, Bangkok, Thailand. May 10-12, 2009. *Drug Met. Rev.* 2009, 41(2): 88-89.
4. Taya S., Punvittayagul C., Chewonarin T., and Wongpoomchai R. Effect of aqueous extract from *Cleistocalyx nervosum* on oxidative status in rat liver. Proceeding at the 2nd National Conference in Toxicology, Bangkok, Thailand. December 17-18, 2009. *Thai J Toxicol.* 24(2) 101-105.

Presentations

1. Punvittayagul C., Pompimon W., Wongpoomchai R. Rat liver mutagenicity and antimutagenicity of pinocembrin extracted from fingerroot rhizome (*Boesenbergia pandurata*). Poster presentation at the 8th Annual Biochemical Research Meeting. October 13-14, 2008, Department of Biochemistry, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand, page 43.
2. Charoensin S., Punvittayagul C., Taya S., Pompimon W., Mevatee U., and Wongpoomchai R. The inhibition of micronucleus formation of pinostrobin from fingerroot (*Boesenbergia pandurata*) in Wistar rat liver induced by diethylnitrosamine and its possibly inhibiting mechanism. Oral presentation at the Second International Conference on Natural Products for Health and Beauty (NATPRO). December 17-19, 2008, Naresuan University at Phayao, Phayao Province, Thailand, page 126.
3. Punvittayagul C., Pompimon W., Wongpoomchai R. Mutagenicity and antimutagenicity of pinocembrin extracted from fingerroot rhizome (*Boesenbergia pandurata*) in rat liver. Poster presentation at the 2nd International Conference on Natural Products for Health and Beauty (NATPRO). December 17-19, 2008, Naresuan University at Phayao, Phayao Province, Thailand, page 185.
4. Taya S., Charoensin S., Punvittayagul C., and Wongpoomchai R. Antioxidative properties of various extracts of ma-kiang (*Cleistocalyx nervosum* var. *paniala*). Poster presentation at the 2nd International Conference on Natural Products for Health and Beauty (NATPRO). December 17-19, 2008, Naresuan University at Phayao, Phayao Province, Thailand, page 223.
5. Taya S., Charoensin S., Punvittayagul C., Wongpornchai S, and Wongpoomchai R. Toxicological study of antioxidative substances extracted from Ma-kiang (*Cleistocalyx nervosum* var. *paniala*) in rat. Poster presentation at the International Congress for Innovation in Chemistry (PERCH-CIC Congress VI). May 3-6, 2009, Jomtien Plam Beach Hotel & Resort, Pattaya, Thailand, page 273.

6. Punvittayagul C., Charoensin S., Taya S., Pompimon W., Wongpoomchai R. Mutagenicity and antimutagenicity of pinocembrin isolated from *Boesenbergia pandurata* rhizome in rat liver. Poster presentation at the International Congress for Innovation in Chemistry (PERCH-CIC Congress VI). May 3-6, 2009, Jomtien Plam Beach Hotel & Resort, Pattaya, Thailand, page 299.
7. Punvittayagul C., Taya S., Charoensin S., Pompimon W., Wongpoomchai R. Evaluation of mutagenicity and antimutagenicity of pinocembrin by rat liver micronucleus test. Oral presentation at the 9th Annual Biochemical Research Meeting. October 8-9, 2009, Department of Biochemistry, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand, page 9.
8. Taya S., Punvittayagul C., and Wongpoomchai R. Effect of *Cleistocalyx nervosum* extract on oxidative stress in early stages of chemicals induced hepatocarcinogenesis. Oral presentation at the 9th Annual Biochemical Research Meeting. October 8-9, 2009, Department of Biochemistry, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand, page 10.
9. Punvittayagul C., Taya S., Pompimon W., Wongpoomchai R. Effect of pinocembrin on promotion stage in diethylnitrosamine-induced rat hepatocarcinogenesis. Poster presentation at the 2nd National Conference in Toxicology. December 17-18, 2009, Miracle Grand Convention Hotel, Bangkok, Thailand, page 170.

Scholarship

1. Center of excellence for innovation in chemistry (PERCH-CIC), year 2008-2009.



