

เอกสารอ้างอิง

- Adisakwattana S, Charoenlertkul P, Yibchok-Anun S. alpha-Glucosidase inhibitory activity of cyanidin-3-galactoside and synergistic effect with acarbose J Enzyme Inhib Med Chem. 2009; 24(1):65-69.
- Aebi H. Catalase in vitro. In: Packer L (ed.). Methods in Enzymology, Academic Press Inc., San Diego. 1984; 105:121-126.
- American Diabetic Association. Report of the expert committee on the diagnosis and classification of diabetes mellitus. Diabetes Care 1997; 20: 1183-1197.
- Asami DK, Hong YJ, Barrett DM, Mitchell AE. Comparison of the total phenolic and ascorbic acid content of freeze-dried and air-dried mulberry, strawberry, and corn grown using conventional, organic, and sustainable agricultural practices. Agric Food Chem 2003; 51: 1237-1241.
- Beutler E. Glutathion peroxidase (GSH-Px). In: Red Cell Metabolism: A Manual of Biochemical Methods. Grune & Stratton, New York. 1984; 74-76.
- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal Biochem 1976; 72: 248-254.
- Brownlee M, Cerami A, Vlassara H. Advanced glycosylation end products in tissue and biochemical basis of diabetic complication. N Engl J Med. 1988; 318: 1315-1322.
- Cao H, Hininger-Favier I, Kelly MA, Benaraba R, Dawson HD, Coves S, Roussel AM, Anderson RA. Green tea polyphenol extract regulates the expression of genes involved in glucose uptake and insulin signaling in rats fed a high fructose diet. J Agric Food Chem. 2007;55(15): 6372-6378.
- Chang CC, Yang MH, Wen HM, Chern JC. Estimation of total flavonoids content in propolis by two complementary colorimetric methods. J Food Drug Anal 2002; 10(2): 178-182.
- Chase CK, McQueen CE. Cinnamon in diabetes mellitus. Am J Health Syst Pharm. 2007; 64(10):1033-1035.
- Cheng AY, Fantus IG. Oral antihyperglycemic therapy for type 2 diabetes mellitus. CMAJ. 2005;172(2):213-26.
- Chiasson JL. Acarbose for the prevention of diabetes, hypertension, and cardiovascular disease in subjects with impaired glucose tolerance: the Study to Prevent Non-Insulin-Dependent Diabetes Mellitus (STOP-NIDDM) Trial. Endocr Pract. 2006 ;12 Suppl 1:25-30.
- Curtis J, Wilson C. Preventing type 2 diabetes mellitus. J Am Board Fam Pract. 2005;18(1):37-43.
- D'Autreaux B, Toledano MB. ROS as signalling molecules: mechanisms that generate specificity in ROS homeostasis. Nat Rev Mol Cell Biol. 2007;8(10):813-824.

- El-Alfy AT, Ahmed AA, Fatani AJ. Protective effect of red grape seeds proanthocyanidins against induction of diabetes by alloxan in rats. *Pharmacol Res.* 2005; 52(3):264-270.
- Elliou ss, Keim NL, Stern JS, Teff K, Havel PJ. Fructose, weight gain and the insulin resistance syndrome. *Am J Clin Nutr.* 2002; 76: 911-922.
- Eriksson JW. Metabolic stress in insulin's target cells leads to ROS accumulation - a hypothetical common pathway causing insulin resistance. *FEBS Lett.* 2007; 581(19):3734-42.
- Fine AM. Oligomeric proanthocyanidin complexes: history, structure, and phytopharmaceutical applications. *Alter Med Rev.* 2000; 5: 144–151.
- Garvey WT, Olefsky JM, Marshall S. Insulin induces progressive insulin resistance in cultured rat Adipocytes: sequential effects at receptor and multiple postreceptor sites *Diabetes* 1986; 35: 258–267.
- Gérard-Monnier D, Chaudiere J. Metabolism and antioxidant function of glutathione. *Pathol Biol (Paris):* 1996; 44(1):77-85.
- Guo H, Ling W, Wang Q, Liu C, Hu Y, Xia M, Feng X, Xia X. Effect of Anthocyanin-Rich Extract from Black Rice (*Oryza sativa* L. *indica*) on Hyperlipidemia and Insulin Resistance in Fructose-Fed Rats. *Plant Foods Hum Nutr.* 2007; 62(1):1-6.
- Halliwell B, Gutteridge JM. Role of free radicals and catalytical metal ions in human disease: An overview. *Methods Enzymol* 1990; 186: 1-85.
- Havsteen BH. The biochemistry and medical significance of the flavonoids. *Pharmacol Ther* 2002; 96: 67-202.
- Hermans MP, Levy JC, Morris RJ, Turner RC. Comparison of insulin sensitivity tests across a range of glucose tolerance from normal to diabetes *Diabetologia* 1999; 42: 678–687.
- Hwang IS, Ho H, Hoffman BB, Reaven GM. Fructose-induced insulin resistance and hypertension in rats. *Hypertension* 1987; 10: 512-516.
- Hwang IS, Ho H, Hoffman BB, Reaven GM. Fructose-induced insulin hypertension: parallels between human diseases and animal models. *Diabetes Care* 1991; 14: 195–202.
- Johnson RJ, Perez-Pozo SE, Sautin YY, Manitius J, Sanchez-Lozada LG, Feig DI, Shafiu M, Segal M, Glasscock RJ, Shimada M, Roncal C, Nakagawa T. Hypothesis: could excessive fructose intake and uric acid cause type 2 diabetes? *Endocr Rev.* 2009; 30(1):96-116.
- Joshi SS, Kuszynski A, Bagchi D. The cellular and molecular basis of health benefits of grape seed proanthocyanidin extract. *Curr Pharm Biotech.* 2001; 2: 187–200.

- Kamisoyama H, Honda K, Tominaga Y, Yokota S, Hasegawa S. Investigation of the anti-obesity action of licorice flavonoid oil in diet-induced obese rats. *Biosci Biotechnol Biochem*. 2008; 72(12):3225-3231.
- Kao YH, Hiipakka RA, Liao S. Modulation of endocrine systems and food intake by green tea epigallocatechin gallate. *Endocrinology*. 2000;141(3):980-7.
- Kaplan F, Al-Majali K, Betteridge DJ. PPARs, insulin resistance and type 2 diabetes. *J Cardiovasc Risk*. 2001; 8: 211–217.
- Karthikeyan K, Bai BR, Devaraj SN. Cardioprotective effect of grape seed proanthocyanidins on isoproterenol-induced myocardial injury in rats. *Int J Cardiol*. 2007; 115: 326–333.
- Krawinkel MB, Keding GB. Bitter melon (*Momordica charantia*): A dietary approach to hyperglycemia. *Nutr Rev*. 2006; 64(7 Pt 1):331-337.
- Kumazawa S, Taniguchi M, Suzuki Y, Shimura M, Kwon MS, Nakayama T. Antioxidant activity of polyphenols in carob pods. *J Agric Food Chem*. 2002; 50(2):373-377.
- Leiter LA, Ceriello A, Davidson JA, Hanefeld M, Monnier L, Owens DR, Tajima N, Tuomilehto J; International Prandial Glucose Regulation Study Group. Postprandial glucose regulation: new data and new implications. *Clin Ther*. 2005; 27 Suppl B:S42-56.
- Lewis GF, Carpentier A, Adeli K, Giacca A. Disordered fat storage and mobilization in the pathogenesis of insulin resistance and type 2 diabetes. *Endocr Rev*. 2002; 2: 201-229.
- Lindmark S, Buren J, Eriksson JW. Insulin resistance, endocrine function and adipokines in type 2 diabetes patients at different glycaemic levels: Potential impact for glucotoxicity in vivo. *Clin Endocrinol*. 2006; 65: 301–309.
- Liu IM, Tzeng TF, Liou SS, Lan TW. Myricetin, a naturally occurring flavonol, ameliorates insulin resistance induced by a high-fructose diet in rats. *Life Sci*. 2007; 81:1479-1488.
- Maeda N, Shimomura I, Kishida K, Nishizawa H, Matsuda M, Nagaretani H, Furuyama N, Kondo H, Takahashi M, Arita Y, Komuro R, Ouchi N, Kihara S, Tochino Y, Okutomi K, Horie M, Takeda S, Aoyama T, Funahashi T, Matsuzawa Y. Diet-induced insulin resistance in mice lacking adiponectin/ACRP30. *Nat Med*. 2002; 8: 731-737.
- Maiese K, Chong ZZ, Shang YC. Mechanistic insights into diabetes mellitus and oxidative stress. *Curr Med Chem*. 2007;14 (16):1729-1738.
- Marklund S and Marklund G. Involvement of the superoxide anion radical in the autoxidation of pyragalol and a convenient assay for superoxide dismutase. *Eur J Biochem*. 1974; 47: 469-474.
- Matsuzawa Y, Shimomura I, Nakamura T, Keno Y, Tokunaga K. Pathophysiology and pathogenesis of visceral fat obesity. *Diabetes Res Clin Pract*. 1994;24 Suppl: s111-s116.

- Mayes PA. Intermediary metabolism of fructose. *Am J Clin Nutr.* 1993; 58: S754-S765.
- Meneilly GS, Tessier D. Diabetes in elderly adults. *J Gerontol A Biol Sci Med Sci.* 2001; 56(1):M5-M13.
- Miele C, Riboulet A, Maitan MA, Oriente R, Romano C, Formisano R. Human glycated albumin affects glucose metabolism in L6 skeletal muscle cells by impairing insulin-induced insulin receptor substrate (IRS) signaling through a protein kinase Ca-mediated mechanism. *J Biol Chem.* 2003; 278: 47376-47387.
- Miller A, Adeli K. Dietary fructose and the metabolic syndrome. *Curr Opin Gastroenterol.* 2008; 24(2):204-209.
- Moreno DA, Ilic N, Poulev A, Brasaemle DL, Fried SK, Raskin I. Inhibitory effects of grape seed extract on lipases. *Nutrition* 2003; 19: 876-879.
- Moskaug J, Carlsen H, Myhrstad MCW, Blomhoff R. Polyphenols and glutathione synthesis regulation. *Am J Clin Nutr.* 2005; 81: 277S-283S.
- Murase T, Nagasawa A, Suzuki J, Hase T, Tokimitsu I. Beneficial effects of tea catechins on diet-induced obesity: stimulation of lipid catabolism in the liver. *Int J Obes Relat Metab Disord.* 2002 ; 26(11):1459-1464.
- Olefsky, JM. Treatment of insulin resistance with peroxisome proliferator-activated receptor gamma agonists. *J Clin Invest.* 2000; 106: 467-472.
- Pinet M, Blay M, Blade MC, Salvado MJ, Arola L, Ardevol A. Grape seed-derived procyanidins have an antihyperglycemic effect in streptozotocin-induced diabetic rats and insulinomimetic activity in insulin-sensitive cell lines. *Endocrinology.* 2005 ;145(11):4985-4990.
- Rosak C. The pathophysiologic basis of efficacy and clinical experience with the new oral antidiabetic agents. *J Diabetes Complications.* 2002;16(1):123-132.
- Rotter V, Nagaev I, Smith U. Interleukin-6 (IL-6) induces insulin resistance in 3T3-L1 adipocytes and is, like IL-8 and tumor necrosis factor-alpha, overexpressed in human fat cells from insulin-resistant subjects. *J Biol Chem.* 2003; 278: 45777-45784.
- Ruzaidi A, Amin I, Nawalyah AG, Hamid M, Faizul HA. The effect of Malaysian cocoa extract on glucose levels and lipid profiles in diabetic rats. *J Ethnopharmacol.* 2005; 98: 55- 60.
- Saito M, Hosoyama H, Ariga T, Kataoka S, Yamaji N. Antiulcer activity of grape seed extract and procyanidins. *J Agric Food Chem* 1998; 46(4): 1460-1464.
- Saltiel AR, Olefsky JM. Thiazolidinediones in the treatment of insulin resistance and type II diabetes. *Diabetes* 1996; 56: 1661-1669.

- Sedlak J, Lindsay RHL. Estimation of total and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Anal Biochem.* 1968; 25: 192-205.
- Semenkovich CF. Insulin resistance and atherosclerosis. *J Clin Invest.* 2006;116(7):1813-1822.
- Thorburn AW, Storlien LH, Jenkins AB, Khouri S, Kraegen EW. Fructose-induced in vivo insulin resistance and elevated plasma triglyceride levels in rats. *Am J Clin Nutr.* 1989; 49: 1155-1163.
- Uchiyama M, Mihara M. Determination of malonaldehyde precursor in tissue by thiobarbituric acid test. *Anal Biochem.* 1978; 86: 271-278.
- Vats V, Grover JK, Rathi SS. Evaluation of anti-hyperglycemic and hypoglycemic effect of *Trigonella foenum-graecum* Linn, *Ocimum sanctum* Linn and *Pterocarpus marsupium* Linn in normal and alloxanized diabetic rats. *J Ethnopharmacol.* 2002; 79: 95-100.
- Watson, R.T., Pessin, J.E. Intracellular organization of insulin signaling and GLUT4 translocation. *Recent Progress in Hormone Research.* 2001; 56: 175-193.
- Yamakoshi J, Kataoka S, Koga T, Ariga T. Proanthocyanidin-rich extract from grape seeds attenuates the development of aortic atherosclerosis in cholesterol-fed rabbits. *Atherosclerosis.* 1999; 142 (1):139-149.
- Yilmaz Y, Toledo RT. Major flavonoids in grape seeds and skins: antioxidant capacity of catechin, epicatechin, and gallic acid. *J Agric Food Chem.* 2004; 52(2): 255-260.
- Zavaroni I, Sander S, Scott S, Reaven G M. Effect of fructose feeding on insulin secretion and insulin action in the rat. *Metabolism* 1980; 10: 970-973.
- Zern TL, Wood RJ, Greene C, West KL, Liu Y, Aggarwal D, Shachter NS, Fernandez ML. Grape polyphenols exert a cardioprotective effect in pre- and postmenopausal women by lowering plasma lipids and reducing oxidative stress. *J Nutr.* 2005;135(8):1911-1917.



