



## REFERENCES

- Ahma, D.F., Ahmad, I., Khan, M.S., 2005. Indole acetic acid production by the indigenous isolates of *Azotobacter* and fluorescent *Pseudomonas* in the presence and absence of tryptophan. *Turk. J. Biol.*, 29:29-34.
- Anderson, A.B., 1991. Symbiotic and asymbiotic germination and growth of *Spiranthes magnicamporum* (Orchidaceae). *Lindleyana.*, 6:183-186.
- Arditti, J., Ernst, R., Yam, T.W., Glabe, C., 1990. The contributions of orchid mycorrhizal fungi to seed germination: a speculative review. *Lindleyana.*, 5:249-255.
- Arditti, J., Krikorain, A.D., 1996. Orchid micropropagation: the part from laboratory to commercialization and an account of several unappreciated investigators. *Bot. J. Linn. Soc.*, 122:183-241.
- Arditti, J., Michaud, J.D., Oliva, A.P., 1981. Seed germination of North American orchids. I. Native California and related species of *Calypso*, *Epipactis*, *Goodyera*, *Piperia*, and *Platanthera*. *Bot. Gaz.*, 142:442-453.
- Athipunyakom, P., Manoch, L., Piluek, C., 2004a. Isolation and identification of mycorrhizal fungi from eleven terrestrial orchids. *Kasetsart. J. (Nat Sci).*, 38:216-228.

- Athipunyakom, P., Manoch, L., Piluek, C., Artjariyasripong, S., Tragulrung, S., 2004b. Mycorrhizal fungi from *Spathoglottis Plicata* and the use of these fungi to germinate seeds of *S. plicata in vitro*. *Kasetsart. J. (Nat Sci)*, 37:83-93.
- Atkin, C.L., Neiland, J.B., Phaff, H.J., 1970. Rhodotorulic acid from species of *Leucosporidium*, *Rhodospondium*, *Rhodotorula*, *Sporidiobolus* and *Sporobolomyces*, and a new alanine-containing ferrichrome from *Cryptococcus meliviosum*. *J. Bacteriol.*, 103:722-733.
- Baakza, A., Vala, A.K., Dave, B.P., Dube, H.C., 2004. A comparative study of siderophore production by fungi from marine and terrestrial habitats. *J. Exp. Mar. Biol. Ecol.*, 311:1-9
- Banks, P.D. (2003). *Handy Pocket Guide to Orchids*. Singapore; Periplus Editions (HK) Ltd.
- Barnett, H.L., and Hunter, B.B. (1987). *Illustrated Genera of Imperfect Fungi* 4<sup>th</sup> ed. New York: Macmillan Publishing Company.
- Batty, A.L., Brundrett, M.C., Dixon, K.W., Sivasithamparam, K., 2006a. *In situ* symbiotic seed germination and propagation of terrestrial orchid seedlings for establishment at field sites. *Aust. J. Bot.*, 54:375-381.
- Batty, A.L., Brundrett, M.C., Dixon, K.W., Sivasithamparam, K., 2006b. New method to improve symbiotic propagation of temperate terrestrial orchid seedlings from axenic culture to soil. *Aust. J. Bot.*, 54:367-374.

- Batty, A.L., Dixon, K.W., Brundrett, M.C., Sivasithamparam, K., 2001. Long-term storage of mycorrhizal fungi and seed as a tool for the conservation of endangered Western Australian terrestrial orchids. *Aust. J. Bot.*, 49:619-628.
- Bayman, P., Lebro'n, L.L., Tremblay, R.L., Lodge, D.J., 1997. Variation in endophytic fungi from roots and leaves of *Lepanthes* (Orchidaceae). *New Phytol.*, 135:143-149.
- Beyrle, H.F., Smith, S.E., Peterson, R.L., Franco, C.M.M., 1995. Colonization of *Orchis morio* protocorms by a mycorrhizal fungus: effects of nitrogen nutrition and glyphosate modifying the responses. *Can. J. Bot.*, 73:1128-1140.
- Bougoure, J.J., Bougoure, D.S., Cairney, J.W.G., Dearnaley, J.D.W., 2005. ITS-RFLP and sequence analysis of endophytes from *Acianthus*, *Caladenia* and *Pterostylis* (Orchidaceae) in southern Queensland. *Mycol. Res.*, 109:452-460.
- Brundrett, M., 2004. Diversity and classification of mycorrhizal associations. *Biol. Rev.*, 79:473-495.
- Brundrett, M., Bougher, N., Dell, B., Grove, T., and Malajczuk, N. (1996). *Working with Mycorrhizas in Forestry and Agriculture*. Canberra: Australian Centre for International Agricultural Research.
- Brundrett, M., Scade, C.A., Batty, A.L., Dixon, K.W., 2003. Sivasithamparam K., Development of *in situ* and *ex situ* seed baiting techniques to detect mycorrhizal fungi from terrestrial orchid habitats. *Mycol. Res.*, 107:1210-1220.

- Brundrett, M., Sivasithamparam, K., Ramsay, M., Krauss, S., Taylor, R., Bunn, E., Hicks, A., Karim, N.A., Debeljak, N., Mursidawati, S., Dixon, B., Batty, A., Bower, C., and Brown, A. (2001). *Orchid Conservation Techniques Manual, First International Orchid Conservation Congress-Training Course*. Perth: Plant Science, King Park & Botanic Garden.
- Castañeda, M., Pardio, S., Orrantia, B., Lango, F., 2005. Influence of water temperature and salinity on seasonal occurrences of *Vibrio cholerae* and enteric bacteria in oyster-producing areas of Veracruz, México. *Mar. Pollut. Bull.*, 50:1641-1648.
- Chaiharn, M., Chunnaleuchanon, S., Lumyong, S., 2009. Screening siderophore producing bacteria as potential biological control agent for fungal rice pathogens in Thailand. *World. J. Microbiol. Biotechnol.*, 25:1919–1928.
- Chaiharn, M., Lumyong, S., 2011. Screening and optimization of indole-3-acetic acid production and phosphate solubilization from rhizobacteria aimed at improving plant growth. *Curr. Microbiol.*, 62:173-181.
- Chou, L.C., Chang, D.C.N., 2004. Asymbiotic and symbiotic seed germination of *Anoectochilus formosanus* and *Haemaria discolor* and their F<sub>1</sub> hybrids. *Botany. Bulletin. Academy. Science.*, 45:143-147.
- Clements, M., 1988. Orchid mycorrhizal associations. *Lindleyana.*, 3:73-86.
- Cordier, C., Gianinazzi, S., Gianinazzi-Pearson, V., 1998. Colonization pattern of root tissue by *Phytophthora nicotianae* var. *parasitica* related to reduced disease in mycorrhizal tomato. *Plant. Soil.*, 185:223–232.

- Currah, R.S., Zelmer, C., 1992. A key and notes for the genera of fungi mycorrhizal with orchids and a new species in the genus *Epulorhiza*. *Rept. Tottori. Mycol. Inst.*, 30:43-59.
- Currah, R.S., Zettler, I.W., McInnis, T.M., 1997. *Epulorhiza inquilina* sp. nov. from *Plantanthera* (Orchidaceae) and a key to *Epulorhiza* species. *Mycotaxon.*, 61:335-342.
- Dasa, A., Majo'n, J.L., Camacho, M., 2006. Effect of carbon and nitrogen sources, pH and temperature on *in vitro* culture of several isolates of *Amanita caesarea* (Scop.:Fr.) Pers. *Mycorrhiza.*, 16:133-136.
- Davies, P.J. (2004). *Plant Hormones, Biosynthesis and Signal Transduction and Action*. (pp. 1-35), Netherlands: Kluwer Academic Publishers.
- Dearnaley, J.D.W., 2007. Further advances in orchid mycorrhizal research. *Mycorrhiza.*, 17:475-486.
- Deshwal, V.K., Dubey, R.C., Maheshwari, D.K., 2003. Isolation of plant-growth promoting strains of *Bradyrhizobium* (*Arachis* sp.) with biocontrol potential against *Macrophomina phaseolina* causing charcoal rot of peanut. *Curr. Sci.*, 84: 443-448.
- Douds, D.D., Nagahashi, G., Reider, C., Hepperly, P.R., 2007. Inoculation with arbuscular mycorrhizal fungi increases the yield of potatoes in a high P soil. *Biol. Agric. Hortic.*, 25:67-78.
- Duchesne, L.C, Peterson, R.L, Ellis, B.E., 1989. The future of ectomycorrhizal fungi as biological control agents. *Phytoprotection.*, 70:51-57.

- Duke University. 2011. "Conserved primer sequences for PCR amplification and sequencing from nuclear ribosomal RNA" [Online]. Available <http://www.biology.duke.edu/fungi/mycolab/primers.html> (8 January 2011).
- Eason, W.R., Newman, E.I., Chuba, P.N., 1991. Specificity of interplant cycling of phosphorus: the role of mycorrhizas. *Plant. Soil.*, 137:267-274.
- Ehmann, A., 1977. The van Urk-Salkowski reagent-a sensitive and specific chromogenic reagent for silica gel thin-layer chromatographic detection and identification of indole derivatives. *J. Chromatogr.*, 132:267-276.
- El-Abyad, M.S.H., Webster, J., 1968. Studies on pyrophilous discomycetes: II. Competition. *Br. Mycol. Soc.* 51:369-375.
- El-Mehalawy, A.A., Hassanin, S.M., Hassanin, N.M., Zaki, S.A., 2007. Induction of resistance and biocontrol of rhizoctonia in cotton against damping-off disease by rhizosphere yeasts and fungi. *New Egypt. J. Microbiol.*, 17:148-167.
- El-Tarabily, K.A., Nassar, A.H., Sivasithamparam, K., 2008. Promotion of growth of bean (*Phaseolus vulgaris* L.) in a calcareous soil by a phosphate-solubilizing, rhizosphere-competent isolate of *Micromonospora endolithica*. *Appl. Soil. Ecol.*, 39:161-171.
- Erturk, Y., Ercisli, S., Sekban, R., Haznedar, A., Donmez, M.F., 2008. The effect of PGPR on rooting and growth of tea (*Camellia sinensis* var. *Sinensis*) cuttings. *Romanian Biotechnol. Lett.*, 13:3747-3756.

- Ezilon maps*. 2011. "Thailand Map - Political Map of The Kingdom of Thailand"  
 [Online]. Available <http://www.ezilon.com/maps/asia/thailand-maps.html>  
 (21 November 2011).
- Fogel, R., Peck, S.B., 1975. Ecological studies of hypogeous fungi. I. *Coleoptera* associated with sporocarps. *Mycologia.*, 67:741-747.
- Fries, L., 1956. Studies in the physiology of *Coprinus*. II. Influence of pH, metal factors and temperature. *Sven. Bot. Tidskr.*, 50:47-96.
- Goodfellow, M., Williams, S.T., 1983. Ecology of actinomycetes. *Ann. Rev. Microbiol.*, 37:189-215.
- Gordon, S.A., Weber, R.P., 1951. Colorimetric estimation of indoleacetic acid. *Plant. Physiol.*, 26:192-195.
- Grandmaison, J., Olah, G.M., Calsteren, M.R., Furlan, V., 1993. Characterisation and localisation of plant phenolics likely involved in the pathogen resistance expressed by endomycorrhizal roots. *Mycorrhiza.*, 3:155-164.
- Hadley, G. (1982). Orchid mycorrhiza. In: Arditti, J., ed. *Orchid Biology Reviews and Perspectives II*. (pp. 85-115), Comstock: Cornell University Press.
- Harley, J.L., and Smith, S.E. (1983). *Mycorrhizal Symbiosis*. London: Academic Press.
- Hasan, H.A.H., 2002. Gibberellin and auxin production by plant root-fungi and their biosynthesis under salinity-calcium interaction. *Acta. Microbiol. Immunol. Hung.*, 49:105-118.
- Haselwandter, K., 1995. Mycorrhizal fungi: siderophore production. *Crit. Rev. Biotechnol.*, 15:287-91.

- Horton, T.R., Bruns, T.D., Parker, V.T., 1999. Ectomycorrhizal fungi associated with *Arctostaphylos* contribute to *Pseudotsuga menziesii* establishment. *Can. J. Bot.*, 77: 93–102.
- Hung, L.L., Trappe, J.M., 1987. Ectomycorrhizal inoculation of Douglas-fir transplanted container seedlings with commercially produced inoculum. *New Forests.*, 1:141-152.
- Ingham, E.R., Massicotte, H.B., 1994. Protozoan communities around conifer roots colonized by ectomycorrhizal fungi. *Mycorrhiza.*, 5:53-61.
- Jeon, J.L., Lee, S.S., Kim, H.Y., Ahn, T.S., Song, H.G., 2003. Plant growth promotion in soil by some inoculated microorganisms. *J. Microbiol.*, 41:271-276.
- Jing, L., Qing, L.S., Yi, L., Li, C.X., 2009. Effects of increased ammonia on root/shoot ratio, grain yield and nitrogen use efficiency of two wheat varieties with various N supply. *Plant. Soil. Environ.*, 7:273-280.
- Johnson, T.R., Stewart, S.L., Dutra, D., Kane, M.E., 2007. Richardson L., Asymbiotic and symbiotic seed germination of *Eulophia alta* (Orchidaceae)-preliminary evidence for the symbiotic culture advantage. *Plant. Cell. Tiss. Organ. Cult.*, 90:313-323.
- Jonbloed, R.H., Borst-Pauwels, G.W.F.H., 1990. Effect of ammonium and pH on growth of some ectomycorrhizal fungi *in vitro*. *Acta. Bot. Neerl.*, 39:349-358.

- Kahiluoto, H., Vestberg, M., 1998. The effect of arbuscular mycorrhiza on biomass production and phosphorus uptake from sparingly soluble sources by leek (*Allium porrum* L.) in Finnish field soils. *Biol. Agric. Hort.*, 16:65-85.
- Kalinowski, B.E., Liermann, L.J., Givens, S., Brantley, S.L., 2000. Rates of bacteria-promoted solubilization of Fe from minerals: A review of problems and approaches. *Chem. Geol.*, 169:357-370.
- Kasana, R.C., Salwan, R., Dhar, H., Dutt, S., Gulati, A., 2008. A rapid and easy method for the detection of microbial celluloses on agar plates using Gram's iodine. *Curr. Microbiol.*, 57:503-507.
- Kato, R., Takatsuna, S., Wada, T., Narihara, Y., Suzuki, T., 2003. *In vitro* phosphorylation of proteins in IAA-treated primary roots and coleoptiles in *Zea mays*. *Bull. Yamagata Univ. (Nat Sci)*, 15: 89-104.
- Khamna, S. (2009). *Biodiversity and secondary metabolites of Actinomycetes from rhizosphere soils of some medicinal plants*. Ph.D. Thesis, Chiang Mai University.
- Khamna, S., Yokota, A., Peberdy, J.F., Lumyong, S., 2010. Indole-3-acetic acid production by *Streptomyces* sp. isolated from some Thai medicinal plant rhizosphere soils. *Eur. Asia. J. BioSci.*, 4:23-32.
- Khan, S.A., Hamayun, M., Kim, H.Y., Yoon, H.J., Lee, I.J., Kim, J.G., 2009. Gibberellin production and plant growth promotion by a newly isolated strain of *Gliomastix murorum*. *World. J. Microbiol. Biotechnol.*, 25:829-833.

- Kim, Y.K., Xiao, C.L., Rogers, J.D., 2005. Influence of culture media and environmental factors on mycelia growth and pycnidial production of *Sphaeropsis pyriputrescens*. *Mycologia.*, 97:25-32.
- Koga, J., Adachi, T., Hidaka, H., 1991. IAA biosynthetic pathway from tryptophan via indole-3-pyruvic acid in *Enterobacter cloaceae*. *Agric. Biol. Chem.*, 55:701-706.
- Kristiansen, K.A., Freudenstein, J.V., Rasmussen, F.N., Rasmussen, H.N., 2004. Molecular identification of mycorrhizal fungi in *Neuwiedia veratrifolia* (Orchidaceae). *Mol. Phylogenet. Evol.*, 33:251-258.
- Kristiansen, K.A., Taylor, D.L., Kjølner, R., Rasmussen, H.N., 2001. Rosendahl S., Identification of mycorrhizal fungi from single pellets of *Dactylorhiza majalis* (Orchidaceae) using single-strand conformation polymorphism and mitochondrial ribosomal large subunit DNA sequences. *Mol. Ecol.*, 10:2089-2093.
- Kumla, J., Danell, E., Bussaban, B., Lumyong, S., 2011. Suitable growth conditions and nutrition factors on *in vitro* culture of *Phlebopus portentosus* (Boletales). *Chiang Mai. J. Sci.* 38:156–159.
- Lee, S.S., 2002. A review of orchid mycorrhizae in Korea. *Plant. Pathol. J.*, 18:169-178.
- Lee, S.S., Patahayah, M., Chong, W.S., Lapeyrie, F., 2008. Successful ectomycorrhizal inoculation of twodipterocarp species with a locally isolated fungus in peninsular Malaysia. *J. Trop. Forest. Sci.*, 20:237-247.

- Lindahl, B., Stenlid, J., Olsson, S., Finlay, R., 1999. Translocation of  $^{32}\text{P}$  between interacting mycelia of a wood decomposing fungus and ectomycorrhizal fungi in microcosm systems. *New. Phytol.*, 144:183–193.
- Linderman, R., Call, C., 1977. Enhanced rooting of woody plant cuttings by mycorrhizal fungi. *J. Am. Soc. Hortic. Sci.*, 102:629–632.
- Little, L.R., Maun, M.A., 1996. The ‘*Ammophila* problem’ revisited: A role for mycorrhizal fungi. *J. Ecol.*, 84:1-7.
- Lussenhop, J., Fogel, R., 1999. Seasonal changes in phosphorus content of *Pinus strobus*-*Cenococcum geophilum* ectomycorrhizae. *Mycologia.*, 91:742-746.
- Ma, M., Tan, T.K., Wong, S.M., 2003. Identification and molecular phylogeny of *Epulorhiza* isolates from tropical orchids. *Mycol. Res.*, 107:1041-1049.
- Mahmoud, A.L.E., Abd-allah, M.A., 2001. Siderophores production by some microorganisms and their effect on *Bradyrhizobium*-mungbean symbiosis. *Int. J. Agric. Biol.*, 3:157-162.
- Maor, R., Haskin, S., Levi-Kedmi, H., Sharon, A., 2004. In planta production of indole-3-acetic acid by *Colletotrichum gloeosporioides* f. sp. *aeschynomene*. *Appl. Environ. Microbiol.*, 70:1852-1854.
- Marx, D.H., Bryan, W.C., Davey, C.B., 1970. Influence of temperature on aseptically synthesized ectomycorrhizae by *Thelephora terrestris* and *Pisolithus tinctorius* on loblolly pine. *For. Sci.*, 16:424-431.
- McCormick, M.K., Whingham, D.F., O’Neill, J., 2004. Mycorrhizal diversity in photosynthetic terrestrial orchids. *New. Phytol.*, 163:425-438.

- McGonigle, T.P., Miller, M.H., Evans, D.G., Fairchild, G.L., Swan, J.A., 1990. A new method which gives an objective measure of colonization of roots by vesicular-arbuscular mycorrhizal fungi. *New. Phytol.*, 115:495-501.
- McKendrick, S.L., Leake, J.R., Taylor, D.L., Read, D.J., 2000. Symbiotic germination and development of myco-heterotrophic plants in nature: ontogeny of *Corallorhiza trifida* and characterization of its mycorrhizal fungi. *New. Phytol.*, 145:523-537.
- McKendrick, S.L., Leake, J.R., Taylor, D.L., Read, D.J., 2002. Symbiotic germination and development of the mycoheterotrophic orchid *Neottia nidus-avis* in nature and its requirement for locally distributed *Sebacina* spp. *New. Phytol.*, 154: 233–247.
- Mei-ling, X.U., Jiao-jun, Z.H.U., Hong-zhang, K., Ai-hua, X.U., Jin-xin, Z., Feng-qin, L.I., 2008. Optimum conditions for pure culture of major ectomycorrhizal fungi obtained from *Pinus sylvestris* var. *mogolica* plantations in southeastern Keerqin sandy, China. *J. Forest. Res.*, 19:113-118.
- Mikola, P., 1970. Mycorrhizal inoculation in afforestation. *Int. Rev. For. Res.*, 3:123-196.
- Milagres, A.M.F., Machuca, A., Napoleão, D., 1999. Detection of siderophore production from several fungi and bacteria by a modification of chrome azurol S (CAS) agar plate assay. *J. Microbiol. Methods.*, 37:1–6.
- Miyoshi, K., Mii, M., 1995. Enhancement of seed germination and protocorm formation in *Calanthe discolor* (Orchidaceae) by NaOCl and polyphenol absorbent treatments. *Plant. Tiss. Cult. Lett.*, 12:267–272.

- Molina, R., and Trappe, J.M. (1984). *Mycorrhiza Management in Bareroot Nurseries*. Oregon: College of Forestry Oregon State University.
- Moore, R.T., 1987. The genera of *Rhizoctonia*-like fungi; *Ascorhizoctonia*, *Ceratorhiza* gen. nov., *Moniliopsis* and *Rhizoctonia*. *Mycotaxon.*, 29: 91-99.
- Nanakorn, W., Indharamusika, S., 1998. *Ex-situ* conservation of native Thai orchids at Queen Sirikit Botanic Garden. *Pure. Appl. Chem.*, 70:2115-2122.
- Nasim, G., Rahman, M., Shabbir, A., 2004. Effect of indole acetic acid on *in vitro* growth and biomass production of some soil fungi. *Pak. J. Biol. Sci.*, 12:2039-2044.
- Neilands, J.B., 1995. Siderophores: structure and function of microbial iron transport compounds. *J. Biol. Chem.*, 270:26723-26726.
- Niemi, K., Vuorinen, T., Ernstsén, A., Häggman, H., 2002. Ectomycorrhizal fungi and exogenous auxins influence root and mycorrhiza formation of Scots pine hypocotyl cuttings *in vitro*. *Tree. Physiol.*, 22:1231-1239.
- Nontachaiyapoom, S., Sasirat, S., Manoch, L., 2010. Isolation and identification of *Rhizoctonia*-like fungi from roots of three orchid genera, *Paphiopedilum*, *Dendrobium*, and *Cymbidium*, collected in Chiang Rai and Chiang Mai provinces of Thailand. *Mycorrhiza.*, 20:459-471.
- Ovando, I., Damon, A., Bello, R., Ambrosio, D., Albores, V., Adriano L., Salvador, M., 2005. Isolation of endophytic fungi and their potential for the tropical epiphytic orchids *Cattleya skinneri*, *C. aurantiaca* and *Brassavola nodosa*. *Asian. J. Plant. Sci.*, 4:309-315.

- Pal, K.K., Gardener, B.M. 2006. "Biological Control of Plant Pathogens." *The Plant Health Instructor*. DOI: 10.1094/PHI-A-2006-1117-02. [Online] Available <http://www.apsnet.org/edcenter/advanced/topics/Pages/BiologicalControl.aspx> (21 November 2011).
- Pérez-Miranda, S., Cabirol, N., George-Téllez, R., Zamudio-Rivera, L.S., Fernández, F.J., 2007. O-CAS, a fast and universal method for siderophore detection. *J. Microbiol. Meth.*, 70:127-131.
- Porrás-Alfaro, A., Bayman, P., 2007. Mycorrhizal fungi of Vanilla: diversity, specificity and effects on seed germination and plant growth. *Mycologia.*, 99:510-525.
- Promptutha, I., Jeewon, R., Lumyong, S., McKenzie, E.H.C., Hyde, K.D., 2005. Ribosomal DNA fingerprinting in the identification of non sporulating endophytes from *Magnolia liliifera* (Magnoliaceae). *Fungal Divers.*, 20:167-186.
- Puplett, D. 2006. "Tree for life: restoration the Caledonian forest; mycorrhizas." [Online]. Available <http://www.treesforlife.org.uk/forest/ecological/mycorrhizas.html> (28 April 2006).
- Rabatin, S.C., Stinner, B.R., 1989. The significance of vesicular-arbuscular mycorrhizal fungal- soil macroinvertebrate interactions in agroecosystems. *Agric. Ecosyst. Environ.*, 27:195-204.
- Rasmussen, H.N. (1995). *Terrestrial Orchids from Seed to Mycotrophic Plants*. (pp. 142-233), New York: Cambridge University Press.
- Robert, P. (1999). *Rhizoctonia-Forming Fungi a Taxonomic Guide*. Great Britain: The Trustees, Royal Botanic Gardens, Kew.

- Robinson, M., Riov, J., Sharon, A., 1998. Indole-3-acetic acid biosynthesis in *Colletotrichum gloeosporioides* f. sp. *aeschynomene*. *Appl. Environ. Microbiol.*, 64:5030-5032.
- Roy, M., Watthana, S., Stier, A., Richard, F., Vessabutr, S., Selosse, M.A., 2009. Two mycoheterotrophic orchids from Thailand tropical dipterocarpacean forests associate with a broad diversity of ectomycorrhizal fungi. *BMC. Biol.*, 7:51.
- Ruanpanun, P., Tangchitsomkid, N., Hyde, K.D., Lumyong, S., 2010. *Actinomycetes* and fungi isolated from plant-parasitic nematode infested soils: screening of the effective biocontrol potential, indole-3-acetic acid and siderophore production. *World. J. Microbiol. Biotechnol.*, 26:1569-1578.
- Rygiewicz, P.T., Andersen, C.P., 1994. Mycorrhizae alter quality and quantity of carbon allocated below ground. *Nature.*, 369:58-60.
- Sánchez, F., Honrubia, M., Torres, P., 2001. Effects of pH and temperature on *in vitro* cultures of ectomycorrhizal fungi from Mediterranean forests. *Mycol.*, 22:243-258.
- Sangthong, S. (2002). *Effect of Endomycorrhizal Fungus on Growth of Dendrobium scabrilingue Lindl. and its Hybrids*. M.S. Thesis, Chiang Mai University.
- Sanmee, R., Lumyong, P., Dell, B., Lumyong, S., 2010. *In vitro* cultivation and fruit body formation of black bolete, *Phlebopus portentus*, a popular edible ectomycorrhizal fungus in Thailand. *Mycoscience.*, 51:15-22.

- Santisuk, T., Chayamarit, K., Pooma, R., and Suddee, S. (2006). *Thailand Red Data: Plants*. Bangkok: Office of Natural Resources and Environmental Policy and Planning (ONEP).
- Schweiger, P.F., Robson, A.D., Barrow, N.J., 1995. Root hair length determines beneficial effect of a *Glomus* species on shoot growth of some pasture species. *New. Phytol.*, 131:247-254.
- Selosse, M.A., Tacon, F.L., 1998. The land flora: a phototroph-fungus partnership?. *Trends. Ecol. Evol.*, 13:15-20.
- Setälä, H., 1995. Growth of birch and pine seedlings in relation to grazing by soil fauna on ectomycorrhizal fungi. *Ecology.*, 76:1844-1851.
- Shefferson, R.P., Weiss, M., Kull, T., Taylor, D.L., 2005. High specificity generally characterizes mycorrhizal association in rare lady's slipper orchids, genus *Cypripedium*. *Mol. Ecol.*, 14:613-626.
- Shimura, H., Sadamoto, M., Matsuura, M., Kawahara, T., Naito, S., Koda, Y., 2009. Characterization of mycorrhizal fungi isolated from the threatened *Cypripedium macranthos* in a northern island of Japan: two phylogenetically distinct fungi associated with the orchids. *Mycorrhiza.*, 19:525-534.
- Sivasithamparam, K., Dixon, K.W., and Barrett, R.L. (2002). *Microorganisms in Plant Conservation and Biodiversity*. (pp.195-226), Netherlands: Kluwer, Dordrecht.
- Smith, S.E., and Read, D.J. (2008). *Mycorrhizal Symbiosis 3<sup>rd</sup> ed.* (pp. 419-463), San Diego: Academic Press.

- Sridevi, M., Mallaiah, K.V., 2008. Production of indole-3-acetic acid by *Rhizobium* isolates from *Sesbania* species. *Plant. Sci. Res.*, 1:13-16.
- Stewart, S.L., Kane, M.E., 2006. Symbiotic seed germination of *Habenaria macroceratitis* (Orchidaceae), a rare Florida terrestrial orchid. *Plant. Cell. Tissue. Organ. Cult.*, 86:159-167.
- Stewart, S.L., Zettler, L.W., 2002. Symbiotic germination of three semi-aquatic rein orchids (*Habenaria repens*, *H. quinquisetata*, *H. macroceratitis*) from Florida. *Aquat. Bot.*, 72:25-35.
- Suarez, J.P., Weiss, M., Abele, A., Garnica, S., Oberwinkler, F., Kottke, I., 2006. Diverse tulasnellid fungi from mycorrhizas with epiphytic orchids in an Andean cloud forest. *Mycol. Res.*, 110:1257-1270.
- Swarts, N.D., Dixon, K.W., 2009. Terrestrial orchid conservation in the age of extinction. *Ann. Bot.*, 104:543-556.
- Tamura, K., Dudley, J., Nei, M., Kumar, S., 2007. MEGA4: Molecular Evolutionary Genetic Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.*, 24:1596-1599.
- Tao, G., Liu, Z.Y., Hyde, K.D., Lu, X.Z., Yu, Z.N., 2008. Whole rDNA analysis reveals novel and endophytic fungi in *Bletilla ochracea* (Orchidaceae). *Fungal Divers.*, 33:101-122.
- Tarafdar, J.C., Marschner, H., 1994. Efficiency of VAM hyphae in utilisation of organic phosphorus by wheat plants. *Soil. Sci. Plant. Nutr.*, 40: 593-600.
- Taylor, D.L., McCormick, M.K., 2007. Internal transcribes spacer primers and sequences for improved characterization of basidiomycetous orchid mycorrhizas. *New. Phytol.*, 177:1020-1033.

- Trappe, J.M., 1977. Selection of fungi for ectomycorrhizal inoculation in nurseries. *Annu. Rev. Phytopathol.*, 15:203-222.
- Tsavkelova, E.A., Bömke, C., Netrusov, A.I., Weiner, J., Tudzynski, B., 2008. Production of gibberellic acids by an orchid-associated *Fusarium proliferatum* strain. *Fungal Genet. Biol.*, 45:1393-1403.
- Tsavkelova, E.A., Cherdyntseva, T.A., Botina, S.G., Netrusov, A.I., 2007. Bacteria associated with orchid roots and microbial production of auxin. *Microbiol. Res.*, 162:69-76.
- Vacin, E., Went, F., 1949. Some pH changes in nutrient solutions. *Bot. Gaz.*, 110:605-613.
- Vaddhanaphuti, N. (2005). *A Field Guide to The Wild Orchids of Thailand 4<sup>th</sup> ed.* Chiang Mai: Silkworm Books.
- Valois, D., Fayad, K., Barbasubiye, T., Garon, M., Déry, C., Brzezinski, R., Beaulieu, C., 1996. Glucanolytic actinomycetes antagonistic to *Phytophthora fragariae* var. *rubi*, the causal agent of raspberry root rot. *App. Environ. Microbiol.*, 62:1630–1635.
- Verma, A. (1998). *Mycorrhiza Manual*. (pp. 243- 270), Germany: Springer-Verlag Berlin Heidelberg.
- Warcup, J.H., 1973. Symbiotic germination of some Australian terrestrial orchids. *New. Phytol.*, 72:387-392.
- Watanabe, T. (2001). *Pictorial Atlas of Soil and Seed Fungi Morphologies of Cultured Fungi and Key to Species 2<sup>nd</sup> ed.* New York: CRC Press.

- Weber, R.W.S., Webster, J., 2001. Teaching techniques for mycology: 14. mycorrhizal infection of orchid seedlings in the laboratory. *Mycologist.*, 15:55-59.
- Wehner, J., Antunes, P.M., Powell, J.R., Mazukatow, J., Rillig, C., 2009. Plant pathogen protection by arbuscular mycorrhizas: A role for fungal diversity?. *Pedobiologia.*, 53:1970201.
- White, T.J., Bruns, T.D., Lee, S., and Taylor, J.W. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis, M.A., Gelfand, D.H., Sninsky, J.J., and White, T.J. (eds), *PCR Protocols: a Guide to Methods and Applications*. (pp. 315-322), San Diego: Academic.
- Withner, C.L. (1974). *The Orchids, Scientific Studies*. New York: Wiley-Interscience.
- Wright, M., Cross, R., Dixon, K., Huynh, T., Lawrie, A., Nesbitt, L., Prichard, A., Swarts, N., Thomson, R., 2009. Propagation and reintroduction of *Caladenia*. *Aust. J. Bot.*, 57:373-387.
- Yadav, J.S., Tripathi, J.P., 1991. Optimization of cultivation and nutrition conditions and substrate pretreatment for solid-substrate fermentation of wheat straw by *Coriolus versicolor*. *Microbiol.*, 36:294-301.
- Yamanaka, T., 2003. The effect of pH on growth of saprotrophic and ectomycorrhizal ammonia fungi *in vitro*. *Mycologia.*, 95:584-589.
- Yamato, M., Yagame, T., Suzuki, A., Iwase, K., 2005. Isolation and identification of mycorrhizal fungi associating with an achlorophyllus plant, *Epipogium rosesum* (Orchidaceae). *Mycoscience.*, 46:73-77.

- Yoder, J.A., Zettler, L.W., Stewart, S.L., 2000. Water requirements of terrestrial and epiphytic orchid seeds and seedlings, and evidence for water uptake by means of mycotrophy. *Plant Sci.*, 56:145-150.
- Zelmer, C.D., Cuthbertson, L., Currah, R.S., 1996. Fungi associated with terrestrial orchid mycorrhizas, seeds and protocorms. *Mycoscience.*, 37:439-448.
- Zettler, L.W., Hofer, C.J., 1998. Propagation of the little club-spur orchid (*Platanthera clavellata*) by symbiotic seed germination and its ecological implications. *Environ. Exp. Bot.*, 39:189-195.
- Zhu, G.S., Yu, Z.N., Gui, Y., Liu, Z.Y., 2008. A novel technique for isolating orchid mycorrhizal fungi. *Fungal Divers.*, 33:123-137.

## APPENDICES

## APPENDIX A

### MEDIA

Various media were used in this study as isolation medium, storage medium or test medium and prepared as followed formula (per liter).

#### **Carboxymethylcellulose (CMC) agar or Cellulose agar (Kasana *et al.* 2008)**

Sodium nitrate	2	g
Dipotassium hydrogen phosphate	1	g
Magnesium sulfate	0.5	g
Potassium chloride	0.5	g
CMC sodium salt	2	g
Peptone	0.2	g
Agar	15	g
pH	7.0	

The media was sterilized by autoclaved at 121°C for 15 min.

#### **Chrome azurol S agar (Milagres *et al.* 1999)**

##### *Potato dextrose agar (PDA)*

Potato	200	g
Glucose	20	g
Agar	15	g
Distilled water	900	ml
pH	7.0	

Dispensed into containers and sterilized by autoclaved at 121°C for 15 min. Let it cool to 55-60°C. Added 100 ml of CAS sterile solution and mixed well.

*CAS solution*

CAS	60.5	mg
Hexadecyltrimethylammonium	72.9	mg

CAS was dissolved in 50 ml deionized water, and mixed with 10 ml iron (III) solution (1 mM FeCl<sub>6</sub>H<sub>2</sub>O, 10 mM HCl). Under stirring, this solution was slowly mixed with HDTMA dissolved in 40 ml water. The resultant dark blue solution was autoclaved at 121 °C for 15 min.

**Czapek Dox modified agar** (Yamato *et al.* 2005)

Sodium nitrate	0.33	g
Potassium chloride	1	g
Potassium phosphate	0.2	g
Magnesium sulphate	0.2	g
Sucrose	0.5	g
Yeast extract	0.1	g
Agar	15	g
pH	6.8-7.0	

The media was sterilized by autoclaved at 121°C for 15 min.

**Oatmeal agar** (Stewart and Kane, 2006)

Pulverized rolled oat	3	g
Agar	15	g
pH	6.0	

The rolled oat was dissolved with 200 ml of distilled water and heat at 60°C in water bath for 4 hours. Then, the suspension and agar was dissolved in 800 ml of distilled water and sterilized by autoclaved at 121°C for 15 min.

**Potato dextrose agar (PDA)**

Potato	200	g
Glucose	20	g
Agar	15	g
pH	7.0	

**Potato dextrose broth (PDB)**

Potato	200	g
Glucose	20	g
pH	7.0	

The pieces of potato (200 g) were boiled in 500 ml distilled water for 15 – 20 min. Discarded the boiled potato and (added agar, PDA) adjusted volume to 1,000 ml with distilled water (PDB). The media was sterilized by autoclaved at 121°C for 15 min.

**Vacin and Went (VW) agar** (Vacin and Went, 1949)

Tricalcium phosphate	0.2	g
Potassium nitrate	0.525	g
Monopotassium acid phosphate	0.25	g
Magnesium sulphate	0.25	g
Ammonium sulphate	0.5	g
Ferric tartrate	0.028	g
Maganese sulphate	0.75	mg
Sucrose	20	g
Agar	16	g
pH	7.0	

The media was sterilized by autoclaved at 121°C for 15 min.

## APPENDIX B

### CHEMICAL REAGENTS AND BUFFER

#### 10% Cetyltrimethylammonium bromide (CTAB)

CTAB	10	g
0.7 M sodium chloride	100	ml

Added CTAB slowly to NaCl solution at 65 °C and sterilized by autoclaved at 121°C for 15 min.

#### 0.5 M Ethylenediaminetetraacetic acid (EDTA)

Na <sub>2</sub> EDTA.2H <sub>2</sub> O	18.6	g
Sodium hydroxide	2	g
Distilled water	88	ml
pH	8.0	

#### 10X TE buffer

1 M Tris-HCl	10	ml
0.5 M EDTA	2	ml
Distilled water	88	ml
pH	8.0	

**1 M Tris-(hydroxymethyl) aminomethane (Tris-HCl)**

Tris-base	121	g
Conc HCl (37% v/v)	48.3	g
Distilled water	845	ml
pH	8.0	

**0.05% Trypan blue solution**

Trypan blue	5	g
Lactoglycerol	100	ml

*Lactoglycerol*; mixed lactic acid: distilled water: glycerol, 1: 1: 1 (v/v/v).

## APPENDIX C

### GRAIN AND POTTING MEDIA

#### **Grain media** (black bean, corn, kidney bean and sorghum)

Each grain was soaked with water for 4 – 6 hours before transferred into a container (test tube 16 × 150 mm) and sterilized by autoclaved at 121°C for 15 min.

#### **Potting media**

##### *Coconut coir*

Coconut coir were soaked with water for 12 hours before transferred into a container (test tube 16 × 150 mm or flask 250 ml) and sterilized by autoclaved at 121°C for 15 min.

##### *Coconut coir with PDB*

Coconut husk were soaked with PDB for 12 hours before transferred into a container (test tube 16 × 150 mm or flask 250 ml) and sterilized by autoclaved at 121°C for 15 min.

##### *Coconut husk*

Coconut husk were soaked with water for 12 hours before transferred into a container (test tube 16 × 150 mm or flask 250 ml) and sterilized by autoclaved at 121°C for 15 min.

*Coconut husk with PDB*

Coconut husk were soaked with PDB for 12 hours before transferred into a container (test tube 16 × 150 mm or flask 250 ml) and sterilized by autoclaved at 121°C for 15 min.

## APPENDIX D

### STANDARD CRUVE OF INDOLE ACETIC ACID (IAA) PREPARATION

(Gordon and Weber, 1951)

#### Reagent

Salkowski reagent: added 1 ml 0.5 M  $\text{FeCl}_3$  into 50 ml of 35%  $\text{HClO}_4$  and mixed well.

#### Procedure

1. Mixed 1 ml of standard IAA with 2 ml of Salkowski reagent and incubated at room temperature for 30 min in the dark. The appearance of pink color indicated IAA production.

2. The absorbance of the pink pigmentation that developed was measured at 530 nm, uninoculated media mixed with reagent used as a blank.

Table D Absorbance at 530 nm of various concentration of standard IAA

IAA concentration ( $\mu\text{mol/ml}$ )	Absorbance 530 nm
0	0.000
10	0.032
20	0.083
50	0.273
100	0.598
150	0.810
200	0.998




Figure D Standard curve of IAA concentration against absorbance at 530 nm

## APPENDIX E

### DETERMINATION OF HYDROXAMATE-TYPE SIDERPHORE CONCENTRATION BY IRON-PERCHLORATE ASSAY (Atkin *et al.* 1970)

The iron-perchlorate assay is a colorimetric assay used for detection and estimation of hydroxamate-type siderophores under acidic condition.

#### Reagent

Ferric perchlorate solution: containing 5 mM  $\text{Fe}(\text{ClO}_4)_3$  in 0.1 M  $\text{HClO}_4$

#### Procedure

1. Added 0.5 ml of culture supernatant sample to 2.5 ml ferric perchlorate solution and incubated at room temperature for approximately five minutes. The presence of a hydroxamate-type siderophore is shown by the development of an orange-red color.
2. Recorded the absorbance at 480 nm using uninoculated media mixed with reagent as a blank.

Table E Absorbance at 480 nm of various concentration of deferoxamine mesylate by iron-perchlorate assay

Deferoxamine mesylate concentration ( $\mu\text{mol/ml}$ )	Absorbance 480 nm
0	0.000
50	0.158
100	0.344
150	0.473
200	0.667
250	0.827
300	0.976
350	1.132




Figure E Standard curve for hydroxamate concentration determination using deferoxamine mesylate as standard

## CURRICULUM VITAE



**Name** Mr. Ruangwut Chutima

**Date of Birth** 8<sup>th</sup> April 1982

### Education Background

**2003** Bachelor of Science (Microbiology) Chiang Mai University  
 Department of Biology Faculty of Science  
*Special project:* Effect of Crude Extract from some Edible  
 Agarics on Microbail Growth

### Scholarship

**2005-2009** The Strategic Scholarship for frontier Research Network for  
 Join Ph.D. Program and National Research University

**2006** Japan Student Service (JASSO) scholarship, 1 year short-  
 term exchange student

### Plublication

1. Chutima R, Dell B, Vessabutr S, Lumyong S (2011) Endophytic fungi from *Pecteilis susannae* (L.) Rafin (Orchidaceae), a threatened terrestrial orchid in Thailand. *Mycorrhiza* 21: 221-229.
2. Chutima R, Dell B, Lumyong S (2011) Effects of mycorrhizal fungi on symbiotic seed germination of *Pecteilis susannae* (L.) Rafin (Orchidaceae), a terrestrial orchid in Thailand. *Symbiosis* 53: 149-156.

