

LIST OF FIGURES

FIGURE	PAGE
2.1 General diagram flows of processing operation for vegetables and fruits	10
2.2 The interrelationship among the many effects of wounding on physiological processes in fresh-cut produces	20
2.3 Ascorbate-glutathione cycle (Halliwell-Asada pathway) of H ₂ O ₂ scavenging	23
2.4 Localization of reactive oxygen species (ROS) scavenging pathways in plant cells	23
2.5 Structure of ascorbic acid, a main antioxidant present in fruits and vegetables	24
2.6 Proposed mechanism for the degradation of chlorophylls in plant tissues	26
2.7 Structures of the major classes of phenolic acids present in fruits and vegetables	28
2.8 Structures of the major classes of flavonoids present in fruits and vegetables	29
2.9 General structures of glucosinolate	30
2.10 Bioactivation of glucosinolates	31
3.1 Calibration curve with L-(+)-ascorbic acid as standard	53
3.2 Calibration curve with gallic acid as standard	54
3.3 Calibration curve with quercetin as standard	55
3.4 Calibration curve with sinigrin as standard	56
3.5 Calibration curve with trolox as standard	57
3.6 Calibration curve with BSA as standard for Chlase	59
3.7 Calibration curve with BSA as standard for CAT	60
3.8 Calibration curve with BSA as standard for POD	61
3.9 Calibration curve with BSA as standard for SOD	62
3.10 Calibration curve with BSA as standard for APX	63
3.11 Calibration curve with BSA as standard for CAT	64

FIGURE	PAGE
4.1 Changes in microbial populations of total bacteria (A), coliforms (B) and <i>Salmonella-Shigella</i> spp. (C) counts of fresh-cut broccoli florets treated with tap water (control) and hot water (HW) at 50, 55 and 60 °C for 3 min	70
4.2 Changes in L* value (A), a* value (B), b* value (C) and hue angle (h°) (D) color of fresh-cut broccoli florets treated with tap water (control) and hot water (HW) at 50, 55 and 60 °C for 3 min	71
4.3 Changes in the percentage of oxygen (O ₂) (A), carbon dioxide (CO ₂) (B) concentration in package, and respiration rate (mg CO ₂ kg ⁻¹ hr ⁻¹) (C) of fresh-cut broccoli florets treated with tap water (control) and hot water (HW) at 50, 55 and 60 °C for 3 min	72
4.4 Changes in the sensory attributes: overall visual quality (A), visual color (B), and odor (C) score of fresh-cut broccoli florets treated with tap water (control) and hot water (HW) at 50, 55 and 60 °C for 3 min	73
4.5 Changes in microbial populations of total bacteria (A), coliforms (B) <i>Salmonella-Shigella</i> spp. (C), and yeast and mold (D) counts of fresh-cut broccoli florets treated with tap water (control) and vapor heat (VH) treatment at 90 °C for 15, 30 and 45 sec	74
4.6 Changes in L* value (A), a* value (B), b* value (C) and hue angle (h°) (D) color of fresh-cut broccoli florets treated with tap water (control) and vapor heat (VH) treatment at 90 °C for 15, 30 and 45 sec	75
4.7 Changes in percentage of oxygen (O ₂) (A), carbon dioxide (B) concentration in package, and respiration rate (mg CO ₂ kg ⁻¹ hr ⁻¹) (C) of fresh-cut broccoli florets treated with tap water (control) and vapor heat treatment (VH) at 90 °C for 15, 30 and 45 sec	76
4.8 Changes in the sensory attributes: overall visual quality (A), visual color (B), and odor (C) score of fresh-cut broccoli florets treated with tap water (control) and vapor heat (VH) treatment at 90 °C for 15, 30 and 45 sec	77

FIGURE	PAGE
4.9 Changes in microbial populations of total bacteria (A), coliforms (B) <i>Salmonella-Shigella</i> spp. (C), and yeast and mold (D) counts of fresh-cut broccoli florets treated with tap water (control) and ozonated water (O ₃) for 5 min (0.56 ppm), 10 min (1.00 ppm) and 15 min (1.50 ppm)	78
4.10 Changes in L* value (A), a* value (B), b* value (C) and hue angle (h°) (D) color of fresh-cut broccoli florets treated with tap water (control) and ozonated water (O ₃) for 5 min (0.56 ppm), 10 min (1.00 ppm) and 15 min (1.50 ppm)	79
4.11 Changes in total chlorophyll (A), chlorophyll <i>a</i> (B), chlorophyll <i>b</i> (C) and carotenoid (D) content of fresh-cut broccoli florets treated with tap water (control) and ozonated water (O ₃) for 5 min (0.56 ppm), 10 min (1.00 ppm) and 15 min (1.50 ppm)	80
4.12 Changes in the sensory attributes: overall visual quality (A), visual color (B), and odor (C) score of fresh-cut broccoli florets treated with tap water (control) and ozonated water (O ₃) for 5 min (0.56 ppm), 10 min (1.00 ppm) and 15 min (1.50 ppm)	81
4.13 Changes in microbial populations of total bacteria (A), <i>Salmonella-Shigella</i> spp. (B), and yeast and mold (C) counts of fresh-cut broccoli florets with tap water (control) and 500, 750 and 1,000 ppm of sodium chlorite (SC) solutions	83
4.14 Changes in weight loss (A) and respiration rate (B) of fresh-cut broccoli florets treated with tap water (control) and 500, 750 and 1,000 ppm of sodium chlorite (SC) solutions	84
4.15 Changes in L* value (A), a* value (B), b* value (C) and hue angle (h°) (D) color of fresh-cut broccoli florets treated with tap water (control) and 500, 750 and 1,000 ppm of sodium chlorite (SC) solutions	84

FIGURE	PAGE
4.16 Changes in total chlorophyll (A), chlorophyll <i>a</i> (B), chlorophyll <i>b</i> (C) and carotenoid (D) content of fresh-cut broccoli florets treated with tap water (control) and 500, 750 and 1,000 ppm of sodium chlorite (SC) solutions	85
4.17 Changes in the sensory attributes: overall visual quality (A), visual color (B), and odor (C) score of fresh-cut broccoli florets treated with tap water (control) and 500, 750 and 1,000 ppm of sodium chlorite (SC) solutions	86
4.18 Changes in microbial populations of total bacteria (A), coliforms (B), <i>Salmonella-Shigella</i> spp. (C), and yeast and mold (D) counts of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 100 or 300 ppm of sodium chlorite (SC) solutions for 1 min	88
4.19 Changes in L* value (A), a* value (B), b* value (C) and hue angle (h°) (D) color of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 100 or 300 ppm of sodium chlorite (SC) solutions for 1 min	89
4.20 Changes in percentage of weight loss (A), oxygen (O ₂) (B), carbon dioxide (CO ₂) (C) and respiration rate (mg CO ₂ kg ⁻¹ hr ⁻¹) (D) of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 100 or 300 ppm of sodium chlorite (SC) solutions for 1 min	90
4.21 Changes in total chlorophyll (A), chlorophyll <i>a</i> (B), chlorophyll <i>b</i> (C) and carotenoid (D) content of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 100 or 300 ppm of sodium chlorite (SC) solutions for 1 min	91
4.22 Changes in the sensory attributes: overall visual quality (A), color (B), and odor (C) score of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 100 or 300 ppm of sodium chlorite (SC) solutions for 1 min	92

FIGURE	PAGE
4.23 Changes in total ascorbic acid (A) and ascorbic acid (B) content of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 300 ppm of sodium chlorite (SC) solutions for 1 min	94
4.24 Changes in chlorophyll (A) and carotenoid (D) content of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 300 ppm of sodium chlorite (SC) solutions for 1 min	95
4.25 Changes in total phenol (A) and flavonoid (B) content of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 300 ppm of sodium chlorite (SC) solutions for 1 min	96
4.26 Changes in total glucosinolate content of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 300 ppm of sodium chlorite (SC) solutions for 1 min	96
4.27 Changes in antioxidant capacity by %DPPH inhibition of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 300 ppm of sodium chlorite (SC) solutions for 1 min	97
4.28 Changes in chlorophyllase (Chlase) activity of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 300 ppm of sodium chlorite (SC) solutions for 1 min	98
4.29 Changes in catalase (CAT) activity, peroxidase (POD) activity, superoxide dismutase (SOD) activity, ascorbate peroxidase (APX), hydrogen peroxide (H ₂ O ₂) content and glutathione reductase (GR) activity of fresh-cut broccoli florets with tap water (control) and treated with hot water (HW) at 45 °C for 1 min and 300 ppm of sodium chlorite (SC) solutions for 1 min	101