

## เอกสารอ้างอิง

- วีรพล คู่คงวิริยพันธุ์ เคมีป้องกันการก่อมะเร็ง: กลไกการป้องกันของยาและสารจากธรรมชาติ. *ศรีนครินทร์เวชสาร*. 2548; 20(3): 164–173.
- Akihisa T., Yasukawa K., Yamaura M., Ukiya M., Kimura Y., Shimizu N., Arai K. Triterpene alcohol and sterol ferulates from rice bran and their an-inflammatory effects. *J. Agri. Food Chem.* 2000; 48: 2313–2319.
- Bradford M.M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, 1976; 72: 248–254.
- Boonsit P., Pongpiachan P., Julsrigival S., Karladee D. Gamma oryzanol content in glutinous purple rice landrace varieties. *CMU J. Nat. Sci.*, 2010; 9(1): 151–157
- Britz S.J., Prasad P.V.V., Moreau R.A., Hartwell Allen L.JR., Kremer D.F., Boote K.J. Rice Influence of Growth Temperature on the Amounts of Tocopherols, Tocotrienols, and  $\gamma$ -Oryzanol in Brown. *J Agric Food Chem.* 2007; 55: 7559–7565.
- Butsat S. and Siriamornpun, S. Antioxidant capacities and phenolic compounds of the husk, bran and endosperm of Thai rice. *Food Chem.*, 2010; 119(2): 606–613.
- Chen C. and Kong A.N. Dietary Chemopreventive compounds and ARE/EpRE signaling. *Free Radic. Biol. Med.*, 2004; 36: 1505–1516.
- Choi C.W., Kim S.C., Hwang S.S., Choi B.K., Ahn H.J., Lee M.Y. Antioxidant activity and free radical scavenging capacity between Korean medicinal plants and flavonoids by assay-guided comparison. *Plant Sci.*, 2002; 163: 1161–1168.
- Choi S.P., Kim S.P., Kang M.Y., Nam S.H., Friedman M. Protective effects of black rice bran against chemically-induced inflammation of mouse skin. *J. Agri. Food Chem.*, 2010; 58(18): 10007–10015.
- Chun K.S. and Surh Y.J. Signal transduction pathways regulating cyclooxygenase-2 expression: potential molecular targets for chemoprevention. *Biochem. Pharmacol.*, 2004; 68: 1089–1100.
- Das D., Arber N., Jankoski J. Chemoprevention of colorectal cancer. *Digestion.* 2007; 76: 51–67.

- Dewanto V., Wu X.Z., Adom K.K. Liu R.H. Thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity. *J. Agric. Food Chem.*, 2002; 50: 3010–3014.
- Eberhart C.E., Coffey R.J., Radhika A., Giardiello F.M., Ferrenbach S., DuBois R.N. Up-regulation of cyclo-oxygenase 2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology*. 1994; 107: 1183–1188.
- Frame L.T., Ozawa S., Nowell S.A., Chou H.C., Delongchamp R.R., Doerge D.R., Lang N.P., Kadlubar F.F. A simple colorimetric assay for phenotyping the major human thermostable phenol sulfotransferase (SULT1A1) using platelet cytosols. *Drug Metab. Dispos.*, 2000; 28: 1063–1068.
- Gerhardt G.L. and Gallo N.B. Full fat rice bran and oat bran similarly reduced of hypercholesterolemia in humans. *J. Nut.*, 1998; 128: 865–869.
- Gonda T.J. and Metcalf D. Expression of *myb*, *myc* and *fos* proto-oncogenes during the differentiation of a murine myeloid leukemia. *Nature*, 1984; 310: 249–251.
- Guengerich F.P. Common and uncommon cytochrome P450 reactions related to metabolism and chemical toxicity. *Chem. Res. Toxicol.*, 2001; 14: 611–650.
- Harada S., Fujii C., Hayashi A., Ohkoshi N. An association between idiopathic Parkinson's disease and polymorphisms of Phase II detoxification enzymes: Glutathione S-transferase M1 and Quinone oxidoreductase 1 and 2. *Biochem. Bioph. Res. Commun.*, 2001; 288: 887–892.
- Hawk E., Lubert R., Limburg P. Chemoprevention in hereditary colorectal cancer syndrome. *Cancer*. 1999; 86: 2551–2563.
- Hein D.W., Rustan T.D., Doll M.A., Bucker K.D., Ferguson R.J., Feng Y. Acetyltransferase and susceptibility to chemicals. *Toxicol. Letter.*, 1992; 64–65: 123–130.
- Huber W.W., Rossmanith W., Grusch M., Haslinger E., Prustomersky S., Peter-Vorosmarty P., Parzefall W., Scharf G., Schulte-Hermann R. Effects of coffee and its chemopreventive components kahweol and cafestol on cytochrome P450 and sulfotransferase in rat liver. *Food Chem. Toxicol.*, 2008; 46: 1230–1238.
- Ikehashi H. Why are There Indica Type and Japonica Type in Rice — History of the Studies and a View for Origin of Two Types. *Rice Science.*, 2009; 16(1): 1–13.

- Islam M.S., Murata T., Fujisawa M., Nagasaka R., Ushio H., Hori M., Ozaki H. Anti-inflammatory effects of phytosteryl ferulates in colitis induced by dextran sulphate sodium in mice. *Br. J. Pharmacol. Sci.*, 2008; 154(4): 812–824.
- Jang S. and Xu Z. Lipophilic and hydrophilic antioxidants and their antioxidant activities in purple rice bran. *J. Agric. Food Chem.*, 2009; 57: 858–862.
- Jarray R., Allain B., Borriello L., Biard D., Loukaci A., Larghero J., Hadj-Slimane R., Garbay C., Lepelletier Y., Raynaud F. Depletion of the novel protein PHACTR-1 from human endothelial cells abolishes tube formation and induces cell death receptor apoptosis. *Biochimie.*, 2011; 93: 1668–1675.
- Juliano, B.O and Hicks P.A. Rice functional properties and rice food products. *Food Rev. Int.*, 1996; 12: 71–103.
- Kaewkool P. Characterization of cold pressed organic rice bran oil. *As. J. Food Ag-Ind.*, 2011; 4(01): 16–21.
- Kelloff G.J., Boone C.W., Steele V.E., Fay J.R., Lubet R.A., Crowell J.A., Sigman C.C. Mechanistic considerations in chemopreventive drug development. *J. Cell. Biochem. Suppl.*, 1994; 20: 1–24.
- Kikuchi G. and Yochida T. Heme degradation by the microsomal heme oxygenase system. *Trends Biochem. Sci.*, 1998; 5: 323–325.
- Kim D.O., Heo H.J., Kim Y.J., Yang H.S., Lee C.Y. Sweet and sour cherry phenolics and their protective effects on neuronal cells. *J. Agric. Food Chem.*, 2005; 52: 9921–9927.
- Kushi L. and Giovannucci E. Dietary fat and cancer. *Am. J. Clin. Nutr.*, 2002; 113: 63–70.
- Kwak D.W., Wakabayashi N., Kensler T.W. Chemoprevention through the keap1-Nrf2 signaling pathway by phase 2 enzyme inducer. *Mutation Res.*, 2004; 555: 133–148.
- Lai P., Li K.Y., Lu S., Chen H.H. Phytochemicals and antioxidant properties of solvent extracts from *Japonica* rice bran. *Food Chem.*, 2009; 117: 538–544
- Land H., Parada L.F., Weinberg R.A. Cellular oncogenes and multistep carcinogenesis. *Science*, 1983; 222: 771–778.

- Lodish H., Berk A., Zipursky S.L. Molecular Cell Biology 4<sup>th</sup> Edition: Proto-oncogenes and tumor-suppressor genes (Section 24.2). New York: W.H. Freeman; 2000
- Lubet R.A., Steele V.E., DeCoster R., Bowden C., You M., Juliana M.M., Eto I., Kelloff G.J., Grubbs C.J. Chemopreventive effects of the aromatase inhibitor vorozole (R 83842) in the methylnitrosourea-induced mammary cancer model. *Carcinogenesis*, 1998; 19(8): 1345–1351.
- Nagendra Prasad M.N., Sanjay K.R., Shravya Khatokar M., Vismaya M.N., Nanjunda Swamy S. Health benefit of rice bran– a review. *Nutr. Food Sci.*, 2011; 1(3): 1–7.
- Nguyen T., Brunsan D., Crespi C.L., Penman B.W., Wishnork J.S., Tannenbaum S.R. DNA damage and mutation in human cells exposed to nitric oxide *in vitro*. *Proc. Natl. Acad. Sci.*, 1992; 89: 3030–3034.
- Qureshi A.A., Mo H., Packer L., Peterson D.M. Isolation and identification of novel tocotrienols from rice bran with hypocholesterolemic, antioxidant and antitumor properties. *J. Agri. Food Chem.*, 2000; 48: 3130–3140.
- Saenjum C., Chaiyasut C., Kadchumsang S., Chansakaow S., Suttajit S., Antioxidant activity and protective effects on DNA damage of *Caesalpinia sappan* L. extract. *J. Med. Plants Res.*, 2010; 4: 1594–1600.
- Seetharamaiah G.S. and Prabhakar J.V. Oryzanol content of Indian rice bran oil and its extraction from soap stock. *J. Food Sci. Technol.*, 1986; 23: 270–273.
- Seetharamaiah G.S., Krishnakantha T.P., Chandrasekhara N. Influence of oryzanol on platelet aggregation in rats. *J. Nutr. Sci. Vitaminol.*, 1990; 36: 291–297.
- Seitz L.M. Stanol and sterol esters of ferulic and *p*-coumaric acids in wheat, corn, rye, and triticale. *J. Agric. Food Chem.*, 1989; 37: 662–667.
- Sheehan K.M., Sheahan K., O'Donoghue D.P., MacSweeney F., Conroy R.M., Fitzgerald D.J. The relationship between cyclooxygenase–2 expression and colorectal cancer. *JAMA*. 1999; 282: 1254–1257.
- Stuehr D.J. and Nathan C.F. Nitric oxide: a macrophage product responsible for cytostasis and respiratory inhibition in tumor target cells. *J. Exp. Med.*, 1989; 169(5): 1543–1555.

- Surh Y.J. Cancer chemoprevention with dietary phytochemicals. *Nat. Rev. Cancer.*, 2003; 3: 768–780.
- Surh Y.J., Han S.S., Keum Y.S., Seo H.J., Lee S.S. Inhibitory effects of curcumin and capsaicin on phorbol ester–induced activation of eukaryotic transcription factors, NF–kappaB and AP–1. *Biofactors.* 2000; 12: 107–112.
- Szabo C., Ischiropoulos H., Radi R. Peroxynitrite: Biochemistry, pathophysiology and development of therapeutics. *Nat. Rev. Drug Discov.*, 2007; 6: 662–680.
- Tsao A.S., Kim E.S., Hong W.K. Chemoprevention of cancer. *CA Cancer J. Clin.*, 2004; 54: 150–180.
- Verschoye R.D., Greaves P., Cai H., Edwards R.E., Steward W.P., Gescher A.J. Evaluation of the cancer chemopreventive efficacy of rice bran in genetic mouse models of breast, prostate and intestinal carcinogenesis. *British J. of Cancer.*, 2007; 96: 248–254.
- Xu Z. and Godber J.S. Purification and identification of components of gamma–oryzanol in rice bran oil. *J. Agri. Food Chem.*, 1999; 47: 2724–2728.
- Xu Z., Godber J.S., Xu Z. Antioxidant activities of major components of gamma–oryzanol from rice bran using a linoleic acid model. *J. Am. Oil Chem. Soc.*, 2001; 78: 465–469.
- Yanping L., Yanhua L., Dongzhi W. Antioxidant activity of a flavonoid–rich extract of *Hypericum perforatum* L in vitro. *J. Agric. Food Chem.*, 2004; 52: 5032–5029.



