

LIST OF FIGURES

FIGURE	PAGE
2.1 Vegetable production (including melons) in the top ten producing countries and world (2007)	5
2.2 Anatomy of select leafy vegetables	6
2.3 Cabbage growth stage	11
2.4 Good quality of cabbage head	12
2.5 Schematic illustration of dropping treatment of cabbage at 80, 40, 20, and 10 cm	22
2.6 Injuries caused by dropping treatment from the high of 80 cm: the first layer (A) and the fourth layer (B) of the head leaf pack of cabbage	22
2.7 Grades and standards for cabbage under Royal Project	23
2.8 The leafy vegetable supply chain in Shanghai, China	29
2.9 A schematic flow chart of the cold chain	36
3.1 The location of study areas (Chiang Mai and Phetchabun provinces, Thailand)	42
3.2 Preparation for transport from farm to retail market	44
3.3 The sketch of positioning of cabbages on the truck during transport from farms in Phetchabun province to wholesale and retail markets in Bangkok, Thailand	45
3.4 The numbers of cabbage leaves were divided in to 4 groups	47
4.1 Practices of cabbage supply chain in Phetchabun province, Thailand	52
4.2 Practices of cabbage supply chain in Royal Project Foundation, Chiang Mai province, Thailand	53
4.3 The cabbage supply chain in Phetchabun province (traditional chain)	54
4.4 The cabbage supply chain in Royal Project Foundation (good practice chain)	55
4.5 Temperature (A) and relative humidity (B) of cabbages during transport from farms in Phetchabun province (1) to wholesale (2) and retail market (3) in Bangkok, Thailand	72
4.6 Severity of physical damage of cabbages with different numbers of wrapper leaves during transportation from farms in Phetchabun Province (1), to wholesale market (2) and retail markets (3) in Bangkok, Thailand	73
4.7 Physical damage of cabbages with different numbers of wrapper leaves	74
4.8 Cabbage marketing channel in Phetchabun province, Thailand	84
4.9 Total soluble solids (TSS) content in different leaves of cabbage	90
4.10 Total chlorophyll contents in various leaves of cabbage	90
4.11 Carotenoids content in different leaves of cabbage	91
4.12 Total ascorbic acid contents in different leaves of cabbage	91

	PAGE	
4.13	Price of cabbage at the collection center of Phetchabun provincial market, Thailand, March 2010	92
4.14	Weight loss (A) and firmness (B) changes in cabbages during storage at 4, 10 and 28 °C	93
4.15	Respiration rate (A) and ethylene production (B) of cabbages during storage at 4, 10 and 28 °C	94
4.16	Yellowing (A), L^* (B), b^* values (C) and chlorophyll content (D) of cabbage during storage at 4, 10 and 28 °C	95
4.17	Appearance of top (A) and bottom (B) parts of cabbage heads stored at 4, 10 and 28 °C	96
4.18	Total soluble solids (A) and ascorbic acid content (B) of cabbages during storage at 4, 10 and 28 °C	97
4.19	Storage life of cabbages at 4, 10 and 28 °C	98
5.1	Recommended models to produce a good quality of cabbage in the supply chain	101