

REFERENCES

- Anand, P., Kunnumakara, A., Sundaram, C., Harikumar, K., Tharakan, S., Lai, O., Sung, B., and Aggarwal, B. (2008). Cancer is a preventable disease that requires major lifestyle changes. *Pharm. Res.*, 25(9): 2097–2116.
- Anzenbacher, P. and Anzenbacherová, E. (2001). Cytochromes P450 and metabolism of xenobiotics. *Cell Mol. Life Sci.*, 58(5-6): 737-47.
- Arsura, M. and Cavin, L.G. (2005). Nuclear factor-kappaB and liver carcinogenesis. *Cancer Lett.*, 229(2): 157-69.
- Benson, A.M., Hunkeler, M.J. and Talalay, P. (1980). Increase of NAD(P)H:quinone reductase by dietary antioxidants: possible role in protection against carcinogenesis and toxicity. *Proc. Natl. Acad. Sci.*, 77(9): 5216-20.
- Bjeldanes, L.F. and Chang, G.W. (1977). Mutagenic activity of quercetin and related compounds. *Science*, 197(4303): 577-8.
- Bolton, J.L., Trush, M.A., Penning, T.M., Dryhurst, G. and Monks, T.J. (2000). Role of quinones in toxicology. *Chem. Res. Toxicol.*, 13(3): 135-60.
- Boonsit, P., Pongpiachan, P., Julsrigival, S., and Karladee, D. (2010). Gamma oryzanol content in glutinous purple rice landrace varieties. *CMU Journal of Natural Science*, 9: 151-157
- Breinholt, V., Lauridsen, S.T., Daneshvar, B., and Jakobsen, J. (2000). Dose-response effects of lycopene on selected drug-metabolizing and antioxidant enzymes in the rat. *Cancer Lett.*, 154(2): 201-10.
- Buck, E. A. (2001). The effect of the flavonoid quercetin on phase 1 and phase 2 enzyme activities in the SW-480 human colon carcinoma cell line. M.S. Thesis, The University of Florida.
- Butsat, S. and Siriamornpun, S. (2010). Phenolic acids and antioxidant activities in husk of different thai rice varieties. *Food Sci. Technol. Int.*, 16(4):329-36.
- Cao, P., Vadhanam, M.V., Spencer, W.A., Cai, J. and Gupta, R.C. (2011). The antioxidant potency of *Punica granatum L.* fruit peel reduces cell proliferation and induces apoptosis on breast cancer. *J. Med. Food*, 14(12): 1638-46.

- Cabibbo, G., Maida, M., Genco, C., Antonucci, M., and Cammà, C. (2012). Causes of and prevention strategies for hepatocellular carcinoma. *Semin. Oncol.* 39(4): 374-83.
- Chen, C. and Kong, A.N. (2004). Dietary chemopreventive compounds and ARE/EpRE signaling. *Free Radic. Biol. Med.*, 36(12): 1505-16.
- Chung, Y.C., Lin, C.C., Chou, C.C., and Hsu, C.P. (2010). The effect of Longan seed polyphenols on colorectal carcinoma cells. *Eur. J. Clin. Invest.*, 40(8): 713-21.
- Charoensin, S., Punvittayagul C., Pompimon, W., Mevatee, U. and Wongpoomchai R. (2010). Toxicological and clastogenic evaluation of pinocembrin and pinostrobin isolated from *Boesenbergia pandurata* in Wistar rats. *Thai. J. Toxicology*, 25(1): 29-40.
- Cliet, I., Fournier, E., Melcion, C., and Cordier, A. (1989). *In vivo* micronucleus test using mouse hepatocytes. *Mutat. Res.*, 216(6): 321-6.
- Dadkhah, A., Allameh, A., Khalafi, H., and Ashrafihelan J. (2011). Inhibitory effects of dietary caraway essential oils on 1,2-dimethylhydrazine-induced colon carcinogenesis is mediated by liver xenobiotic metabolizing enzymes. *Nutr. Cancer*, 63(1): 46-54.
- Daiponmaka, W., Theerakulpisutb, P., Thanonkaoc, P., Vanavichitd, A., and Prathephaa, P. (2010). Changes of anthocyanin cyanidin-3-glucoside content and antioxidant activity in Thai rice varieties under salinity stress. *ScienceAsia*, 36: 286-291.
- Dinkova-Kostova, A.T., Holtzclaw, W.D., Cole, R.N., Itoh, K., Wakabayashi, N., Katoh, Y., Yamamoto, M. and Talalay, P. (2002). Direct evidence that sulfhydryl groups of Keap1 are the sensors regulating induction of phase 2 enzymes that protect against carcinogens and oxidants. *Proc. Natl. Acad. Sci. U.S.A.*, 99(18): 11908-13.
- Doostdar, H., Burke, M.D., and Mayer, R.T. (2000). Bioflavonoids: selective substrates and inhibitors for cytochrome P450 CYP1A and CYP1B1. *Toxicology*, 144(1-3): 31-8

- Fenech, M., Kirsch-Volders, M., Natarajan, A.T., Surralles, J., Crott, J.W., Parry, J., Norppa, H., Eastmond, D.A., Tucker, J.D. and Thomas, P. (2011). Molecular mechanisms of micronucleus, nucleoplasmic bridge and nuclear bud formation in mammalian and human cells. *Mutagenesis*, 26(1): 125-32
- Ferguson, L.R., Philpott, M., and Karunasinghe, N. (2004). Dietary cancer and prevention using antimutagens. *Toxicology*, 198(1-3): 147-59.
- Ferreira, J.F., Luthria, D.L., Sasaki, T., and Heyerick, A. (2010). Flavonoids from *Artemisia annua* L. as antioxidants and their potential synergism with artemisinin against malaria and cancer. *Molecules*, 15(5): 3135-70.
- Fiddler (1975). The occurrence and determination of N-nitroso compounds. *Toxicol. Appl. Pharmacol.*, 31(3): 352-60.
- Fisher, M.B., Paine, M.F., Strelevitz, T.J. and Wrighton, S.A. (2001). The role of hepatic and extrahepatic UDP-glucuronosyltransferases in human drug metabolism. *Drug Metab. Rev.*, 33(3-4): 273-97.
- Goldman, R. and Shields, P.G. (2003). Food mutagens. *J. Nutr.*, 133 Suppl 3: 965S-973S.
- Gozzelino, R., Jeney, V., and Soares, M.P. (2010). Mechanisms of cell protection by heme oxygenase-1. *Annu. Rev. Pharmacol. Toxicol.*, 50: 323-54.
- Habig, W.H. and Jakoby, W.B. (1981). Assays for differentiation of glutathione S-transferases. *Methods Enzymol.*, 77: 398-405.
- Hayes, J.D. and McMahon, M. (2001). Molecular basis for the contribution of the antioxidant responsive element to cancer chemoprevention. *Cancer Lett.*, 174(2): 103-13.
- Higgins, G.M. and Anderson, R.M. (1931). Experimental pathology of the liver. I. Restoration of the liver of the white rat following partial surgical removal. *Arch. Pathol.*, 12: 186-202.
- Huang, S.H. and Ng, L.T. (2011). Quantification of tocopherols, tocotrienols, and gamma-oryzanol contents and their distribution in some commercial rice varieties in Taiwan. *J. Agric. Food Chem.*, 59(20): 11150-9
- Hui, C., Bin, Y., Xiaoping, Y., Long, Y., Chunye, C., Mantian, M. and Wenhua, L. (2010). Anticancer activities of an anthocyanin-rich extract from black rice against breast cancer cells *in vitro* and *in vivo*. *Nutr. Cancer*, 62(8): 1128-36.

- Ip, C. and Ganther, H.E. (1991). Combination of blocking agents and suppressing agents in cancer prevention. *Carcinogenesis*, 12(2): 365-7.
- Ito, N., Imaida, K., Asamoto, M. and Shirai, T. (2000). Early detection of carcinogenic substances and modifiers in rats. *Mutat. Res.*, 462(2-3): 209-17.
- Jancova, P., Anzenbacher, P. and Anzenbacherova, E. (2010). Phase II drug metabolizing enzymes. *Biomed. Pap. Med. Fac. Univ. Palacky. Olomouc. Czech. Repub.*, 154(2): 103-16.
- Jeon, K.I., Park, E., Park, H.R., Jeon, Y.J., Cha, S.H. and Lee, S.C. (2006). Antioxidant activity of far-infrared radiated rice hull extracts on reactive oxygen species scavenging and oxidative DNA damage in human lymphocytes. *J. Med. Food*, 9(1): 42-8.
- Johnson, I.T. (2002). Glucosinolates: bioavailability and importance to health. *Int. J. Vitam. Nutr. Res.*, 72(1): 26-31.
- Johnson, I.T., Williamson, G., and Musk, S.R. (1994). Anticarcinogenic factors in plant foods: a new class of nutrients?. *Nutr. Res. Rev.*, 7(1): 175-204.
- Jozkowicz, A., Was, H. and Dulak, J. (2007). Heme oxygenase-1 in tumors: is it a false friend?. *Antioxid. Redox Signal.*, 9(12): 2099-117.
- Karladee, D. and Suriyong, S. (2012). γ -Aminobutyric acid (GABA) content in different varieties of brown rice during germination. *ScienceAsia*, 38: 13-17.
- Klaassen, C.D., and Watkins, J.B. (2010). Casarett & Doull's Essentials of Toxicology. New York : McGraw-Hill Professional.
- Kennedy, C.H., Mitchell, C.E., Fukushima, N.H., Neft, R.E. and Lechner, J.F. (1996). Induction of genomic instability in normal human bronchial epithelial cells by ²³⁸Pu alpha-particles. *Carcinogenesis*, 17(8):1671-6.
- Kong, A.N., Yu, R., Chen, C., Mandlekar, S., and Primiano, T. (2000). Signal transduction events elicited by natural products: role of MAPK and caspase pathways in homeostatic response and induction of apoptosis. *Arch. Pharm. Res.* 23(1):1-16.
- Kong, C.S., Jeong, C.H., Choi, J.S., Kim, K.J. and Jeong, J.W. (2013). Antiangiogenic effects of p-coumaric Acid in human endothelial cells. *Phytother. Res.*, 27(3): 317-23.

- Krajka-Kuźniak, V., Szafer, H. and Baer-Dubowska W. (2005). Modulation of cytochrome P450 and phase II enzymes by protocatechuic acid in mouse liver and kidney. *Toxicology*. 216(1): 24-31.
- Krajka-Kuźniak, V., Kaczmarek, J. and Baer-Dubowska, W. (2008). Effect of naturally occurring phenolic acids on the expression of glutathione S-transferase isozymes in the rat. *Food Chem. Toxicol.* 46(3): 1097-102.
- Kwak, M.K., Egner, P.A., Dolan, P.M., Ramos-Gomez, M., Groopman, J.D., Itoh, K., Yamamoto, M. and Kensler, T.W. (2001). Role of phase 2 enzyme induction in chemoprotection by dithiolethiones. *Mutat. Res.* 48(81): 305-15.
- Lee, J., Robert, W., Durst, W., and Ronald, E. (2005). Determination of total monomeric anthocyanin pigment content of fruit juices, beverages, natural colorants, and wines by the pH differential method: collaborative study. *J. AOAC Int.* 88(5): 1269-1278.
- Lin, X.F., Min, W. and Luo, D. (2010). Anticarcinogenic effect of ferulic acid on ultraviolet-B irradiated human keratinocyte HaCaT cells. *J. Med. Plant. Res.*, 4(16): 1686-1694.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951). Protein measurement with the Folin phenol reagent. *J. Biol. Chem.*, 193(1): 265-75.
- Ma, Y.T. and Cheung, P.C. (2007). Spectrophotometric determination of phenolic compounds by enzymatic and chemical methods a comparison of structure activity relationship. *J. Agric. Food Chem.*, 55(10): 4222-8.
- Maeda, S., Kamata, H., Luo, J.L., Leffert, H. and Karin, M. (2005). IKKbeta couples hepatocyte death to cytokine-driven compensatory proliferation that promotes chemical hepatocarcinogenesis. *Cell*, 121(7): 977-90.
- Majewska-Wierzbicka, M., and Czczot, H. (2012). Anticancer activity of flavonoids. *Pol. Merkur. Lekarski.* 33(198): 364-9.
- Maksimović, Z., Malencić, D. and Kovacević, N. (2005). Polyphenol contents and antioxidant activity of *Maydis stigma* extracts. *Bioresour. Technol.* 96(8): 873-7.
- Martins, P.N., Theruvath, T.P. and Neuhaus, P. (2008). Rodent models of partial hepatectomies. *Liver Int.*, 28(1): 3-11.

- Miao, W., Hu ,L., Scrivens, P.J. and Batist, G. (2005). Transcriptional regulation of NF-E2 p45-related factor (NRF2) expression by the aryl hydrocarbon receptor-xenobiotic response element signaling pathway: direct cross-talk between phase I and II drug-metabolizing enzymes. *J. Biol. Chem.*, 280(21): 20340-8.
- Mizutani, M. and Ohta, D. (1998). Two isoforms of NADPH: cytochrome P450 reductase in *Arabidopsis thaliana*: Gene structure, heterologous expression in insect cells, and differential regulation. *Plant Physiol.*, 116(1):357-67.
- Monks, T. J. and Jones, D. C. (2002). The metabolism and toxicity of quinones, quinonimines, quinone methides, and quinone-thioethers. *Curr. Drug. Metab.*, 3(4): 425-38.
- Pray, L. (2008). DNA replication and causes of mutation. *Nature Education*, 1(1):1-6
- Punyatong, M., Pongpiachan, P., Pongpiachan, P., Karladee, D. and Mankhetkorn, S. (2008). Cytotoxicity of crude proanthocyanidin extract from purple glutinous rice bran (*Oryza sativa* L.) (Kum Doi Saket) compared with cyanidin 3-glucoside on X63 myeloma cancer cell lines. *Kasetsart J. (Nat. Sci.)*, 42: 676-681.
- Morita, T., MacGregor, J.T. and Hayashi, M. (2011). Micronucleus assays in rodent tissues other than bone marrow. *Mutagenesis*, 26(1): 223-30.
- Müller-Tegethoff, K., Kasper, P. and Müller, L. (1995). Evaluation studies on the *in vitro* rat hepatocyte micronucleus assay. *Mutat. Res.*, 335(3): 293-307.
- Nair, S., Li, W. and Kong, AN. (2007). Natural dietary anti-cancer chemopreventive compounds: redox-mediated differential signaling mechanisms in cytoprotection of normal cells versus cytotoxicity in tumor cells. *Acta. Pharmacol. Sin.*, 28(4): 459-72.
- Narumi, K., Ashizawa, K., Takashima, R., Takasawa, H., Katayama, S., Tsuzuki, Y., Tatemoto, H., Morita, T., Hayashi, M. and Hamada, S. (2012). Development of a repeated-dose liver micronucleus assay using adult rats: an investigation of diethylnitrosamine and 2,4-diaminotoluene. *Mutat. Res.*, 747(2): 234-9.

- Nioi, P. and Hayes, J.D. (2004). Contribution of NAD(P)H: quinone oxidoreductase 1 to protection against carcinogenesis, and regulation of its gene by the Nrf2 basic-region leucine zipper and the arylhydrocarbon receptor basic helix-loop-helix transcription factors. *Mutat. Res.*, 555(1-2): 149-71.
- Nishimura, R., Tabata, K., Arakawa, M., Ito, Y., Kimura, Y., Akihisa, T., Nagai, H., Sakuma, A., Kohno, H. and Suzuki, T. (2007). Isobavachalcone, a chalcone constituent of *Angelica keiskei*, induces apoptosis in neuroblastoma. *Biol. Pharm. Bull.*, 30(10): 1878-83.
- OECD. (2008). OECD guidelines for the testing of chemicals (Acute oral toxicity – up and down procedure).
- Oh, C.H. and Oh, S.H. (2004). Effects of germinated brown rice extracts with enhanced levels of GABA on cancer cell proliferation and apoptosis. *J. Med. Food*, 7(1):19-23.
- Oliveira, P.A., Colaço, A., Chaves, R., Guedes-Pinto, H., De-La-Cruz P, L.F. and Lopes, C. (2007). Chemical carcinogenesis. *An Acad. Bras. Cienc.*, 79(4): 593-616.
- Okamoto T. (2005). Safety of quercetin for clinical application (Review). *Int. J. Mol. Med.*, 16(2):275-8.
- Omiecinski, C.J., Rimmel, R.P. and Hosagrahara, V.P. (1999). Concise review of the cytochrome P450s and their roles in toxicology. *Toxicol. Sci.*, 48(2): 151-6.
- Owuor, E. D. and Kong A. N. (2002). Antioxidants and oxidants regulated signal transduction pathways. *Biochem. Pharmacol.*, 64(5-6): 765-70.
- Patterson, A.D., Gonzalez, F.J. and Idle, J.R. (2010). Xenobiotic metabolism: a view through the metabolometer. *Chem. Res. Toxicol.*, 23(5): 851-60.
- Pitot, H.C. (1989). Progression: The terminal stage of carcinogenesis. *Jpn. J. Cancer Res.*, 80: 599-607.
- Pool-Zobel, B., Veeriah, S. and Böhmer, F.D. (2005). Modulation of xenobiotic metabolising enzymes by anticarcinogens focus on glutathione S-transferases and their role as targets of dietary chemoprevention in colorectal carcinogenesis. *Mutat. Res.*, 591(1-2): 74-92.

- Prasad, K.N., Yang, B., Shi, J., Yu, C., Zhao, M., Xue, S. and Jiang, Y. (2010). Enhanced antioxidant and antityrosinase activities of longan fruit pericarp by ultra-high-pressure-assisted extraction. *J. Pharm. Biomed. Anal.*, 51(2): 471-7.
- Punvittayagul C., Kanokwan S., Karladee D., Wongpoomchai R. (2011). Mutagenic study of various extracts from black glutinous rice (*Oryza sativa* L. var. *glutinosa*) by using Salmonella mutation assay. *The 3rd Biochemistry and Molecular Biology conference manual*, 2011: 214.
- Reagan-Shaw, S., Eggert, D., Mukhtar, H. and Ahmad, N. (2010). Antiproliferative effects of apple peel extract against cancer cells. *Nutr. Cancer*, 62(4): 517-24.
- Rietjens, I.M., Boersma, M.G., van der Woude, H., Jeurissen, S.M., Schutte, M.E. and Alink, G.M. (2005). Flavonoids and alkenylbenzenes: mechanisms of mutagenic action and carcinogenic risk. *Mutat. Res.*, 574(1-2): 124-38.
- Rueff, J., Laires, A., Gaspar, J. and Rodrigues A. (1990). Mutagenic activity in the wine-making process: correlations with rutin and quercetin levels. *Mutagenesis*. 5(4): 393-6.
- Sangkitikomol, W., Tencomnaol, T., and Rocejanasaroj, A. (2010). Effects of Thai black sticky rice extract on oxidative stress and lipid metabolism gene expression in HepG2 cells. *Genet. Mol. Res.*, 9(4): 2086-2095.
- Sheehan, D., Meade, G., Foley, V.M. and Dowd, C.A. (2001). Structure, function and evolution of glutathione transferases: implications for classification of non-mammalian members of an ancient enzyme superfamily. *Biochem. J.*, 360 (Pt 1): 1-16.
- Sheweita, S.A., and Tilmisany, A.K. (2003). Cancer and phase II drug-metabolizing enzymes. *Curr. Drug Metab.*, 4(1): 45-58.
- Shih, P.H., Yeh, C.T., and Yen, G.C. (2007). Anthocyanins induce the activation of phase II enzymes through the antioxidant response element pathway against oxidative stress-induced apoptosis. *J. Agric. Food Chem.*, 55(23): 9427-35.
- Sogawa, K. and Fujii-Kuriyama, Y. (1997). Ah receptor, a novel ligand-activated transcription factor. *J. Biochem.*, 122(6): 1075-9.
- Srivastava, A., Akoh, C.C., Fischer, J., and Krewer, G. (2007). Effect of anthocyanin fractions from selected cultivars of Georgia-grown blueberries on apoptosis and phase II enzymes. *J. Agric. Food Chem.*, 55(8): 3180-5.

- Surh, Y.J. (2003). Cancer chemoprevention with dietary phytochemicals. *Nat. Rev. Cancer*, 3(10): 768-80.
- Sutandyo, N. (2010). Nutritional carcinogenesis. *Acta. Med. Indones.*, 42(1): 36-42.
- Tan, X.L. and Spivack, S.D. (2009). Dietary chemoprevention strategies for induction of phase II xenobiotic-metabolizing enzymes in lung carcinogenesis: A review. *Lung Cancer*, 65(2): 129-37.
- Talalay, P., Fahey, J.W., Holtzclaw, W.D., Prester, T., and Zhang, Y. (1995). Chemoprotection against cancer by phase 2 enzyme induction. *Toxicol. Lett.*, 82-83:173-9.
- Tenhunen, R., Marver, H.S., and Schmid, R. (1968). The enzymatic conversion of heme to bilirubin by microsomal heme oxygenase. *Proc. Natl. Acad. Sci. U. S. A.*, 61: 748-755.
- Terradas, M., Martín, M., Tusell, L and Genesca, A. (2010). Genetic activities in micronuclei: is the DNA entrapped in micronuclei lost for the cell?. *Mutat Res.*, 705(1): 60-7.
- Tsuda, H., Ohshima, Y., Nomoto, H., Fujita, K., Matsuda, E., Iigo, M., Takasuka, N. and Moore, M.A. (2004). Cancer prevention by natural compounds. *Drug Metab. Pharmacokinet.*, 19(4): 245-63.
- Uda, Y., Price, K.R., Williamson, G. and Rhodes, M.J. (1997). Induction of the anticarcinogenic marker enzyme, quinone reductase, in murine hepatoma cells in vitro by flavonoids. *Cancer Lett.*, 120(2): 213-6.
- Utesch, D., Feige, K., Dasenbrock, J., Broschard, T.H., Harwood, M., Danielewska-Nikiel, B. and Lines, T.C. (2008). Evaluation of the potential in vivo genotoxicity of quercetin. *Mutat. Res.*, 654(1): 38-44.
- Vakkila, J. and Lotze, M. T. (2004). Inflammation and necrosis promote tumour growth. *Nat. Rev. Immunol.*, 4(8): 641-8.
- Van Bladeren, P. J. (2000). Glutathione conjugation as a bioactivation reaction. *Chem. Biol. Interact.*, 129(1-2): 61-76.
- Vermilion, J.L., Ballou, D.P., Massey, V. and Coon, M.J. (1981). Separate roles for FMN and FAD in catalysis by liver microsomal NADPH-cytochrome P-450 reductase. *J. Biol. Chem.*, 256(1): 266-77.

- Verna, L., Whysner, J. and Williams, G.M. (1996). N-nitrosodiethylamine mechanistic data and risk assessment: bioactivation, DNA-adduct formation, mutagenicity, and tumor initiation. *Pharmacol. Ther.*, 71(1-2): 57-81.
- Vichaponga, J., Sookserma, M., Srijesdaruk, V., Swatsitangc, P. and Srijaranai S. (2010). High performance liquid chromatographic analysis of phenolic compounds and their antioxidant activities in rice varieties. *LWT. Food Science and Technology International*, 43: 1325e1330.
- Wells, P.G., Mackenzie, P.I., Chowdhury, J.R., Guillemette, C., Gregory, P.A., Ishii, Y., Hansen, A.J., Kessler, F.K., Kim, P.M., Chowdhury, N.R. and Ritter, J.K. (2004). Glucuronidation and the UDP-glucuronosyltransferases in health and disease. *Drug. Metab. Dispos.*, 32(3): 281-90.
- Williams, G. M. (2001). Mechanisms of chemical carcinogenesis and application to human cancer risk assessment. *Toxicology*, 166(1-2): 3-10.
- Yang, L., Browning, J.D. and Awika, J.M. (2009). Sorghum 3-deoxyanthocyanins possess strong phase II enzyme inducer activity and cancer cell growth inhibition properties. *J. Agric. Food Chem.*, 57(5): 1797-804.
- Yang, C.S., Smith, T.J. and Hong, J.Y. (1994). Cytochrome P-450 enzymes as targets for chemoprevention against chemical carcinogenesis and toxicity: opportunities and limitations. *Cancer Res.*, 54(7 Suppl): 1982s-1986s.
- Yoshimura, Y., Zaima, N., Moriyama, T. and Kawamura, Y. (2012). Different localization patterns of anthocyanin species in the pericarp of black rice revealed by imaging mass spectrometry. *PLoS One*, 7(2): e31285.
- Zhu, B.T., Suchar, L.A., Gwynne, M. and Huang, M.T. (1995). Effect of age and sex on the inducibility of hepatic microsomal 4-nitrophenol UDP-glucuronyltransferase activity in rats. *Life Sci.*, 57(1): PL19-24.
- Zhu, X., Fan, W.G., Li, D.P., Lin, M.C. and Kung, H. (2010). Heme oxygenase-1 system and gastrointestinal tumors. *World J. Gastroenterol.*, 16(21): 2633-7.