

## LIST OF FIGURES

FIGURE	PAGE
2.1	Shrimp life cycle <span style="float: right;">6</span>
2.2	Bacterially mediated processes in a shrimp culture system <span style="float: right;">10</span>
2.3	Outline of the cloning procedure for studying a microbial community <span style="float: right;">17</span>
2.4	Schematic representation of DGGE <span style="float: right;">18</span>
3.1	(A) Experimental set up of 35 L of plastic tank with Set 1: shrimp culture (control); Set 2: shrimp culture with <i>Spirulina</i> mat; Set 3: shrimp culture with artificial mat, respectively. (B) Top view of the closed-recirculation system in 35 L plastic tank <span style="float: right;">19</span>
3.2	(A) Fibrous polyester mats; (B) <i>Spirulina</i> mat immobilization <span style="float: right;">21</span>
3.3	(A) Filter medium; (B) Filtration systems for suspended removal in closed-recirculation system during shrimp cultivation <span style="float: right;">22</span>
3.4	Outline procedures to analyze microbial community between samples from hatchery and grow-out pond <span style="float: right;">25</span>
4.1	Microbial community in water column of without mat condition. CW0-CW10 was microbial community in water column from day 0 to day 10 that reared shrimp from PL2 to PL12 in without <i>Spirulina</i> mat condition. The arrow shows the water exchanging and the water samples were taken before water exchanging <span style="float: right;">31</span>
4.2	16S rDNA-based denaturing gradient gel electrophoresis (DGGE) profiles (Left) with cluster analysis or percentage of Nei and Li's similarity coefficients[87](Right) of the bacterial community in water column of Set 1. Lane CW0 to CW10 was bacteria community in water column in Set 1 <span style="float: right;">32</span>
4.3	Microbial community in water column of Set 2 (co-cultured between <i>Spirulina</i> mat and shrimp). WH0-WH10 was microbial community in water column from day 0 to day 10 that reared shrimp from PL2 to PL12 in co-culturing with <i>Spirulina</i> mat condition. Marker was <i>Spirulina platensis</i> . The arrow shows the water exchanging and the water samples were taken before water exchanging <span style="float: right;">34</span>
4.4	16S rDNA-based denaturing gradient gel electrophoresis (DGGE) profiles (Left) with cluster analysis or percentage of Nei and Li's similarity coefficients (Right) of the bacterial community in water column of Set 2. Marker was pure culture of <i>Spirulina platensis</i> <span style="float: right;">35</span>
4.5	Microbial community in water column of Set 3 (co-cultured between artificial mat and shrimp). WA1-WA10 was microbial community in water column from day 1 to day 10 that reared shrimp from PL2 to PL12 in co-culturing with artificial mat condition. The arrow shows the water exchanging and the water samples were taken before water exchanging <span style="float: right;">37</span>

FIGURE	PAGE	
4.6	16S rDNA-based denaturing gradient gel electrophoresis (DGGE) profiles (Left) with cluster analysis or percentage of Nei and Li's similarity coefficients (Right) of the bacterial community in water column of Set 3. Lane WHO and WA1 to WA10 was bacteria community in water column in Set 3	38
4.7	The comparison of microbial community on <i>Spirulina</i> mat (S) and artificial mat (A) on day 0, 2, 5, 8 and 10, respectively	40
4.8	Changes of TAN in water column of Set 1, (○); Set 2, (●); and Set 3, (◇); respectively during PL2 to PL12 cultivation. The Set 1, 2 and 3 represented the only shrimp cultivation, shrimp co-cultured with <i>Spirulina</i> mat and shrimp co-cultured with artificial mat, respectively. White arrows indicate a 50% water exchange rate in Set 1 and Set 3 while Black indicate a 50% in Set 2, respectively	45
4.9	Changes in nitrite in water column of Set 1, (○); Set 2, (●); and Set 3, (◇); respectively during PL2 to PL12 cultivation. The set 1, 2 and 3 represented the only shrimp cultivation, shrimp co-cultured with <i>Spirulina</i> mat and shrimp co-cultured with artificial mat, respectively. White arrows indicate a 50% water exchange rate in Set 1 and Set 3 while Black indicate a 50% in Set 2, respectively	46
4.10	Changes in nitrate in water column of Set 1, (○); Set 2, (●); and Set 3, (◇); respectively during PL2 to PL12 cultivation. The Set 1, 2 and 3 represented the only shrimp cultivation, shrimp co-cultured with <i>Spirulina</i> mat and shrimp co-cultured with artificial mat, respectively. White arrows indicate a 50% water exchange rate in Set 1 and Set 3 while Black indicate a 50% in Set 2, respectively	47
4.11	Survival rates of shrimp among Set 1, (○); Set 2, (●); and Set 3, (◇); respectively during shrimp rearing periods for 10 days	49
4.12	Schematic flow of the co-culturing in shrimp larvae (PL2-PL12) with <i>Spirulina</i> mat. Solid line indicates Nitrogen flow; dash line indicates the relationship between modules	50
4.13	Using <i>Spirulina</i> mats during shrimp growing period in grow-out pond. A to F was <i>Spirulina</i> mat on day 0, 7 14, 16, 17 and 21, respectively. Figure G represented microbial community on <i>Spirulina</i> mat after 21 days under microscope	52
4.14	DGGE profile of bacterial amplicons from shrimp grow-out water; W <sub>0</sub> , microbial community before shrimp released to grow-out pond. W <sub>1</sub> , W <sub>2</sub> , and W <sub>3</sub> , after shrimp growing period 1,2 and 3 months, respectively. Ladder was the 22 clones mixture	53
4.15	Phylogenetic tree constructed based on 16S rDNA 1.5-kb clone sequences recovered from water during shrimp growing period in <i>Beta-proteobacteria</i> . The scale bar length of 0.1 denotes the number of amino acid replacements per site validated with 1000 bootstraps. The number on the branches indicates the support proportion of each branch	55

<b>FIGURE</b>	<b>PAGE</b>
4.16 Phylogenetic tree constructed based on 16S rDNA 1.5-kb clone sequences recovered from water during shrimp growing period in <i>Alpha</i> -, <i>Gamma</i> -proteobacteria and <i>Cytophaga-Flavobacterium Bacteroides</i> (CFB). The scale bar length of 0.1 denotes the number of amino acid replacements per site validated with 1000 bootstraps. The number on the branches indicates the support proportion of each branch	56
4.17 Phylogenetic tree constructed based on 16S rDNA 1.5-kb clone sequences recovered from water during shrimp growing period in Cyanobacteria and Chloroplast. The scale bar length of 0.1 denotes the number of amino acid replacements per site validated with 1000 bootstraps. The number on the branches indicates the support proportion of each branch	57
4.18 Ammonia oxidation rate: ammonia oxidation process with aerobic condition. $W_0$ , the activity in water before shrimp growing period; $W_1$ , $W_2$ , and $W_3$ , the activity in water after shrimp growing period 1, 2 and 3 months, respectively. The initial ammonia concentration was 10 mg-N L <sup>-1</sup>	65
4.19 Nitrite oxidation rate: nitrite oxidation process with aerobic condition. $W_0$ , the activity in water before shrimp growing period; $W_1$ , $W_2$ , and $W_3$ , the activity in water after shrimp growing period 1, 2 and 3 months, respectively. The initial nitrite concentration was 5 mg-N L <sup>-1</sup>	66
4.20 DGGE profile of bacterial diversity from the hatchery and shrimp grow-out pond water column. $WH_0$ , microbial community in hatchery water column and $WG_0$ , microbial community in grow-out pond water column	72