

## LIST OF ILLUSTRATIONS

Figure	Page
1.1 Structure of azo dye: PAR, Br-PADAP and Br-PAPS	5
3.1 Absorption spectra of PAPS-fluoride, Zr-PAPS-fluoride ( $0.5 \text{ mg L}^{-1}$ ) and Hf-PAPS-fluoride ( $2 \text{ mg L}^{-1}$ ) at pH 3.7. Conditions [PAPS] = $0.1 \text{ mM}$ , [fluoride] = $0.02 \text{ mM}$ .	9
3.2 Effect of PAPS concentration on peak height of Zr(IV) ( $0.1 \text{ mg L}^{-1}$ )	10
3.3 Effect of fluoride concentration on peak height of Zr(IV) ( $0.1 \text{ mg L}^{-1}$ )	10
3.4 Chromatogram of (a) Hf(IV)-PAPS-fluoride ( $0.4 \text{ mg L}^{-1}$ ), (b) Zr(IV)-PAPS-fluoride ( $0.1 \text{ mg L}^{-1}$ ) and (c) standard mixture of Zr(IV) ( $0.05 \text{ mg L}^{-1}$ ) and Hf(IV) ( $0.2 \text{ mg L}^{-1}$ ). Mobile phase was methanol-water 65% v/v containing $0.05 \text{ mM}$ fluoride and 10 mM acetate buffer at pH 5, analytical column 75 mm x 4 mm I.D., 3 $\mu\text{m}$ , Purospher C <sub>18</sub> ; flow rate 1 ml/min; temperature 30 °C; detection at 585 nm.	11
3.5 Effect of TBABr concentration on $k'$ of Zr-PAPS-fluoride and Hf-PAPS-fluoride. Other conditions are the same as in Figure 3.4	12
3.6 Chromatogram of (a) blank, (b) Hf(IV)-PAPS-fluoride ( $20 \text{ }\mu\text{g L}^{-1}$ ), (c) Zr(IV)-PAPS-fluoride ( $5 \text{ }\mu\text{g L}^{-1}$ ) and (d) standard mixture of Zr(IV) ( $10 \text{ }\mu\text{g L}^{-1}$ ) and Hf(IV) ( $40 \text{ }\mu\text{g L}^{-1}$ ). Mobile phase was methanol-water 65% v/v containing $0.05 \text{ mM}$ fluoride, 5 mM TBABr and 10 mM acetate buffer at pH 5. Other conditions are the same as in Figure 3.4	13
3.7 Effect of pH on peak height of Zr(IV)-PAPS-fluoride ( $50 \text{ }\mu\text{g L}^{-1}$ ) and Hf(IV)-PAPS-fluoride ( $200 \text{ }\mu\text{g L}^{-1}$ ). Other conditions are the same as in Figure 3.4	13
3.8 Chromatogram of (a) blank, (b) granite rock sample KP112 and (c) granite rock sample KP112 spiked with $5 \text{ }\mu\text{g L}^{-1}$ Zr(IV) and $20 \text{ }\mu\text{g L}^{-1}$ Hf(IV). Top: complete view. Bottom: zoom view. The sample contained [PAPS] = $0.25 \text{ mM}$ , [fluoride] = $0.02 \text{ mM}$ . Other conditions are the same as Figure 3.6	15

7.1	Structure of imidazole	33
7.2	Spectrum of background electrolyte containing 12 mM imidazole, 3 mM 18-crown-6 ether and 15 mM alanine at pH 6	33
7.3	Electropherogram of standard (a) $\text{NH}_4^+$ (2.5 mg L <sup>-1</sup> ), (b) $\text{K}^+$ (5 mg L <sup>-1</sup> ), (c) $\text{Ca}^{2+}$ (1.25 mg L <sup>-1</sup> ), (d) $\text{Mg}^{2+}$ (0.5 mg L <sup>-1</sup> ) and (e) mixture of $\text{NH}_4^+$ (2.5 mg L <sup>-1</sup> ), $\text{K}^+$ (5 mg L <sup>-1</sup> ), $\text{Ca}^{2+}$ (1.25 mg L <sup>-1</sup> ), and $\text{Mg}^{2+}$ (0.5 mg L <sup>-1</sup> ). Electrolyte: 12 mM imidazole, 3 mM 18-crown-6 ether pH 4.3. Separation voltage: +25 kV.	34
7.4	Electrolyte: 12 mM imidazole, 3 mM 18-crown-6 ether pH 6.0. Concentration of standard (a), (b), (c) and (d) are the same as in Figure 7.3	35
7.5	Electrolyte: 12 mM imidazole, 3 mM 18-crown-6 ether and 15 mM alanine at pH 6.0. Concentration of standard (a), (b), (c) and (d) are the same as in Figure 7.3	35
7.6	Electropherogram of $\text{NH}_4^+$ (2.5 mg L <sup>-1</sup> ), $\text{K}^+$ (5 mg L <sup>-1</sup> ), $\text{Ca}^{2+}$ (1.25 mg L <sup>-1</sup> ), $\text{Na}^+$ (1 mg L <sup>-1</sup> ), $\text{Mg}^{2+}$ (0.5 mg L <sup>-1</sup> ), $\text{Mn}^{2+}$ (1 mg L <sup>-1</sup> ), $\text{Co}^{2+}$ (1 mg L <sup>-1</sup> ) and $\text{Cd}^{2+}$ (5 mg L <sup>-1</sup> ) in a standard solution; (a) electrolyte: 12 mM imidazole and 15 mM alanine at pH 6; (b) electrolyte: 12 mM imidazole, 3 mM 18-crown-6 ether and 15 mM alanine at pH 6.0	36
7.7	Co-EOF of standard $\text{NH}_4^+$ (2.5 mg L <sup>-1</sup> ), $\text{K}^+$ (5 mg L <sup>-1</sup> ), $\text{Ca}^{2+}$ (1.25 mg L <sup>-1</sup> ), $\text{Na}^+$ (1 mg L <sup>-1</sup> ), $\text{Mg}^{2+}$ (0.5 mg L <sup>-1</sup> ), $\text{Mn}^{2+}$ (1 mg L <sup>-1</sup> ), $\text{Co}^{2+}$ (1 mg L <sup>-1</sup> ) and $\text{Cd}^{2+}$ (5 mg L <sup>-1</sup> ). The electrolyte was 12 mM imidazole, 3 mM 18-crown-6 ether and 15 mM alanine at pH 6.0; the separation temperature 25°C, the separation voltage +25 kV and injection time 7 s under pressure 50 mbar	37
7.8	Electropherogram of solid fertilizer sample: N-P-K = 18-4-6; (a) blank; (b) unspiked N-P-K fertilizer; (c) N-P-K fertilizer spiked with 2 mg L <sup>-1</sup> $\text{NH}_4^+$ , 2.5 mg L <sup>-1</sup> $\text{K}^+$ , 0.5 mg L <sup>-1</sup> $\text{Ca}^{2+}$ , 0.3 mg L <sup>-1</sup> $\text{Na}^+$ and 0.2 mg L <sup>-1</sup> $\text{Mg}^{2+}$ . Other conditions are the same as in Figure 7.7	41
7.9	Electropherogram of Ca-Mg fluid fertilizer sample; (a) unspiked Ca-Mg fertilizer; (b) Ca-Mg fertilizer spiked with 0.5 mg L <sup>-1</sup> $\text{Ca}^{2+}$ and 0.2 mg L <sup>-1</sup> $\text{Mg}^{2+}$ . Other conditions are the same as in Figure 7.7	41