

ปัจจัยที่มีความสัมพันธ์ต่อการปฏิบัติงานตามแนวคิดการดูแลระดับปฐมภูมิสำหรับโรคหลอดเลือดของหัวใจ ของ
 พยาบาลสาธารณสุข ศูนย์บริการสาธารณสุข กรุงเทพมหานคร
 CORRELATES OF PRACTICE ON PRIMARY CARE CONCEPT FOR CORONARY HEART DISEASES OF
 PUBLIC HEALTH NURSES IN PUBLIC HEALTH CENTERS, BANGKOK METROPOLITAN
 ADMINISTRATION

สุคฤทัย เรืองฤทธิ์ 5037011 PHPH/M

วท.ม. (สาธารณสุขศาสตร์) สาขาวิชาเอกการพยาบาลสาธารณสุข

คณะกรรมการที่ปรึกษาวิทยานิพนธ์ : ปัญญรัตน์ ลาภวงศ์วัฒนา, Ph.D., สุนิย์ ละกำป็น กศ.ด.

บทคัดย่อ

การวิจัยพรรณนาเชิงวิเคราะห์นี้มีวัตถุประสงค์เพื่อทดสอบหาปัจจัยที่มีความสัมพันธ์ต่อการปฏิบัติงานตามแนวคิดการดูแลระดับปฐมภูมิสำหรับโรคหลอดเลือดของหัวใจ ของพยาบาลสาธารณสุข ศูนย์บริการสาธารณสุข กรุงเทพมหานคร กลุ่มตัวอย่างได้จากการสุ่มอย่างง่ายเป็นพยาบาลสาธารณสุขปฏิบัติหน้าที่เยี่ยมบ้านของศูนย์บริการสาธารณสุข 68 แห่ง จำนวน 136 คน จากทั้งหมด 330 คน เก็บรวบรวมข้อมูล ด้วยการให้ตอบแบบสอบถามชนิดตอบเอง เครื่องมือที่ใช้ในการเก็บข้อมูลเป็นแบบวัดมี 5 ส่วน คือ ปัจจัยนำ ประกอบด้วย คุณลักษณะส่วนบุคคล ความรู้และเจตคติ ปัจจัยเอื้อ ได้แก่ นโยบายหน่วยงาน การเข้าถึงข้อมูลสุขภาพ และการได้รับการฝึกอบรม ปัจจัยเสริม ได้แก่ การสนับสนุนจากผู้บังคับบัญชา การสนับสนุนผู้ร่วมงาน และการสนับสนุนอาสาสมัครสาธารณสุข และการปฏิบัติงานตามแนวคิดการดูแลระดับปฐมภูมิสำหรับโรคหลอดเลือดของหัวใจ วิเคราะห์ข้อมูลโดยการแจกแจงความถี่ ร้อยละ ค่าเฉลี่ย ส่วนเบี่ยงเบนมาตรฐาน สัมประสิทธิ์สหสัมพันธ์ของเพียร์สัน และการวิเคราะห์ถดถอยเชิงพหุ

ผลการวิจัยพบว่า พยาบาลสาธารณสุขปฏิบัติงานตามแนวคิดการดูแลระดับปฐมภูมิสำหรับโรคหลอดเลือดของหัวใจ อยู่ในระดับมาก ($\bar{X} = 77.10$, S.D.=9.91) โดยเฉพาะการบริการแบบบูรณาการ การบริการเชิงรุก การบริการที่ส่งเสริมสนับสนุนให้เกิดการดูแลสุขภาพตนเอง และการบริการด้านแรก อยู่ในระดับมาก ($\bar{X} = 13.75, 13.44, 12.90, 12.29$ ตามลำดับ) ปัจจัยที่สามารถทำนายการปฏิบัติงานตามแนวคิดการดูแลระดับปฐมภูมิสำหรับโรคหลอดเลือดของหัวใจ คือ ปัจจัยนำ ได้แก่ ประสิทธิภาพการทำงาน ($\beta = .236, p < .002$) เจตคติต่อการปฏิบัติงานตามแนวคิดการดูแลระดับปฐมภูมิสำหรับโรคหลอดเลือดของหัวใจ ($\beta = .201, p < .011$) ปัจจัยเอื้อ ได้แก่ นโยบายหน่วยงาน ($\beta = .356, p < .004$) การเข้าถึงข้อมูลสุขภาพ ($\beta = -.290, p < .010$) ปัจจัยเสริม ได้แก่ การได้รับการสนับสนุนจากผู้บังคับบัญชา ($\beta = .220, p < .049$) โดยสามารถร่วมกันทำนายได้ร้อยละ 28.8

มีข้อเสนอแนะว่า การพัฒนาความสามารถของพยาบาลสาธารณสุขในการปฏิบัติงานตามแนวคิดการดูแลระดับปฐมภูมิสำหรับโรคหลอดเลือดของหัวใจนั้น ต้องมีการกำหนดนโยบาย โดย ผู้บังคับบัญชา มีหน้าที่สนับสนุน ความรู้และพัฒนาศักยภาพด้านการรักษาโรคเบื้องต้น และการฟื้นฟูสุขภาพผู้ป่วยโรคหลอดเลือดของหัวใจที่บ้านอย่างต่อเนื่อง ให้เข้าถึงแหล่งข้อมูลสุขภาพ ได้รับการส่งเสริมเจตคติที่ดีต่อการปฏิบัติงาน และผู้ปฏิบัติงานต้องมีประสิทธิภาพดูแลผู้ป่วยโรคหลอดเลือดของหัวใจในชุมชน เพื่อปฏิบัติงานดูแลระดับปฐมภูมิได้อย่างมีประสิทธิภาพ

คำสำคัญ : แนวคิดการดูแลระดับปฐมภูมิ/ โรคหลอดเลือดของหัวใจ/ พยาบาลสาธารณสุข

CORRELATES OF PRACTICE ON PRIMARY CARE CONCEPT FOR
CORONARY HEART DISEASES OF PUBLIC HEALTH NURSES IN PUBLIC
HEALTH CENTERS, BANGKOK METROPOLITAN ADMINISTRATION

SUDRUETHAI RUEANGRIT 5037011 PHPH/M

MS.c. (PUBLIC HEALTH) MAJOR IN PUBLIC HEALTH NURSING

THESIS ADVISORY COMMITTEE: PUNYARAT LAPVONGWATANA, Ph.D.,
SUNEE LAGAMPAN, Ed.D.

ABSTRACT

This descriptive analytic research aims to examine correlates of primary care concept practice by public health nurses regarding coronary heart diseases. The 136 samples from 330 public health nurses were randomly selected from 68 Public Health Centers, Bangkok Metropolitan Administration. Self administered-questionnaires consisted of 5 parts. The predisposing factors comprised forms on general characteristics knowledge and attitudes. Enabling factors: departmental policies, information accessibility and training, and reinforcing factors: support from head, coworkers and health volunteers, as well as primary care practice for coronary heart disease scales were also investigated. Data were analyzed by using frequency, mean, standard deviation, Pearson's product moment correlation coefficient, and multiple regressions.

Results indicated that overall primary care concept practices of public health nurses for coronary heart diseases were at a high level ($\bar{x} = 77.10$, $SD = 9.91$): specifically, first contact care, preventive care, integrated care, and self care obtained means of 13.75, 13.44, 12.90, and 12.29, respectively. Significant variables that predicted primary care practice for coronary heart diseases were predisposing factors including work experience ($\beta = .236$, $p < .002$) and attitude to coronary heart diseases ($\beta = .201$, $p < .011$); the enabling factors of departmental policies ($\beta = .356$, $p < .004$), and accessible health information ($\beta = -.290$, $p < .010$); and the reinforcing factor of support for the administrators ($\beta = .220$, $p < .049$), which contributed to 28.8 percent of variance.

Recommendations are made to the head nurses to establish a transparent and concise policy for practicing primary care among coronary heart disease patients so that all health nurses could improve their own competencies. In addition, the practitioners should have primary care experience and support knowledge development. They should support continued basic treatment and rehabilitation at home. Moreover, they must enhance the accessibility of health information resources and promote a good attitude to primary care concept practice.

KEY WORDS: PRIMARY CARE CONCEPT / CORONARY HEART DISEASES/
PUBLIC HEALTH NURSES

172 pages