

REFERENCES

- Abollino, O., Aceto, M., Malandrino, M., Sarzanini, C., and Mentasti, E. 2003. Adsorption of heavy metals on Na-montmorillonite. Effect of pH and organic substances. Water Research 37 (7): 1619-1627.
- Abut, M., Skye, R., Rimbaud, A., Potion-Gautier, M., Teller, S., and Bannerman, W. 2003. Improving the environmental management of small-scale gold mining in Ghana: a case study of Dumas. Journal of Cleaner Production 11: 215-221.
- Adhikari, T. and Singh, M.V. 2003. Sorption characteristics of lead and cadmium in some soils of India. Geoderma 114: 81-92.
- Allen, T. 1974. Particle Size Measurement. Chapman & Hall, London.
- Alumaa, P., Kirso, U., Petersell, V., and Steinnes, E. 2002. Sorption of toxic heavy metals to soils. International Journal of Hygiene and Environmental Health 204: 375-376.
- Antoniadis, V., McKinley, J.D., and Zuhairi, W.Y.W. 2007. Single-element and competitive metal mobility measured with column infiltration and batch tests. Journal of Environmental Quality 36: 53-60.
- Appel, C. and Ma, L. 2002. Concentration, pH, and Surface Charge Effects on Cadmium and Lead Sorption in Three Tropical Soils. Journal of Environmental Quality 31: 581-589.
- Arias, M., Perez-Novo, C., Lopez, E., and Soto, B. 2006. Competitive adsorption of copper and zinc in acid soils. Geoderma 133: 151-159.
- Aswathararyana, U. 2003. Mineral Resources Management and the Environment. A.A. Balkema, Tokyo. 294 pp.
- ATSDR (Agency for Toxic Substances and Disease Registry). 2000a. Toxicological Profile for Nickel. U.S.Department of Public Health and Human Services, Public Health Service, Atlanta, GA.
- ATSDR (Agency for Toxic Substances and Disease Registry). 2000b. Toxicological Profile for Zinc (Update). U.S.Department of Public Health and Human Services, Public Health Service, Atlanta, GA.
- ATSDR (Agency for Toxic Substances and Disease Registry). 2000c. Toxicological Profile for Manganese. U.S.Department of Public Health and Human Services, Public Health Service, Atlanta, GA.

- ATSDR (Agency for Toxic Substances and Disease Registry). 2005. Toxicological Profile for Lead (Draft for Public Comment). U.S. Department of Public Health and Human Services, Public Health Service, Atlanta, GA.
- Babut, M., Sekyi, R., Rambaud, A., Potin-Gautier, M., Tellier, S., and Bannerman, W. 2003. Improving the environmental management of small-scale gold mining in Ghana: a case study of Dumasi. Journal of Cleaner Production 11:215-221.
- Bajracharya, K., Tran, Y.T., and Barry, D.A. 1996. Cadmium adsorption at different pore water velocities. Geoderma 73: 197-216.
- Basta, N.T. and Tabatabai, M.A. 1992. Effect of cropping system on adsorption of metals by soils. Soil Science 153: 331-337.
- Bedient, P.B., Rifai, H.S., and Newell, C.J. 1999. Ground Water Contamination. Prentice Hall.
- Beigel, C. and Pietro, L.D. 1999. Transport of Trifluoromethylazole in homogeneous soil columns: Influence of nonequilibrium sorption. Soil Science Society of American Journal 63: 1077-1086.
- Bellir, K., Bencheikh-Lehocine, M., Meniai, A-H., and Gherbi, N. 2005. Study of the retention of heavy metals by natural material used as liners in landfills. Desalination 185: 111-119.
- Bin-Shafique, M.S., Benson, C.H., and Edil T.B. 2002. Leaching of Heavy Metals from Fly Ash Stabilized Soils used in Highway Pavements. Geo Engineering Report No.02-14: 1-204.
- Blowes, D.W. 1997. The environmental effects of mine wastes. Mapping and monitoring the mine environment 119: 887-892.
- Brady, K.B.C. and Hornberger, R.H. 1990. The prediction of mine drainage quality in Pennsylvania. Water Pollution Control Association Pa Magazine 23: 8-15.
- Brunauer, S., Emmett, P.H., and Teller, E. 1938. Adsorption of gases in multimolecular layers. Journal of the American Chemical Society 60: 309-319.
- Brusseau, M.L., Larsen, T., and Christensen, T.H. 1991. Rate-limited sorption and non-equilibrium transport of organic chemicals in low organic carbon aquifer materials. Water Resources Research 28 (9): 2485-2497.
- Burckhard, S.R., Schwab, A.P., and Banks, M.K. 1995. The effects of organic acids on the leaching of heavy metals from mine tailings. Journal of Hazardous Materials 41: 135-145.

- Bürgisser, C.S., Černik, M., Borkovec, M., and Sticher, H. 1993. Determination of nonlinear adsorption isotherms from column experiments: An alternative to batch studies. Environmental Science and Technology 27: 943-948.
- Cavallaro, N. and McBride, M.B. 1980. Activities of Cu^{2+} and Cd^{2+} in soil solutions as affected by pH. Soil Science Society of America Journal 44: 729-732.
- Celardin, F., Trouillet, C., and Tisiot, R. 2004. pH dependence of copper adsorption in vineyard soils of Geneva. Environmental Chemistry Letters 1:225-227.
- Chalermyanont, T., Arrykul, S., and Charoenthaisong, N. 2008. Potential use of lateritic and marine clay soils as landfill liners to retain heavy metals. Waste Management (Article in Press).
- Chang, C.M., Wang, M.K., Chang, T.W., Lin, C., and Chen, Y.R. 2001. Transport modeling of copper and cadmium with linear and nonlinear retardation factors. Chemosphere 43 : 1133-1139.
- Cidu, R., Biagini, C., Fanfani, L., La Ruffa, G., and Marras, I. 2000. Mine closure at Monteponi (Italy): Effect of the cessation of dewatering on the quality of shallow groundwater. Applied Geochemistry 16: 489-502.
- Concas, A., Arda, C., Cristini, A., Zuddas, P., and Cao, G. 2005. Mobility of heavy metals from tailings to stream waters in a mining activity contaminated site. Chemosphere (Article in Press).
- Connaughton, D.F., Stedinger, J.G., Lion, L.W., and Shuler, M.L. 1993. Description of time-varying desorption kinetics: release of naphthalene from contaminated soils. Environmental Science & Technology 27: 2397-2403.
- Darland, J.E. 1997. Effect of pore water velocity on the transport of arsenate. Environmental Science and Technology 31: 704-709.
- Di Torro, D.M., Mahony, J.D., Kirchgraber, P.R., O'Byrne, A.L., Pasquale, L.R., and Piccirilli, D.C. 1986. Effect of nonreversibility, particle concentration and ionic strength on heavy metal sorption. Environmental Science and Technology 20 (1): 55-61.
- Drillia, P., Stamateletou, K., and Lyberatos, G. 2005. Fate and mobility of pharmaceuticals in solid matrices. Chemosphere 60: 1034-1044.
- Dubrovsky, N.M. 1986. Geochemical Evolution of Inactive Pyritic Tailings in the Elliot Lake Uranium District. Doctoral dissertation. University of Waterloo Canada.
- Elzahabi, M. and Yong, R.N. 2001. pH influence on sorption characteristics of heavy

- metal in the vadose zone. Engineering Geology 60: 61-68.
- EPA (US Environmental Protection Agency). 1993. Acid generation prediction in mining: Draft. Crystal City, VA.
- EPA (US Environmental Protection Agency). 1994. Method 3051, Micro-wave assisted acid digestion of sediments, sludges, soils and oils. [online]. Available from: URL: <http://www.epa.gov/epaoswer/hazwaste/test3051.pdf>. [May 20, 2007].
- Erses, A.S., Fazal, M.A., Onay, T.T., and Craing, W.H. 2005. Determination of solid waste sorption capacity for selected heavy metals in landfills. Journal of Hazardous Materials B 121: 223-232.
- Esposito, A., Pagnanelli, F., and Veglio, F. 2002. pH-related equilibria models for biosorption in single metal systems. Chemical Engineering Science 57: 307-313.
- Fetter, C.W. 1993. Contaminant Hydrogeology. New Jersey: Practice Hall.
- Freeze, R.A. and Cherry, J.A. 1979. Groundwater. Prentice Hall.
- Gaber, H.M., Inskeep, W.P., Comfort, S.D., and Wraith, J.M. 1995. Nonequilibrium transport of atrazine through large intact soil cores. Soil Science Society of America Journal 59: 60-67.
- Gao, Y. and Bradshaw, A.D. 1995. The containment of toxic wastes: II. Metal movement in leachate and drainage at Parc lead-zinc mine, North Wales. Environmental Pollution 90: 379-382.
- Gomes, P.C., Fontes, M.P.F., Silva, A.G.D, Mendonça, E.D.S., and Netto, A.R. 2001. Selectivity sequence and competitive adsorption of heavy metals by Brazilian soils. Soil Science Society of America Journal 65: 1115-1121.
- Gunsunger, M.R., Ptacek, C.J., Blowes, D.W., and Jambor, J.L. 2006. Evaluation of long-term sulfide oxidation processes within pyrrhotite-rich tailings, Lynn Lake, Manitoba. Journal of Contaminant Hydrology 83: 149-170.
- Gutierrez, M. and Fuentes, H.R., 1991. Competitive Adsorption of Cesium, Cobalt and Strontium in Conditioned Clayey Soil Suspensions. Journal of Environmental Radioactivity 13: 271-282.
- Harter, R.D. 1992. Competitive sorption of Cobalt, Copper, and Nickel ions by a calcium-saturated soil. Soil Science Society of America Journal 56: 444-449.
- Harter, R.D. and Naidu, R. 2001. An assessment of environmental and solution parameters impact on trace-metal sorption by soils. Soil Science Society of

- American Journal 65: 597-612.
- Hinz, C. and Selim, H. 1994. Lead adsorption onto aquifer gravel using batch experiments and XPS. In: Arehart, G.B., Hulston, J.R. (Eds.), Water-Rock Interaction. pp. 939-942. Balkema, Rotterdam.
- Hooda, P.S. and Alloway, B.J. 1998. Cadmium and lead sorption behaviour of selected English and Indian soils. Geoderma 84: 121-134.
- Jain, C.K., and Ram, D.A. 1996. Adsorption of lead and zinc on bed sediments of the river Kali. Water Research 31: 154-162.
- Jeon, B-H., Dempsey, B.A., Burgos, W.D., Roger, R.A., and Roden, E.E. 2004. Modeling the sorption kinetics of divalent metal ions to hematite. Water Research. 38: 2499-2504.
- Jurjovec, J., Ptacek, C., and Blowes, D.W. 2001. Acid neutralization mechanisms and metals release in mine tailings: A laboratory column experiment. Geochimica Et Cosmochimica Acta 66: 1511-1523.
- Kabala, C. and Singh, B.R. 2001. Fractionation and mobility of copper, lead, and zinc in soil profiles in the vicinity of a copper smelter. Journal of Environmental Quality 30: 485-492.
- Kalbitz, K., and Wennrich, R. 1998. Mobilization of heavy metals and arsenic in polluted wetland soils and its dependence on dissolved organic matter. The Science of Total Environment 209: 27-39.
- Kookana, R.S., Naidu, R., and Tiller, K.G. 1994. Sorption non-equilibrium during cadmium transport through soils. Australian Journal of Soil Research 32: 635-651.
- Kookana, R.S. and Naida, R. 1998. Effect of soil solution composition on cadmium transport through variable charge soils. Geoderma 84: 235-248.
- Kwon, J., Yun, S., Kim, S., Mayer, B., and Huntcheon, I. 2005. Sorption of Zn (II) in aqueous solutions by scoria. Chemosphere 60: 1416-1426.
- Lee, S., Chang, L., Yang, H., Chen, C., and Liu, M., Adsorption characteristics of lead onto soils. Journal of Hazardous Materials 63: 37-49.
- Leitao, T.E., Smets, S., Beek, C.V., and Ferreira, J.P.L. 2000. Migration of Heavy Metals in Saturated Porous Media: Simulation in laboratory experiment. 5th Congresso da Agua, 1-15.
- Leitao, T.E. and Zakharova, T. 2003. Competitive adsorption of heavy metals: Simulation in laboratory experiments. Seminario Sobre Aguas Subterraneas

- (APRH/CEAS), pp.1-12. Feb 27-28, 2003. Lisbon.
- Li, Q. 2000. Effect of surfactants on the sorption and transport of Cu (II) in a sandy soil: batch and column experiments and modeling. Master's Thesis. Department of Building, Civil and Environmental Engineering. Concordia University.
- Liu, C.L., Chang, T., Wang, M., and Huang, C. 2006. Transport of cadmium, nickel, and zinc in Taoyuan red soil using one-dimensional convective-dispersive model. Geoderma 131: 181-189.
- Lottermoser, B. 2003. Mine Waste: Characterization, Treatment and Environmental Impacts. pp. 277. Berlin: Springer.
- Macro, D.B., Claudio, G., Veronica, G., Cataldo, L.G., Bruno, P., Carmela, R., and Sergio, S. 2002. Chemical and geological characterization of a mine site in Central Italy. 17th International Conference on World Congress of Soil Science, pp. 1-11. August 14-21, 2002. Bangkok.
- Maraq, M.A. 2001. Effects of fundamental differences between batch and miscible displacement techniques on sorption distribution coefficient. Environmental geology 41: 219-228.
- Martínez-Villegas, N., Flores-Vélez, L.M., and Domínguez, O. 2004. Sorption of lead in soil as a function of pH: a study case in México. Chemosphere 57: 1537-1542.
- McKenzie, R.M. 1980. The adsorption of lead and other heavy metals on oxide of manganese and iron. Australian Journal of Soil Research 18: 61-73.
- McLean, J.E. and Bledsoe, B.E. 1992. EPA Ground Water Issue: Behavior of metals in soils. EPA/540/S-92/018, Office of Solid Waste and Emergency Response, Washington, DC.
- Mesquita, M.E., and Vieira e Silva, J.M. Preliminary study of Ph effect in the application of Langmuir and Freundlich isotherms to Cu-Zn competitive adsorption. Geoderma 106: 219-234.
- Ministry of Industry. 1996. Notification of Ministry of Industry No.2 B.E. 2539 (1996) [online]. Available from. URL: <http://www2.DIW.go.th/PIC/download/water/water2.pdf>. [April 17, 2008]
- Miretzky, P., Muñoz, C., and Carrillo-Chávez, A. 2006. Experimental Zn (II) retention in a sandy loam soil by very small columns. Chemosphere 65: 2082-2089.

- Mohan, D. and Chander, S. 2006. Single, binary, and multicomponent of iron and manganese on lignite. Journal of Colloid and Interface Science 299: 76-87.
- Mohan, D. and Singh, K.P. 2002. Single- and multi-component adsorption of cadmium and zinc using activated carbon derived from bagasse- an agriculture waste. Water Research 36: 2304-2318.
- Moncur, M.C., Ptacek, C.J., Blowes, D.W., and Jambor, J.L. 2005 .Release, transport and attenuation of metals from an old tailings impoundment. Applied Geochemistry 20: 639-659.
- Moncur, M.C., Ptacek, C.J., Blowes, D.W., and Jambor, J.L. Spatial variations in water composition at a northern Canadian lake impacted by mine drainage. Applied Geochemistry 21: 1799-1817.
- Moreno, L. and Neretnieks, I. 2006. Long-term environmental impact of tailings deposits. Hydrometallurgy 83: 176-183.
- Muhammad, N., Parr, J., Smith, M.D., and Wheatley, A.D. 1998. Adsorption of heavy metals in slow sand filters. The WEDC Conference on sanitation and water for all. 24th, 1998. pp. 346-349. Islamabad, Pakistan.
- Nedunuri, K.V., Erickson L.E., and Govindaraju, R.S. 1998. Modeling the role of active biomass on fate and transport of a heavy metal in the presence of root exudates. Journal of Hazardous Substance Research 1: 9-24.
- Neves, O. and Matias, M.J. 2008. Assessment of groundwater quality and contamination problems ascribed to an abandoned uranium mine (Cunha Baixa region, Central Portugal). Environmental Geology 53: 1799-1810.
- Nkedi-Kizza, P., Biggar, J.W., van Genuchten, M.Th., Wierenga, P.J., Selim, H.M., Davidson, J.M., and Nielson, D.R. 1983. Modeling tritium and chloride-36 transport an aggregated Oxisol. Water Resources Research 19: 691-700.
- Nkedi-Kizza, P, Rao, P.S.C., Hornsby, A.G. 1987. Influence of organic cosolvent on leaching of hydrophobic organic chemicals through soils. Environmental Science and Technology 21: 1107-1111.
- Oren, A.H. and K, A. 2005. Factors affecting adsorption characteristics of Zn^{2+} on two natural zeolites. Journal of Hazardous Materials B 131: 59-65.
- Ouyang, Y., Shinde, D., Mansell, R.S., and Harris, W. 1996. Colloid-enhanced transport of chemicals in subsurface environments: A review. Critical Review: Environmental Science and Technology 26: 189-204.
- Panteleit, B., Kessels, W., and Schulz, H.D. 2002. Geochemical processes in the salt-

- freshwater transtion zone – Preliminary results of a 2D sand tank experiment. 17th Salt Water Intrusion Meeting, Delft pp. 29-38. May 6-10, 2002. The Netherlands.
- Pang, L. 1995. Contamination of groundwater in the Te Arota area by heavy metals from an abandoned mine. Journal of Hydrology 33: 17-34.
- Pang, L., Close, M., Schneider, D., and Stanton, G. 2002. Effect of pore-water velocity on chemical nonequilibrium transport of Cd, Zn, and Pb in alluvial gravel columns. Journal of Contaminant Hydrology 57: 241-258.
- Pang, L., Close, M., Greenfield, H., and Stanton, G. 2004. Adsorption and transport of cadmium and rhodamine WT in pumice sand columns. New Zealand Journal of Marine and Freshwater Research 38: 367-378.
- Passariello, B., Giuliano, V., Quaresima, S., Barbaro, M., Caroli, S., Forte, G., Carelli, G., and Iavicoli, I. 2002. Evaluation of the environmental contamination at an abandoned mining site. Microchemical Journal 73: 245-250.
- Pazka, T. 2003. Studies of competitive Cu^{2+} , Co^{2+} , and Cr^{3+} sorption in Grey-Brown Podzolic soils. Polish Journal of Environmental Studies 4: 439-446.
- PCD (Pollution Control Department). 2004a. Agricultural soil standard. [online]. Available from: URL: http://www.pcd.go.th/info_serv/reg_std_soil01.html#s1. [May 20, 2007].
- PCD (Pollution Control Department). 2004b. Groundwater standard. [online]. Available from: URL: http://www.pcd.go.th/info_serv/reg_std_water03.html. [May 20, 2007].
- Pinto, M.J., Gritman, H., and Gorelick, S.M. 1997. Laboratory-scale analysis of aquifer remediation by in-well vapor stripping: 2. Modeling results. Journal of Contaminant Hydrology 29: 41-58.
- Plassard, F., Winiarski, T., and Ramel, M.P. 2000. Retention and distribution of three heavy metals in a carbonated soil: Comparison between batch and unsaturated column studies. Journal of Contaminant Hydrology 42: 99-111.
- Ponizovsky, A.A. and Tsadilas, C.D. 2003. Lead (II) retention by Alfisol and clinoptilolite: cation balance and pH effect. Geoderma 115: 303-312.
- Protano, G. and Riccobono, F. 1997a. Antimony and arsenic contents in fresh waters and stream sediments of the Monti Romani area (Southern Tuscany, Italy). Mineralogical Petrografica Acta (Bologna) XL: 213-226.
- Putthividhya, A. and Chotpantararat, S. 2006. Variable water saturation impact on

- sorption/desorption characteristics of multiple species heavy metals from gold mining. The international symposium Darcy 2006. May 30- June 1, 2006. Dijon, France.
- Putthividhya, A. and Chotpantarat, S. 2007. Influence of Pb²⁺ on Competitive Binary Sorption of Zn²⁺, Ni²⁺, and Mn²⁺ onto Lateritic Soil. The IAH congress, groundwater & ecosystems. September 17-21, 2007. Lisbon.
- Pyatt, F.B. and Grant, tan P.J. 2001. Some consequences of ancient mining activities on the health of ancient and modern human populations. Journal of Public Health Medicine 23: 235-236.
- Ramirez-Requelme, M.E., Ramos, J.F.F., Angelica, R.S., and Brabo, E.S. 2002. Assessment of Hg-contamination in soils and stream sediments in the mineral district of Nambija, Ecuadorian Amazon (example of an impacted area affected by artisanal gold mining). Applied Geochemistry 18: 371-381.
- Reed, R.E., and Nonavinakere, S.K. 1992. Metal adsorption by activated carbon: effect of complexing ligands, competing adsorbates, ionic strength, and background electrolyte. Separation Science and Technology 27(14): 1985-2000.
- Reeuwijk, L.P. (ed) 1995. Procedures for soil analysis. Tech. Paper 9. 5th ed. ISRIC, Wageningen, the Netherlands.
- Rodríguez-maroto, J.M., García-delgado, R.A., Gómez-lahoz, C., Vereda-alonso, C., García-herruzo, F., and Muñoz, M.P.P. 2003. Competitive retention of lead and cadmium on an agricultural soil. Environmental Monitoring and Assessment 89: 165-177.
- Roszyk, E. and Szerszen, L. 1988. Accumulation of heavy metals in arable layer of soils in the protection zones of copper smelters. Rocz. Glebozn. 39: 135-156.
- Salomons, W. 1995. Environmental impact of metals derived from mining activities: Processes, predictions, prevention. Journal of Geochmcal Exploration 52: 5-23.
- Schwarzenbach, R.P. and Westall, J. 1981. Transport of nonpolar organic compounds from surface water to groundwater: laboratory sorption studies. Environmental Science and Technology 15: 1360-1367.
- Schwertmann, U., and Taylor, R.M. 1989. Iron oxides. p. 379-438. In J.B. Dixon and S.B. Weed (ed.) Minerals in soil environments. ASA and SSSA, Madison, WI.
- Selim, H.M., Davidson, J.M., and Mansell, R.S. 1976. Evaluation of a two-site

- adsorption-desorption model for describing solute transport in soil. Computer simulation conference, pp. 444-448. Washington, DC.
- Selim, H.M., and Amacer, M.C. 1996. Reactivity and Transport of Heavy Metals in Soils. Boca Raton: CRC Press.
- Selim, H.M. 1999. Modeling the kinetics of heavy metals reactivity in soils. In H.M. Selim and K.I. Iskandar (eds.), Fate and Transport of heavy metal in the vadose zone, pp. 91-106., Boca Raton, Florida: Lewis Publish.
- Selim, H.M., and Sparks, D.L. 2000. Heavy Metals Release in Soils. Boca Raton: CRC Press.
- Serrano, S., Garrido, F., Campbell, C.G., and Garcia-Gonzalez, M.T. 2005. Competitive sorption of cadmium and lead in acid soils of Central Spain. Geoderma 124: 91-104.
- Seuntjens, P., Tirez, K., Simunek, J., van Genuchten, M.Th., Cornelis, C., and Geuzens, P. 2001. Aging effects on cadmium transport in undisturbed contaminated sandy soil columns. Journal of Environmental Quality 30: 1040-1050.
- Sharma, H.D. and Reddy, K.R. 2004. Geoenvironmental engineering: Site remediation, water contaminant, and emerging waste management technologies. New Jersey: John Wiley & Sons Inc.
- Sheoran, A.S. and Sheoran, V. 2006. Heavy metal removal mechanism of acid mine drainage in wetlands: A critical review. Minerals Engineering 19:105-116.
- Simunek, J., Šejna, M., and van Genuchten, M.T. 2008. The HYDRUS-1D software package for simulating the one-dimensional movement of water, heat, and multiple solutes in variably saturated media. Version 4.Beta 1. ICGWMC-TPS-70, International Ground Water Modeling Center, Colorado School of Mines, Golden, Colorado.
- Sparks, D.L. 1995. Environmental Soil Chemistry. San Diego, California: Academic Press.
- Srivastava, P., Singh, B., and Angove, M. 2005. Competitive adsorption behavior of heavy metals on kaolinite. Journal of Colloid and Interface Science 290: 28-38.
- Srivastava, P., Singh, B., and Angrove, M. 2005. Competitive adsorption behavior of heavy metals on kaolinite. Journal of Colloid and Interface Science 290: 28-38.

- Strawn, D.G., Scheidegger, A.M., and Sparks, D.L. 1996. Kinetics and mechanisms of Pb (II) sorption and desorption at the aluminum oxide: water interface. Environmental Science and Technology 32:2596-2601.
- Stumm, W., and Morgan, J.J. 1995. Aquatic Chemistry. A Wiley-Interscience Publication, John Wiley and Sons, Inc., NY.
- Tarras-Wahlberg, N.H., and Nguyen, L.Y. 2006. Environmental regulatory failure and metal contamination at the Giap Lai pyrite mine, Northern Vietnam. Journal of Environmental Management (Article in press).
- Thomas, G.W. 1982. Exchangeable cations. In: A.L. Page, et al., (Eds.), Methods of Soil Analysis, Vol. 2. Chemical and Microbiological Properties. Madison, WI: American Societies of Agronomy, Soil Science Society of American.
- Toride, N., Leij, F.J., and van Genuchten, M.T. 1999. The CXTFIT code for estimating transport parameters from laboratory or field tracer experiments. Version 2.1. US Salinity Laboratory, Agricultural Research Service US Department of Agriculture, Riverside, California, Research Report, vol. 137.
- Trois, C., Marcello, A., Pretti, S., Trois, P., and Rossi, G. 2007. The environmental risk posed by small dumps of complex arsenic, antimony, nickel and cobalt sulphides. Journal of Geochemical exploration 92: 83-95.
- Tsang, D.C.W. and Lo, I.M. 2006. Competitive Cu and Cd sorption and transport in soils: A combined batch kinetics, column, and sequential extraction study. Environmental Science & technology 40: 6655-6661.
- URS Australia PTY LTD, 1999. Description of existing environment in Environmental Impact Assessment, Petchabun.
- van Genuchten, M.Th. and Wierenga, P.J. 1976. Mass transfer studies in sorbing porous media: I. Analytical solutions. Soil Science Society of America Journal 40: 473-481.
- Vega, F.A., Covelo, E.F., and Andrade, M.L. 2006. Competitive sorption and desorption of heavy metals in mine soils: Influence of mine soil characteristics. Journal of Colloid and Interface Science 298: 582-592.
- Voegelin, A., Barmettler, K., and Kretzchmar, R. 2003. Heavy metal release from contaminated soils: comparison of column leaching and batch extraction results. Journal of Environmental Quality 32: 865-875.
- Walkley, A. and Black, I.A. 1934. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic

- acid titration method. Soil Science 34: 29-38.
- Wilczak, A. and Keinath, T.M. 1993. Kinetics of sorption and desorption of copper (II) and lead (II) on activated carbon. Water Environment Research 65(3): 238-244.
- Williams, L.E., Mark, O.B., Kramer, T.A., and Melville, J.G. 2003. Adsorption and transport of Arsenic (V) in experimental subsurface systems. Journal of Environmental Quality 32: 841-850.
- Yamamoto, H., Morley, M.C., Speitel JR, G.E., and Clausen, J. 2004. Fate and transport of high explosive in a sandy soil. Adsorption and Desorption. Soil & Sediment Contamination 13: 361-379.
- Yolcubal, I. and Akyol, N.H. 2007. Retention and transport of hexavalent chromium in Calcareous Karst soils. Turkish Journal of Earth Sciences 16: 363-379.
- Yong, R.N., and Diperno, N. 1991. Sources and characteristics of waste with specific reference to Canada. Geotechnical research centre, McGill University. Geo-Engineering series 91-1.
- Yong, R.N. 2001. Geoenvironmental Engineering: Contaminated soils, pollution fate, and migration. Boca Raton: CRC Press.
- Yukselen, M.A. 2002. Characterization of heavy metal contaminated soils in Northern Cyprus. Environmental Geology 42: 597-603.

APPENDICES

APPENDIX A: CHEMICAL ANALYSIS OF TAILINGS

Table A.1 Chemical analysis of tailing solid samples and top soil at the Akara Mine

No.	Type	Concentration (mg g ⁻¹)																									
		Ag	Al	As	B	Ba	Bi	Ca	Cd	Co	Cr	Cu	Fe	Ga	Hg*	K	Li	Mg	Mn	Na	Ni	Pb	Sr	Tl	Zn		
1		ND	10.88	ND	0.11	0.14	ND	44.08	ND	ND	0.07	20.48	ND	0.14	2.61	ND	5.00	2.54	ND	ND	0.06	0.03	0.04	ND	0.20		
2		ND	10.71	ND	0.10	0.14	ND	45.41	ND	ND	0.07	20.28	ND	0.16	2.56	ND	4.91	2.58	ND	ND	0.08	0.03	0.04	ND	0.14		
3		ND	9.09	ND	0.09	0.14	ND	51.76	ND	ND	0.06	18.94	ND	0.14	2.49	ND	4.66	2.30	ND	ND	0.05	0.02	0.05	ND	0.09		
4		ND	12.52	ND	0.10	0.20	ND	50.51	ND	ND	0.07	19.08	ND	0.23	2.61	ND	5.13	2.69	ND	ND	0.07	0.03	0.04	ND	0.12		
5		ND	12.73	ND	0.11	0.23	ND	48.68	ND	ND	0.08	21.51	ND	0.12	2.54	ND	5.23	2.69	ND	0.48	0.05	0.03	0.05	ND	0.12		
6		ND	10.65	ND	0.09	0.29	ND	49.86	ND	ND	0.07	19.00	ND	0.28	2.47	ND	4.94	2.46	ND	ND	0.06	0.04	0.05	ND	0.15		
7		ND	9.44	ND	0.07	0.28	ND	42.60	ND	ND	0.06	16.76	ND	0.26	2.37	ND	4.71	2.14	ND	ND	0.06	0.05	0.05	ND	0.18		
8		ND	9.43	ND	0.12	0.10	ND	44.44	ND	ND	0.07	19.98	ND	1.72	2.47	ND	4.74	2.41	ND	ND	0.07	0.03	0.03	ND	0.12		
9		ND	8.27	ND	0.13	0.09	ND	40.07	ND	ND	0.09	20.86	ND	0.36	2.42	ND	4.42	2.31	ND	ND	0.08	0.04	0.03	ND	0.21		
10		ND	13.23	ND	0.17	0.17	ND	44.68	ND	ND	0.08	21.05	ND	0.16	2.79	ND	5.08	2.53	ND	ND	0.09	0.04	0.04	ND	0.21		
11		ND	7.28	ND	0.04	0.05	ND	27.23	ND	ND	0.02	11.41	ND	0.17	2.58	ND	3.86	1.67	ND	0.33	0.07	0.00	0.02	ND	0.05		
12		ND	13.52	ND	0.09	0.17	ND	46.35	ND	ND	0.07	20.06	ND	0.19	2.44	ND	5.24	2.84	ND	ND	0.08	0.03	0.04	ND	0.16		
13		ND	11.32	ND	0.10	0.12	ND	40.76	ND	ND	0.09	20.83	ND	0.19	2.42	ND	4.97	2.84	ND	ND	0.08	0.05	0.04	ND	0.22		
14		ND	10.68	ND	0.11	0.10	ND	40.08	ND	ND	0.07	18.62	ND	0.15	2.43	ND	4.85	2.66	ND	ND	0.07	0.02	0.03	ND	0.12		
15		ND	11.36	ND	0.10	0.13	ND	43.40	ND	ND	0.08	20.21	ND	0.40	2.50	ND	4.76	2.67	ND	ND	0.07	0.04	0.03	ND	0.18		
16		ND	8.97	ND	0.13	0.14	ND	41.97	ND	ND	0.08	18.92	ND	1.73	2.52	ND	4.47	2.46	ND	ND	0.05	0.03	0.03	ND	0.15		
17		ND	10.82	ND	0.10	0.09	ND	24.21	ND	ND	0.09	22.09	ND	0.25	2.75	ND	5.16	2.36	ND	ND	0.08	0.13	0.02	ND	0.28		
18		ND	10.39	ND	0.12	0.04	ND	28.60	ND	ND	0.09	22.90	ND	0.94	2.61	ND	5.36	2.39	ND	ND	0.11	0.07	0.02	ND	0.25		
19		ND	13.67	ND	0.09	0.18	ND	45.80	ND	ND	0.07	18.76	ND	0.26	2.71	ND	5.30	2.75	ND	ND	0.06	0.04	0.04	ND	0.16		
20		ND	12.70	ND	0.10	0.17	ND	42.88	ND	ND	0.08	21.43	ND	0.17	2.53	ND	5.25	2.65	ND	ND	0.06	0.05	0.04	ND	0.23		
21		ND	9.49	ND	0.07	0.15	ND	47.11	ND	ND	0.06	18.08	ND	0.34	2.40	ND	4.81	2.39	ND	ND	0.09	0.04	0.04	ND	0.15		
22		ND	10.99	ND	0.11	0.11	ND	29.10	ND	ND	0.09	22.37	ND	0.46	2.56	ND	4.41	2.37	ND	ND	0.06	0.06	0.02	ND	0.24		
1		ND	7.87	ND	0.17	0.01	ND	0.59	ND	ND	0.02	28.04	ND	0.38	0.67	ND	0.38	0.11	ND	ND	ND	ND	ND	ND	ND		
2		ND	7.73	ND	0.06	0.04	ND	1.21	ND	ND	0.06	19.12	ND	0.12	1.26	ND	1.53	0.13	ND	ND	ND	ND	0.01	ND	0.04		
3		ND	5.98	ND	0.09	0.04	ND	1.29	ND	ND	0.05	19.04	ND	0.06	1.28	ND	1.53	0.14	ND	ND	ND	ND	0.01	ND	0.04		
4		ND	8.48	ND	0.13	0.09	ND	1.88	ND	ND	0.05	24.19	ND	0.04	1.86	ND	2.01	0.18	ND	ND	ND	ND	0.02	ND	0.09		
5		ND	6.35	ND	0.02	0.04	ND	3.47	ND	ND	ND	4.83	ND	0.03	2.15	ND	1.94	0.03	ND	ND	ND	ND	0.03	ND	ND		




Figure A.1 3 pH conditions under batch desorption experiments.



Figure A.2 Tailings samplings



Figure A.3 Lateritic soil samplings

APPENDIX B: COLUMN EXPERIMENT DATA FOR SINGLE METAL SYSTEMS

Table B.1 Observed, predicted and simulated breakthrough curves of Lead (Pb) transport through lateritic soil under pH 4 with initial concentration 5.28 mmol L⁻¹

Sample #	Pore volume	Observed data Pb		Predicted		Fitted	
		Conc.(mM)	C/C0	Langmuir C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0
1	0.33	0.00	0.00	0.00	0.00	0.00	0.00
2	2.86	0.00	0.00	0.00	0.00	0.00	0.00
3	5.15	0.00	0.00	0.00	0.01	0.00	0.00
4	7.69	4.77	0.01	0.00	0.03	0.00	0.00
5	10.44	4.93	0.01	0.05	0.10	0.00	0.00
6	11.86	6.65	0.01	0.23	0.14	0.00	0.00
7	16.03	33.02	0.03	0.43	0.31	0.00	0.00
8	18.81	89.81	0.09	0.63	0.42	0.00	0.00
9	21.50	133.95	0.14	0.78	0.52	0.00	0.02
10	24.24	215.70	0.23	0.87	0.61	0.00	0.16
11	26.97	316.60	0.33	0.93	0.69	0.00	0.42
12	29.72	424.65	0.45	0.96	0.75	0.00	0.54
13	32.48	512.50	0.54	0.98	0.80	0.47	0.62
14	35.28	587.50	0.62	0.99	0.85	0.99	0.68
15	38.04	651.50	0.69	0.99	0.88	1.00	0.74
16	40.79	703.50	0.74	1.00	0.91	1.00	0.78
17	43.52	798.50	0.84	1.00	0.93	1.00	0.82
18	46.25	828.50	0.87	1.00	0.94	1.00	0.85
19	49.00	846.00	0.89	1.00	0.95	1.00	0.87
20	51.80	879.50	0.93	1.00	0.96	1.00	0.89
21	53.46	886.50	0.94	1.00	0.97	1.00	0.90
22	55.95	896.00	0.95	1.00	0.98	1.00	0.92
23	58.73	907.50	0.96	1.00	0.98	1.00	0.93
24	61.49	923.50	0.98	1.00	0.99	1.00	0.94
25	64.29	933.50	0.99	1.00	0.99	1.00	0.95
26	67.06	946.50	1.00	1.00	0.99	1.00	0.96
27	69.86	919.50	0.97	1.00	0.99	1.00	0.97
28	72.66	943.50	1.00	1.00	0.99	1.00	0.97
29	75.46	956.00	1.01	1.00	1.00	1.00	0.98
30	78.22	946.00	1.00	1.00	1.00	1.00	0.98
31	80.96	964.50	1.02	1.00	1.00	1.00	0.98
32	83.73	991.00	1.05	1.00	1.00	0.99	0.98
33	84.26	999.00	1.06	1.00	1.00	0.90	0.90
34	84.54	978.00	1.03	1.00	1.00	0.83	0.84
35	84.81	922.00	0.97	0.99	1.00	0.77	0.78
36	85.07	778.50	0.82	0.99	1.00	0.71	0.73
37	85.33	673.50	0.71	0.98	1.00	0.66	0.68
38	85.85	495.50	0.52	0.86	1.00	0.58	0.61
39	86.38	409.40	0.43	0.71	1.00	0.52	0.55
40	88.97	295.90	0.31	0.60	0.98	0.36	0.40
41	91.81	245.85	0.26	0.52	0.93	0.29	0.33
42	94.13	205.95	0.22	0.42	0.85	0.25	0.30
43	96.22	180.85	0.19	0.36	0.77	0.22	0.28

Table B.2 Observed, predicted and simulated breakthrough curves of Zinc (Zn) transport through lateritic soil under pH 4 with initial concentration 4.67 mmol/L

Sample #	Pore volume	Observed data Zn		Predicted		Fitted	TSM C/C0
		Conc.(mM)	C/C0	Langmuir C/C0	Linear C/C0	Langmuir C/C0	
1	0.00	0.00	0.00	0.00	0.02	0.00	0.00
2	3.11	0.08	0.02	0.00	0.03	0.00	0.00
3	4.06	0.09	0.02	0.00	0.07	0.00	0.00
4	7.45	0.43	0.09	0.02	0.33	0.00	0.00
5	10.88	1.24	0.27	0.16	0.58	0.00	0.24
6	13.17	1.99	0.43	0.34	0.71	0.42	0.48
7	16.90	2.52	0.54	0.66	0.83	0.99	0.61
8	20.06	3.01	0.64	0.83	0.89	1.00	0.68
9	23.84	3.50	0.75	0.93	0.93	1.00	0.73
10	27.01	3.76	0.80	0.97	0.95	1.00	0.77
11	33.90	3.85	0.82	0.99	0.96	1.00	0.84
12	38.51	4.37	0.94	1.00	0.96	1.00	0.87
13	41.67	4.29	0.92	1.00	0.96	1.00	0.89
14	47.72	4.33	0.93	1.00	0.97	1.00	0.92
15	51.28	4.28	0.92	1.00	0.97	1.00	0.93
16	54.34	4.42	0.95	1.00	0.97	1.00	0.94
17	58.00	4.46	0.96	1.00	0.97	1.00	0.95
18	61.04	4.40	0.94	1.00	0.97	1.00	0.96
19	64.54	4.39	0.94	1.00	0.97	1.00	0.97
20	67.45	4.58	0.98	1.00	0.97	1.00	0.97
21	73.79	4.43	0.95	1.00	0.97	1.00	0.98
22	78.00	4.38	0.94	1.00	0.97	1.00	0.98
23	79.06	4.31	0.92	1.00	0.96	0.94	0.92
24	79.58	4.29	0.92	0.99	0.96	0.83	0.81
25	80.10	4.11	0.88	0.98	0.95	0.73	0.71
26	80.62	3.07	0.66	0.96	0.94	0.65	0.63
27	81.13	2.30	0.49	0.94	0.92	0.58	0.57
28	81.89	2.07	0.44	0.89	0.88	0.51	0.50
29	82.66	1.90	0.41	0.84	0.83	0.45	0.45
30	83.17	1.83	0.39	0.81	0.79	0.42	0.42
31	83.93	1.69	0.36	0.76	0.72	0.38	0.39
32	86.38	1.32	0.28	0.60	0.49	0.30	0.32
33	88.70	1.06	0.23	0.47	0.31	0.24	0.28
34	91.01	0.93	0.20	0.37	0.19	0.21	0.26
35	93.25	0.80	0.17	0.30	0.11	0.18	0.24
36	95.45	0.71	0.15	0.24	0.07	0.16	0.22
37	97.58	0.64	0.14	0.20	0.04	0.14	0.20
38	99.62	0.61	0.13	0.16	0.02	0.13	0.19

Table B.3 Observed, predicted and simulated breakthrough curves of Nickel (Ni) transport through lateritic soil under pH 4 with initial concentration 5.06 mmol/L

Sample #	Pore volume	Observed data Ni		Predicted		Fitted	
		Conc.(mM)	C/C0	Langmuir C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0
1	0.28	0.00	0.00	0.00	0.01	0.00	0.00
2	0.55	0.05	0.01	0.00	0.03	0.00	0.00
3	0.83	0.25	0.05	0.00	0.12	0.00	0.01
4	1.11	0.45	0.09	0.00	0.15	0.00	0.01
5	1.39	0.58	0.11	0.00	0.22	0.00	0.04
6	1.66	0.78	0.15	0.00	0.25	0.00	0.06
7	1.94	1.22	0.24	0.00	0.43	0.00	0.24
8	2.22	1.99	0.39	0.00	0.68	0.01	0.50
9	2.50	2.36	0.47	0.00	0.77	0.50	0.58
10	2.77	3.38	0.67	0.00	0.89	0.99	0.71
11	3.05	4.11	0.81	0.00	0.95	1.00	0.79
12	3.33	4.61	0.91	0.00	0.97	1.00	0.85
13	3.61	4.58	0.91	0.00	0.99	1.00	0.90
14	3.88	4.76	0.94	0.00	0.99	1.00	0.93
15	4.16	4.80	0.95	0.00	1.00	1.00	0.96
16	4.44	4.95	0.98	0.00	1.00	1.00	0.97
17	4.71	4.96	0.98	0.00	1.00	1.00	0.98
18	4.99	4.92	0.97	0.00	1.00	1.00	0.98
19	5.27	4.98	0.98	0.00	1.00	1.00	0.99
20	5.55	5.07	1.00	0.00	1.00	1.00	0.99
21	5.82	4.72	0.93	0.00	1.00	1.00	0.99
22	6.10	4.90	0.97	0.00	1.00	0.91	0.80
23	6.38	4.10	0.81	0.01	0.99	0.68	0.60
24	6.66	2.57	0.51	0.01	0.98	0.58	0.52
25	6.93	2.12	0.42	0.01	0.95	0.51	0.47
26	7.21	1.59	0.31	0.02	0.87	0.44	0.41
27	7.49	0.88	0.17	0.02	0.54	0.30	0.32
28	7.76	0.62	0.12	0.03	0.28	0.23	0.27
29	8.04	0.48	0.10	0.04	0.14	0.19	0.24
30	8.32	0.56	0.11	0.05	0.07	0.16	0.21
31	8.60	0.44	0.09	0.07	0.03	0.14	0.18

Table B.4 Observed, predicted and simulated breakthrough curves of Manganese (Mn) transport through lateritic soil under pH 4 with initial concentration 4.87 mmol/L

Sample #	Pore volume	Observed data Mn		Predicted		Fitted	TSM C/C0
		Conc.(mM)	C/C0	Langmuir C/C0	Linear C/C0	Langmuir C/C0	
1	1.29	0.06	0.01	0.00	0.02	0.00	0.00
2	2.68	0.04	0.01	0.00	0.06	0.00	0.00
3	3.73	0.09	0.02	0.01	0.29	0.00	0.01
4	7.06	0.74	0.15	0.25	0.48	0.02	0.26
5	9.52	1.42	0.29	0.71	0.66	0.51	0.49
6	12.41	1.90	0.39	0.94	0.83	0.97	0.62
7	16.36	2.56	0.53	0.99	0.88	0.99	0.67
8	18.59	2.70	0.55	1.00	0.94	1.00	0.75
9	22.53	3.21	0.66	1.00	0.96	1.00	0.79
10	25.03	3.38	0.70	1.00	0.98	1.00	0.82
11	27.93	4.15	0.85	1.00	0.99	1.00	0.86
12	31.08	4.47	0.92	1.00	0.99	1.00	0.88
13	34.25	4.53	0.93	1.00	1.00	1.00	0.93
14	37.57	4.48	0.92	1.00	1.00	1.00	0.94
15	41.26	4.70	0.97	1.00	1.00	1.00	0.95
16	44.17	4.75	0.98	1.00	1.00	1.00	0.96
17	46.73	4.80	0.99	1.00	1.00	1.00	0.97
18	51.20	4.92	1.01	1.00	1.00	1.00	0.95
19	52.46	4.93	1.01	1.00	1.00	0.94	0.72
20	57.14	4.90	1.01	1.00	0.99	0.84	0.55
21	57.89	5.16	1.06	0.97	0.98	0.74	0.45
22	58.67	3.53	0.73	0.89	0.96	0.66	0.39
23	59.44	2.12	0.43	0.80	0.70	0.41	0.28
24	60.21	1.62	0.33	0.71	0.53	0.34	0.25
25	63.26	1.07	0.22	0.40	0.31	0.27	0.22
26	65.35	0.85	0.18	0.30	0.19	0.22	0.20
27	68.64	0.67	0.14	0.21	0.11	0.18	0.19
28	71.19	0.57	0.12	0.15	0.06	0.16	0.17
29	74.18	0.49	0.10	0.11	0.03	0.13	0.16
30	76.93	0.44	0.09	0.08	0.02	0.12	0.15
31	79.36	0.41	0.08	0.06	0.00	0.00	0.00
32	81.31	0.37	0.08	0.04	0.00	0.00	0.00

Table B.5 Observed and simulated breakthrough curves of Lead (Pb) transport through lateritic soil under pH 5 with initial concentration 5.05 mol/L

Sample #	Pore volume	Observed data Pb		Linear C/C0	Fitted	TSM C/C0
		Conc.(mM)	C/C0		Langmuir C/C0	
1	1.76	0.00	0.00	0.00	0.00	0.00
2	5.24	0.00	0.00	0.00	0.00	0.00
3	7.84	0.00	0.00	0.01	0.00	0.00
4	10.38	0.00	0.00	0.04	0.00	0.00
5	11.30	0.00	0.00	0.10	0.00	0.00
6	13.85	0.07	0.01	0.16	0.00	0.00
7	16.36	0.45	0.09	0.24	0.00	0.00
8	18.88	0.98	0.19	0.31	0.00	0.03
9	21.42	1.07	0.21	0.39	0.00	0.20
10	23.96	1.47	0.29	0.46	0.00	0.36
11	25.21	1.83	0.36	0.49	0.00	0.40
12	27.73	1.97	0.39	0.55	0.00	0.47
13	30.25	2.21	0.44	0.61	0.00	0.52
14	32.76	2.54	0.50	0.65	0.62	0.57
15	35.27	2.92	0.58	0.70	1.00	0.61
16	37.75	2.94	0.58	0.73	1.00	0.65
17	40.25	3.25	0.64	0.77	1.00	0.69
18	42.75	3.38	0.67	0.80	1.00	0.72
19	45.25	3.71	0.73	0.82	1.00	0.75
20	47.75	3.90	0.77	0.84	1.00	0.77
21	48.98	3.94	0.78	0.85	1.00	0.78
22	51.39	4.05	0.80	0.87	1.00	0.81
23	53.86	4.40	0.87	0.89	1.00	0.83
24	56.36	4.09	0.81	0.90	1.00	0.84
25	58.87	4.44	0.88	0.91	1.00	0.86
26	61.32	4.66	0.92	0.92	1.00	0.87
27	63.79	4.52	0.89	0.93	1.00	0.89
28	66.30	4.54	0.90	0.94	1.00	0.90
29	68.79	4.73	0.94	0.94	1.00	0.91
30	72.58	4.86	0.96	0.95	1.00	0.92
31	75.10	4.89	0.97	0.95	1.00	0.93
32	77.37	4.73	0.94	0.96	1.00	0.94
33	80.17	4.91	0.97	0.96	1.00	0.94
34	82.72	4.89	0.97	0.96	1.00	0.95
35	85.24	4.90	0.97	0.97	1.00	0.95
36	87.78	5.04	1.00	0.97	1.00	0.96
37	90.32	5.09	1.01	0.97	1.00	0.96
38	92.82	5.58	1.11	0.97	1.00	0.97
39	93.33	5.59	1.11	0.97	1.00	0.97
40	93.84	5.61	1.11	0.97	1.00	0.97
41	94.10	5.60	1.11	0.97	1.00	0.97
42	94.36	5.42	1.07	0.97	0.99	0.96
43	94.89	4.60	0.91	0.97	0.85	0.84
44	95.41	3.72	0.74	0.97	0.68	0.69
45	95.93	3.07	0.61	0.97	0.57	0.58
46	96.71	2.55	0.50	0.96	0.47	0.48
47	98.02	2.13	0.42	0.95	0.37	0.39
48	99.31	1.82	0.36	0.94	0.31	0.34

Table B.5 Observed and simulated breakthrough curves of Lead (Pb) transport through lateritic soil under pH 5 with initial concentration 5.05 mol/L (Continued)

49	101.94	1.40	0.28	0.89	0.25	0.28
50	104.44	1.12	0.22	0.82	0.21	0.25
51	107.25	0.97	0.19	0.74	0.18	0.23
52	109.55	0.87	0.17	0.67	0.17	0.22
53	112.11	0.79	0.16	0.60	0.15	0.21
54	114.67	0.75	0.15	0.53	0.14	0.20
55	117.23	0.72	0.14	0.47	0.13	0.19
56	121.08	0.69	0.14	0.38	0.12	0.18
57	123.61	0.68	0.13	0.33	0.11	0.18
58	126.16	0.68	0.13	0.29	0.11	0.17
59	128.68	0.65	0.13	0.25	0.10	0.16
60	131.20	0.64	0.13	0.22	0.10	0.16
61	133.73	0.64	0.13	0.19	0.10	0.15
62	136.25	0.62	0.12	0.16	0.09	0.15
63	138.78	0.59	0.12	0.14	0.09	0.15
64	141.27	0.60	0.12	0.12	0.09	0.14
65	144.96	0.55	0.11	0.10	0.08	0.14
66	147.40	0.56	0.11	0.08	0.08	0.13
67	149.84	0.56	0.11	0.07	0.08	0.13
68	152.25	0.54	0.11	0.06	0.07	0.13
69	154.64	0.52	0.10	0.05	0.07	0.12
70	156.99	0.50	0.10	0.05	0.07	0.12
71	159.32	0.48	0.09	0.04	0.07	0.12
72	161.59	0.47	0.09	0.04	0.07	0.11
73	163.80	0.43	0.09	0.03	0.07	0.11
74	164.02	0.39	0.08	0.03	0.07	0.11
75	169.26	0.39	0.08	0.02	0.06	0.11
76	173.45	0.35	0.07	0.02	0.06	0.10
77	175.51	0.33	0.07	0.01	0.06	0.10

Table B.6 Observed and simulated breakthrough curves of Lead (Pb) transport through lateritic soil under pH 5 with initial concentration 5.01 mol/L (Duplicated Column)

Sample #	Pore volume	Observed data Pb		Linear C/C0	Fitted	TSM C/C0
		Conc.(mM)	C/C0		Langmuir C/C0	
1	0.00	0.00	0.00	0.00	0.00	0.00
2	3.24	0.00	0.00	0.00	0.00	0.00
3	10.08	0.00	0.00	0.03	0.00	0.00
4	17.62	0.00	0.00	0.23	0.00	0.00
5	20.16	0.07	0.01	0.33	0.00	0.00
6	23.22	0.29	0.06	0.44	0.00	0.00
7	27.39	0.94	0.19	0.59	0.00	0.14
8	31.26	1.64	0.33	0.70	0.34	0.51
9	36.58	2.63	0.53	0.81	1.00	0.62
10	40.73	3.20	0.64	0.87	1.00	0.68
11	43.26	3.67	0.73	0.90	1.00	0.71
12	49.56	4.04	0.81	0.95	1.00	0.78
13	51.68	4.24	0.85	0.96	1.00	0.80
14	59.74	4.55	0.91	0.98	1.00	0.86
15	64.22	4.75	0.95	0.99	1.00	0.89
16	69.25	4.75	0.95	0.99	1.00	0.91
17	72.56	4.73	0.94	1.00	1.00	0.93
18	76.11	4.69	0.94	1.00	1.00	0.94
19	81.70	4.80	0.96	1.00	1.00	0.95
20	85.68	4.92	0.98	1.00	1.00	0.96
21	89.48	4.78	0.96	1.00	1.00	0.96
22	95.71	4.77	0.95	1.00	1.00	0.97
23	97.58	4.76	0.95	1.00	1.00	0.98
24	98.64	4.71	0.94	1.00	0.96	0.93
25	99.17	4.65	0.93	1.00	0.87	0.83
26	99.68	4.62	0.92	1.00	0.77	0.72
27	100.20	3.92	0.78	1.00	0.69	0.64
28	100.72	2.97	0.59	1.00	0.63	0.58
29	101.23	2.53	0.51	1.00	0.57	0.53
30	101.74	2.31	0.46	1.00	0.53	0.49
31	102.26	2.16	0.43	1.00	0.50	0.46
32	103.58	1.80	0.36	1.00	0.43	0.40
33	104.90	1.47	0.29	1.00	0.38	0.36
34	107.54	1.01	0.20	0.98	0.32	0.31
35	110.15	0.80	0.16	0.93	0.28	0.28
36	112.80	0.70	0.14	0.86	0.25	0.26
37	115.40	0.59	0.12	0.76	0.22	0.24
38	119.32	0.72	0.14	0.50	0.18	0.21
39	121.89	0.66	0.13	0.41	0.17	0.20
40	124.44	0.65	0.13	0.33	0.16	0.19
41	127.00	0.65	0.13	0.05	0.12	0.15
42	129.58	0.66	0.13	0.02	0.10	0.14
43	132.12	0.68	0.13	0.02	0.10	0.14
44	134.60	0.67	0.13	0.01	0.10	0.13
45	137.11	0.65	0.13	0.01	0.09	0.13
46	139.56	0.71	0.14	0.01	0.09	0.12
47	142.99	0.66	0.13	0.00	0.09	0.12
48	145.39	0.60	0.12	0.00	0.09	0.12
49	147.75	0.63	0.13	0.00	0.00	0.00

Table B.6 Observed and simulated breakthrough curves of Lead (Pb) transport through lateritic soil under pH 5 with initial concentration 5.01 mol/L (Duplicated Column) (Continued)

50	150.05	0.62	0.12	0.00	0.00	0.00
51	152.32	0.54	0.11	0.00	0.00	0.00
52	154.52	0.48	0.10	0.00	0.00	0.00
53	156.63	0.44	0.09	0.00	0.00	0.00
54	158.69	0.40	0.08	0.00	0.00	0.00
55	160.62	0.33	0.07	0.00	0.00	0.00
56	162.74	0.30	0.06	0.00	0.00	0.00
57	164.66	0.26	0.05	0.00	0.00	0.00

Table B.7 Observed and simulated breakthrough curves of Zinc (Zn) transport through lateritic soil under pH 5 with initial concentration 4.61 mol/L

Sample #	Pore volume	Observed data Zn		Linear C/C0	Fitted	
		Conc.(mM)	C/C0		Langmuir C/C0	TSM C/C0
1	2.60	0.10	0.02	0.00	0.00	0.00
2	4.63	0.09	0.02	0.01	0.00	0.00
3	6.96	0.16	0.03	0.04	0.00	0.00
4	11.58	0.47	0.10	0.19	0.00	0.02
5	16.48	1.09	0.24	0.41	0.00	0.20
6	19.67	1.55	0.34	0.54	0.01	0.36
7	23.43	2.02	0.44	0.67	0.34	0.49
8	26.50	2.35	0.51	0.75	0.89	0.56
9	30.34	2.77	0.60	0.82	0.99	0.64
10	33.55	3.00	0.65	0.87	1.00	0.69
11	37.27	3.25	0.70	0.91	1.00	0.74
12	40.64	3.56	0.77	0.93	1.00	0.78
13	44.18	3.79	0.82	0.95	1.00	0.81
14	47.65	3.90	0.85	0.97	1.00	0.84
15	51.44	3.99	0.86	0.98	1.00	0.86
16	54.66	4.11	0.89	0.98	1.00	0.88
17	58.41	4.19	0.91	0.99	1.00	0.90
18	61.64	4.20	0.91	0.99	1.00	0.92
19	66.47	4.39	0.95	0.99	1.00	0.93
20	68.78	4.35	0.94	1.00	1.00	0.94
21	73.59	4.46	0.97	1.00	1.00	0.95
22	75.92	4.44	0.96	1.00	1.00	0.96
23	79.80	4.58	0.99	1.00	1.00	0.96
24	82.82	4.58	0.99	1.00	1.00	0.97
25	86.44	4.61	1.00	1.00	1.00	0.97
26	88.44	4.62	1.05	1.00	1.00	0.98
27	90.73	4.61	1.02	1.00	1.00	0.98
28	94.18	4.71	1.03	1.00	1.00	0.98
29	96.39	4.65	1.07	1.00	0.94	0.91
30	96.95	4.60	1.05	1.00	0.89	0.85
31	97.51	4.62	1.04	1.00	0.83	0.80
32	98.06	3.76	0.82	1.00	0.78	0.75
33	98.61	2.97	0.64	0.99	0.73	0.70
34	98.88	2.84	0.62	0.99	0.71	0.68
35	101.60	1.89	0.41	0.95	0.54	0.54
36	104.26	1.49	0.32	0.86	0.44	0.46
37	105.92	1.36	0.30	0.78	0.39	0.42
38	107.22	1.21	0.26	0.65	0.34	0.38
39	109.19	1.13	0.25	0.48	0.29	0.34

Table B.8 Observed and simulated breakthrough curves of Nickel (Ni) transport through lateritic soil under pH 5 with initial concentration 4.89 mol/L

Sample #	Pore volume	Observed data Ni		Linear C/C0	Fitted	TSM C/C0
		Conc.(mM)	C/C0		Langmuir C/C0	
1	2.40	0.00	0.00	0.00	0.00	0.00
2	2.67	0.00	0.00	0.00	0.00	0.00
3	4.30	0.00	0.00	0.03	0.00	0.00
4	4.57	0.00	0.00	0.03	0.00	0.00
5	10.69	0.65	0.05	0.37	0.00	0.00
6	13.08	0.80	0.19	0.52	0.00	0.00
7	15.52	0.95	0.29	0.65	0.00	0.02
8	18.35	1.14	0.38	0.77	0.37	0.38
9	21.87	1.36	0.46	0.86	1.00	0.53
10	24.82	1.55	0.52	0.91	1.00	0.58
11	28.13	1.77	0.58	0.95	1.00	0.62
12	31.04	1.96	0.65	0.97	1.00	0.65
13	33.84	2.15	0.68	0.98	1.00	0.68
14	37.10	2.37	0.72	0.99	1.00	0.71
15	40.34	2.59	0.75	0.99	1.00	0.74
16	43.08	2.78	0.78	1.00	1.00	0.76
17	46.35	3.00	0.80	1.00	1.00	0.79
18	49.11	3.19	0.79	1.00	1.00	0.80
19	53.10	3.46	0.84	1.00	1.00	0.83
20	54.98	3.59	0.84	1.00	1.00	0.84
21	58.97	3.87	0.85	1.00	1.00	0.86
22	60.94	4.00	0.88	1.00	1.00	0.87
23	64.18	4.24	0.88	1.00	1.00	0.88
24	66.81	4.42	0.86	1.00	1.00	0.89
25	70.63	4.68	0.91	1.00	1.00	0.90
26	72.61	4.82	0.92	1.00	1.00	0.91
27	79.32	5.30	0.94	1.00	1.00	0.93
28	85.01	5.71	1.00	1.00	1.00	0.94
29	89.71	6.05	1.02	1.00	1.00	0.95
30	90.32	6.09	1.01	1.00	0.77	0.73
31	90.80	6.13	0.76	1.00	0.65	0.61
32	91.05	6.14	0.60	1.00	0.60	0.57
33	91.28	6.16	0.60	1.00	0.56	0.53
34	92.22	6.23	0.40	0.98	0.45	0.43
35	92.68	6.26	0.36	0.97	0.41	0.39
36	93.38	6.31	0.29	0.95	0.37	0.35
37	93.85	6.35	0.29	0.92	0.35	0.33
38	94.55	6.40	0.24	0.89	0.32	0.31
39	96.90	6.57	0.19	0.72	0.25	0.25
40	99.19	6.74	0.16	0.54	0.21	0.22
41	101.45	6.91	0.14	0.39	0.19	0.21
42	103.66	7.08	0.13	0.27	0.17	0.19
43	105.88	7.25	0.12	0.19	0.15	0.18
44	108.07	7.42	0.11	0.13	0.14	0.17
45	110.23	7.59	0.11	0.08	0.13	0.16
46	113.49	7.85	0.10	0.05	0.11	0.16
47	115.62	8.02	0.10	0.03	0.11	0.15
48	117.77	8.19	0.10	0.02	0.10	0.15
49	119.87	8.36	0.09	0.01	0.10	0.14

Table B.8 Observed and simulated breakthrough curves of Nickel (Ni) transport through lateritic soil under pH 5 with initial concentration 4.89 mol/L (Continued)

50	121.93	8.53	0.08	0.01	0.09	0.14
51	123.88	8.70	0.08	0.01	0.09	0.13
52	125.51	8.85	0.08	0.00	0.00	0.00
36	93.38	6.31	0.29	0.95	0.37	0.35
37	93.85	6.35	0.29	0.92	0.35	0.33
38	94.55	6.40	0.24	0.89	0.32	0.31

Table B.9 Observed and simulated breakthrough curves of Manganese (Mn) transport through lateritic soil under pH 5 with initial concentration 4.81 mol/L

Sample #	Pore volume	Observed data Mn		Linear C/C0	Fitted	TSM C/C0
		Conc.(mM)	C/C0		Langmuir C/C0	
1	1.67	0.00	0.00	0.00	0.00	0.00
2	2.78	0.00	0.00	0.03	0.00	0.00
3	5.57	0.04	0.01	0.18	0.00	0.03
4	6.12	0.59	0.12	0.22	0.00	0.05
5	6.97	1.04	0.22	0.28	0.00	0.11
6	8.36	1.17	0.24	0.39	0.00	0.24
7	9.77	1.39	0.29	0.48	0.00	0.36
8	12.57	1.96	0.41	0.65	0.21	0.50
9	13.98	2.22	0.46	0.71	0.61	0.54
10	16.79	2.56	0.53	0.81	0.95	0.61
11	19.56	3.14	0.65	0.87	0.99	0.66
12	22.37	3.45	0.72	0.91	1.00	0.71
13	25.15	3.57	0.74	0.94	1.00	0.75
14	29.21	3.96	0.82	0.97	1.00	0.81
15	31.95	4.05	0.84	0.98	1.00	0.83
16	34.69	4.20	0.87	0.99	1.00	0.86
17	37.46	4.25	0.88	0.99	1.00	0.88
18	40.19	4.37	0.91	0.99	1.00	0.90
19	42.98	4.53	0.94	1.00	1.00	0.91
20	45.78	4.54	0.94	1.00	1.00	0.92
21	48.62	4.62	0.96	1.00	1.00	0.94
22	51.47	4.69	0.97	1.00	1.00	0.94
23	55.63	4.51	0.94	1.00	1.00	0.96
24	58.43	4.51	0.94	1.00	1.00	0.96
25	61.24	4.69	0.98	1.00	1.00	0.97
26	64.06	4.62	0.96	1.00	1.00	0.97
27	66.85	4.66	0.97	1.00	1.00	0.98
28	69.66	4.71	0.98	1.00	1.00	0.98
29	72.45	4.69	0.98	1.00	1.00	0.98
30	75.26	4.73	0.98	1.00	1.00	0.99
31	77.77	4.64	0.96	1.00	1.00	0.99
32	78.32	4.66	0.97	1.00	1.00	0.99
33	80.77	4.62	0.96	0.99	0.78	0.69
34	81.31	3.42	0.71	0.98	0.70	0.61
35	82.13	2.19	0.46	0.95	0.60	0.52
36	83.49	1.58	0.33	0.88	0.49	0.44
37	84.85	1.25	0.26	0.79	0.42	0.38
38	87.56	0.91	0.19	0.58	0.32	0.32
39	90.28	0.72	0.15	0.40	0.27	0.29
40	93.00	0.61	0.13	0.27	0.23	0.26
41	95.71	0.54	0.11	0.18	0.20	0.24
42	98.33	0.50	0.10	0.12	0.17	0.23
43	100.79	0.44	0.09	0.08	0.15	0.21
44	103.00	0.41	0.09	0.05	0.14	0.20

Table B.10 Observed, predicted and simulated breakthrough curves of Nickel (Ni) transport through lateritic soil under pH 4 with initial concentration 5.28 mmol/L

Sample #	Pore volume	Observed data Pb		Predicted		Fitted	
		Conc.(mM)	C/C0	Langmuir C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0
1	0.00	0.00	0.00	0.00	0.01	0.00	0.00
2	1.60	0.01	0.00	0.00	0.02	0.00	0.00
3	2.91	0.16	0.03	0.00	0.08	0.00	0.01
4	5.22	0.88	0.17	0.00	0.28	0.01	0.12
5	8.20	2.18	0.41	0.08	0.53	0.22	0.42
6	9.68	2.57	0.49	0.24	0.64	0.53	0.54
7	10.61	2.94	0.56	0.40	0.71	0.71	0.60
8	13.30	3.68	0.70	0.77	0.83	0.94	0.72
9	14.51	4.00	0.76	0.87	0.87	0.97	0.76
10	18.97	4.52	0.86	0.98	0.95	1.00	0.87
11	21.65	4.53	0.86	0.99	0.97	1.00	0.91
12	24.55	5.15	0.97	1.00	0.99	1.00	0.94
13	26.87	5.27	1.00	1.00	0.99	1.00	0.96
14	29.35	5.28	1.00	1.00	1.00	1.00	0.97
15	31.30	5.32	1.01	1.00	1.00	1.00	0.98
16	33.41	5.40	1.02	1.00	1.00	1.00	0.99
17	35.70	5.45	1.03	1.00	1.00	1.00	0.99
18	37.48	5.49	1.04	1.00	1.00	1.00	0.99
19	39.03	5.50	1.04	1.00	1.00	1.00	1.00
20	43.20	5.52	1.04	1.00	1.00	1.00	1.00
21	45.52	5.48	1.04	1.00	1.00	1.00	1.00
22	46.60	5.49	1.04	1.00	1.00	1.00	1.00
23	49.03	5.48	1.04	1.00	1.00	1.00	1.00
24	49.36	5.91	1.12	1.00	1.00	1.00	1.00
25	49.82	5.98	1.13	1.00	1.00	0.99	0.94
26	50.29	4.70	0.89	0.99	0.99	0.93	0.81
27	50.91	3.34	0.63	0.94	0.95	0.82	0.66
28	52.45	1.95	0.37	0.70	0.77	0.58	0.47
29	53.15	1.67	0.32	0.61	0.65	0.49	0.42
30	54.63	1.37	0.26	0.46	0.43	0.37	0.35
31	57.76	1.02	0.19	0.28	0.15	0.22	0.25
32	59.48	0.94	0.18	0.22	0.08	0.17	0.22
33	60.95	0.86	0.16	0.18	0.04	0.14	0.19
34	62.89	0.88	0.17	0.14	0.02	0.11	0.16

APPENDIX C: COLUMN EXPERIMENT DATA FOR BINARY AND MULTI-METAL SYSTEM

Table C.1 Observed, predicted and simulated breakthrough curves of Lead (Pb^{2+}) and Nickel (Ni^{2+}) transport through lateritic soil under pH 5 with initial concentration 4.86 mmol/L for Pb^{2+} and 3.01 mmol/L for Ni^{2+}

Sample #	Observed data Pb				Fitted			Observed data Ni				Fitted		
	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	
1	1.08	0.00	0.00	0.00	0.00	0.00	1	1.08	0.00	0.00	0.00	0.00	0.00	
2	3.52	0.00	0.00	0.01	0.00	0.00	2	3.52	0.00	0.00	0.03	0.00	0.00	
3	5.42	0.00	0.00	0.04	0.00	0.00	3	5.42	0.00	0.00	0.11	0.00	0.00	
4	7.61	0.00	0.00	0.12	0.00	0.00	4	7.61	0.16	0.05	0.26	0.00	0.00	
5	10.08	0.11	0.02	0.26	0.00	0.00	5	10.08	0.63	0.21	0.45	0.00	0.00	
6	10.91	0.36	0.08	0.31	0.00	0.00	6	10.91	0.76	0.25	0.51	0.00	0.00	
7	13.11	0.71	0.15	0.44	0.00	0.00	7	13.11	0.83	0.28	0.64	0.00	0.00	
8	16.70	1.35	0.29	0.62	0.00	0.27	8	14.22	1.16	0.39	0.70	0.00	0.00	
9	19.73	1.97	0.42	0.73	0.43	0.50	9	16.70	1.66	0.55	0.80	0.56	0.00	
10	23.43	2.56	0.55	0.83	1.00	0.59	10	19.73	2.12	0.70	0.88	1.00	0.00	
11	26.48	2.96	0.63	0.88	1.00	0.66	11	23.43	2.43	0.81	0.93	1.00	0.00	
12	30.12	3.33	0.71	0.92	1.00	0.72	12	26.48	2.57	0.85	0.96	1.00	0.00	
13	33.19	3.51	0.75	0.95	1.00	0.77	13	30.12	2.73	0.91	0.98	1.00	0.00	
14	36.81	3.74	0.80	0.97	1.00	0.81	14	33.19	2.79	0.93	0.99	1.00	0.00	
15	39.59	3.89	0.83	0.98	1.00	0.84	15	36.81	2.90	0.96	0.99	1.00	0.00	
16	43.37	4.10	0.88	0.99	1.00	0.87	16	39.59	2.88	0.96	1.00	1.00	0.00	
17	46.96	4.16	0.89	0.99	1.00	0.90	17	43.37	2.98	0.99	1.00	1.00	0.00	
18	49.86	4.45	0.95	1.00	1.00	0.92	18	46.96	2.91	0.97	1.00	1.00	0.00	
19	52.85	4.46	0.95	1.00	1.00	0.93	19	49.86	2.96	0.98	1.00	1.00	0.00	
20	56.40	4.56	0.98	1.00	1.00	0.95	20	52.85	2.97	0.99	1.00	1.00	0.00	
21	59.47	4.54	0.97	1.00	1.00	0.95	21	56.40	3.00	1.00	1.00	1.00	0.00	
22	63.04	4.60	0.98	1.00	1.00	0.96	22	59.47	2.96	0.98	1.00	1.00	0.00	
23	66.07	4.57	0.98	1.00	1.00	0.97	23	63.04	2.99	0.99	1.00	1.00	0.00	

Table C.1 Observed, predicted and simulated breakthrough curves of Lead (Pb^{2+}) and Nickel (Ni^{2+}) transport through lateritic soil under pH 5 with initial concentration 4.86 mmol/L for Pb^{2+} and 3.01 mmol/L for Ni^{2+} (Continued)

24	69.69	4.69	1.00	1.00	1.00	0.97	24	66.07	3.05	1.01	1.00	1.00	0.00
25	72.71	4.77	1.02	1.00	1.00	0.98	25	69.69	3.01	1.00	1.00	1.00	0.00
26	76.27	4.77	1.02	1.00	1.00	0.98	26	72.71	3.07	1.02	1.00	1.00	0.00
27	78.20	4.90	1.05	1.00	1.00	0.98	27	76.27	3.04	1.01	1.00	1.00	0.00
28	82.89	4.80	1.03	1.00	0.99	0.99	28	78.20	3.02	1.00	1.00	1.00	0.00
29	84.64	3.03	0.65	1.00	0.56	0.52	29	82.89	2.86	0.95	1.00	0.94	0.00
30	85.47	2.64	0.57	0.99	0.47	0.43	30	84.37	2.01	0.67	0.99	0.54	0.00
31	85.74	2.14	0.46	0.99	0.44	0.41	31	84.64	1.60	0.53	0.98	0.51	0.00
32	86.02	1.81	0.39	0.98	0.42	0.39	32	85.47	1.33	0.44	0.97	0.43	0.00
33	86.29	1.62	0.35	0.98	0.40	0.37	33	85.74	1.15	0.38	0.96	0.41	0.00
34	89.34	0.92	0.20	0.88	0.29	0.28	34	86.02	0.91	0.30	0.95	0.39	0.00
35	92.35	0.65	0.14	0.71	0.23	0.24	35	86.29	0.78	0.26	0.94	0.37	0.00
36	95.64	0.58	0.12	0.51	0.19	0.21	36	89.34	0.42	0.14	0.74	0.27	0.00
37	98.92	0.39	0.08	0.36	0.17	0.19	37	92.35	0.32	0.11	0.52	0.21	0.00
38	102.54	0.33	0.07	0.25	0.15	0.18	38	95.64	0.34	0.11	0.31	0.18	0.00
39	107.10	0.32	0.07	0.14	0.13	0.16	39	98.92	0.31	0.10	0.19	0.16	0.00
40	108.74	0.29	0.06	0.10	0.12	0.16	40	102.54	0.28	0.09	0.11	0.14	0.00
41	111.90	0.28	0.06	0.07	0.11	0.15	41	107.10	0.25	0.08	0.05	0.12	0.00
42	116.84	0.27	0.06	0.03	0.10	0.13	42	108.74	0.24	0.08	0.03	0.11	0.00
43	0.00	0.00	0.00	0.00	0.00	0.00	43	111.90	0.23	0.08	0.02	0.10	0.00
44	0.00	0.00	0.00	0.00	0.00	0.00	44	116.84	0.20	0.07	0.01	0.09	0.00



Table C.2 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Nickel (Ni²⁺) transport through lateritic soil under pH 5 with initial concentration 4.85 mmol/L for Pb²⁺ and 3.17mmol/L for Ni²⁺ (Duplicated)

Sample #	Observed data Pb				Fitted			Observed data Ni				Fitted		
	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	
1	1.61	0.00	0.00	0.00	0.00	0.00	1	1.61	0.00	0.00	0.00	0.00	0.00	
2	3.93	0.00	0.00	0.02	0.00	0.00	2	3.93	0.00	0.00	0.11	0.00	0.00	
3	6.63	0.00	0.00	0.11	0.00	0.00	3	6.63	0.11	0.04	0.36	0.00	0.00	
4	9.32	0.27	0.05	0.27	0.00	0.00	4	9.32	0.64	0.20	0.60	0.00	0.02	
5	10.67	0.52	0.11	0.35	0.00	0.00	5	10.67	0.94	0.30	0.69	0.00	0.31	
6	12.02	1.02	0.21	0.44	0.00	0.00	6	12.02	1.52	0.48	0.77	0.00	0.51	
7	14.74	1.47	0.30	0.59	0.00	0.28	7	14.74	1.73	0.54	0.87	0.00	0.63	
8	17.33	2.04	0.42	0.70	0.00	0.48	8	17.33	2.13	0.67	0.93	0.00	0.71	
9	20.02	2.50	0.52	0.79	0.52	0.55	9	20.02	2.35	0.74	0.96	0.74	0.77	
10	22.53	2.78	0.57	0.86	1.00	0.61	10	22.53	2.53	0.80	0.98	1.00	0.82	
11	24.99	3.03	0.63	0.90	1.00	0.66	11	24.99	2.66	0.84	0.99	1.00	0.86	
12	29.17	3.37	0.70	0.94	1.00	0.72	12	29.17	2.94	0.93	1.00	1.00	0.90	
13	31.94	3.78	0.78	0.96	1.00	0.75	13	31.94	3.05	0.96	1.00	1.00	0.92	
14	34.72	3.83	0.79	0.97	1.00	0.78	14	34.72	3.02	0.95	1.00	1.00	0.94	
15	37.50	3.88	0.80	0.98	1.00	0.81	15	37.50	3.08	0.97	1.00	1.00	0.95	
16	40.31	4.00	0.83	0.99	1.00	0.84	16	40.31	3.08	0.97	1.00	1.00	0.96	
17	43.13	4.04	0.83	0.99	1.00	0.86	17	43.13	3.15	1.00	1.00	1.00	0.97	
18	45.95	4.12	0.85	0.99	1.00	0.87	18	45.95	3.13	0.99	1.00	1.00	0.98	
19	48.80	4.18	0.86	1.00	1.00	0.89	19	48.80	3.24	1.02	1.00	1.00	0.98	
20	51.66	4.30	0.89	1.00	1.00	0.90	20	51.66	3.26	1.03	1.00	1.00	0.99	
21	55.86	4.57	0.94	1.00	1.00	0.92	21	55.86	3.33	1.05	1.00	1.00	0.99	
22	58.70	4.63	0.95	1.00	1.00	0.93	22	58.70	3.19	1.01	1.00	1.00	0.99	
23	61.49	4.72	0.97	1.00	1.00	0.94	23	61.49	3.20	1.01	1.00	1.00	0.99	
24	64.29	4.83	1.00	1.00	1.00	0.95	24	64.29	3.30	1.04	1.00	1.00	1.00	

Table C.2 Observed, predicted and simulated breakthrough curves of Lead (Pb2+) and Nickel (Ni2+) transport through lateritic soil under pH 5 with initial concentration 4.85 mmol/L for Pb2+ and 3.17mmol/L for Ni2+ (Duplicated) (Continued)

25	67.07	4.82	0.99	1.00	1.00	0.95	25	67.07	3.23	1.02	1.00	1.00	1.00
26	69.89	4.75	0.98	1.00	1.00	0.96	26	69.89	3.26	1.03	1.00	1.00	1.00
27	72.69	4.81	0.99	1.00	1.00	0.97	27	72.69	3.22	1.02	1.00	1.00	1.00
28	75.53	4.73	0.98	1.00	1.00	0.97	28	75.53	3.21	1.01	1.00	1.00	1.00
29	78.40	4.84	1.00	1.00	1.00	0.97	29	78.40	3.18	1.00	1.00	1.00	1.00
30	82.74	4.88	1.01	1.00	1.00	0.98	30	82.74	3.23	1.02	1.00	1.00	1.00
31	85.61	4.89	1.01	1.00	1.00	0.98	31	85.61	3.12	0.99	1.00	1.00	1.00
32	88.50	4.98	1.03	1.00	1.00	0.98	32	88.50	3.12	0.99	1.00	1.00	1.00
33	91.44	4.86	1.00	1.00	1.00	0.99	33	91.44	3.09	0.97	1.00	1.00	1.00
34	92.94	4.98	1.03	1.00	1.00	0.99	34	92.94	3.16	1.00	1.00	1.00	1.00
35	93.52	4.98	1.03	1.00	0.92	0.89	35	93.52	3.26	1.03	1.00	0.87	0.86
36	94.12	4.36	0.90	1.00	0.72	0.67	36	94.12	2.78	0.88	1.00	0.64	0.62
37	94.73	2.99	0.62	1.00	0.58	0.54	37	94.73	1.78	0.56	0.99	0.50	0.48
38	95.34	2.33	0.48	1.00	0.49	0.45	38	95.34	1.23	0.39	0.98	0.41	0.41
39	95.94	1.80	0.37	1.00	0.43	0.40	39	95.94	0.98	0.31	0.97	0.36	0.36
40	96.54	1.43	0.29	0.99	0.38	0.36	40	96.54	0.81	0.26	0.94	0.32	0.32
41	97.14	1.27	0.26	0.98	0.35	0.33	41	97.14	0.64	0.20	0.91	0.29	0.30
42	98.95	1.00	0.21	0.94	0.28	0.28	42	98.95	0.39	0.12	0.77	0.23	0.25
43	100.45	0.90	0.19	0.89	0.25	0.25	43	100.45	0.34	0.11	0.64	0.21	0.23
44	103.43	0.70	0.14	0.73	0.20	0.22	44	103.43	0.25	0.08	0.40	0.17	0.20
45	106.41	0.65	0.13	0.57	0.17	0.20	45	110.72	0.26	0.08	0.10	0.12	0.15
46	110.72	0.56	0.12	0.35	0.14	0.18	46	113.47	0.21	0.06	0.05	0.11	0.14
47	113.47	0.54	0.11	0.25	0.13	0.17	47	116.21	0.18	0.06	0.03	0.10	0.13
48	116.21	0.52	0.11	0.17	0.12	0.16	48	118.90	0.15	0.05	0.02	0.10	0.13
49	118.90	0.42	0.09	0.12	0.11	0.15	49	121.60	0.22	0.07	0.01	0.09	0.12
50	121.60	0.46	0.10	0.08	0.11	0.15	50	124.32	0.21	0.07	0.00	0.08	0.11
51	124.32	0.44	0.09	0.06	0.10	0.14	51	127.01	0.19	0.06	0.00	0.08	0.11
52	127.01	0.40	0.08	0.04	0.09	0.14	52	129.75	0.19	0.06	0.00	0.08	0.10
53	129.75	0.40	0.08	0.03	0.09	0.13	53	132.42	0.18	0.06	0.00	0.07	0.10

Table C.2 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Nickel (Ni²⁺) transport through lateritic soil under pH 5 with initial concentration 4.85 mmol/L for Pb²⁺ and 3.17mmol/L for Ni²⁺ (Duplicated) (Continued)

54	132.42	0.38	0.08	0.02	0.08	0.12	54	136.15	0.22	0.07	0.00	0.07	0.09
55	136.15	0.44	0.09	0.01	0.08	0.12	55	0.00	0.00	0.00	0.00	0.00	0.00

Table C.3 Observed, predicted and simulated breakthrough curves of Lead (Pb2+) and Nickel (Ni2+) transport through lateritic soil under pH 5 with initial concentration 4.93 mmol/L for Pb2+ and 5.13 mmol/L for Ni2+

Sample #	Observed data Pb						Fitted			Observed data Ni						Fitted				
	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0
1	1.07	0.00	0.00	0.00	0.00	0.00	1	1.07	0.00	0.00	0.00	0.00	0.00	1	1.07	0.00	0.00	0.00	0.00	0.00
2	2.96	0.00	0.00	0.00	0.00	0.00	2	2.96	0.00	0.00	0.00	0.00	0.00	2	2.96	0.00	0.00	0.03	0.00	0.00
3	4.94	0.00	0.00	0.03	0.00	0.00	3	4.94	0.00	0.00	0.00	0.00	0.00	3	4.94	0.00	0.00	0.14	0.00	0.00
4	7.00	0.00	0.00	0.11	0.00	0.00	4	7.00	8.07	0.00	0.00	0.00	0.00	4	7.00	8.07	0.03	0.34	0.00	0.00
5	9.46	0.00	0.00	0.24	0.00	0.00	5	9.46	54.94	0.00	0.00	0.00	0.00	5	9.46	54.94	0.18	0.55	0.09	0.09
6	10.27	0.00	0.00	0.29	0.00	0.00	6	10.27	73.24	0.00	0.00	0.00	0.00	6	10.27	73.24	0.24	0.61	0.30	0.27
7	12.43	0.31	0.06	0.42	0.00	0.00	7	12.43	132.21	0.00	0.00	0.00	0.00	7	12.43	132.21	0.44	0.74	0.89	0.56
8	15.62	1.11	0.22	0.61	0.00	0.21	8	15.62	215.24	0.00	0.00	0.21	0.00	8	15.62	215.24	0.72	0.87	1.00	0.73
9	18.55	2.12	0.43	0.73	0.00	0.53	9	18.55	263.58	0.00	0.00	0.53	0.00	9	18.55	263.58	0.88	0.93	1.00	0.82
10	21.85	2.99	0.61	0.83	0.62	0.64	10	21.85	280.78	0.61	0.62	0.64	0.00	10	21.85	280.78	0.93	0.97	1.00	0.89
11	24.41	3.49	0.71	0.88	1.00	0.71	11	24.41	286.66	0.71	1.00	0.71	0.00	11	24.41	286.66	0.95	0.98	1.00	0.93
12	28.32	4.19	0.85	0.93	1.00	0.78	12	28.32	296.03	0.85	1.00	0.78	0.00	12	28.32	296.03	0.98	0.99	1.00	0.96
13	30.45	4.18	0.85	0.95	1.00	0.82	13	30.45	290.58	0.85	1.00	0.82	0.00	13	30.45	290.58	0.97	1.00	1.00	0.97
14	34.08	4.36	0.88	0.97	1.00	0.86	14	34.08	290.30	0.88	1.00	0.86	0.00	14	34.08	290.30	0.96	1.00	1.00	0.98
15	36.79	4.38	0.89	0.98	1.00	0.88	15	36.79	289.04	0.89	1.00	0.88	0.00	15	36.79	289.04	0.96	1.00	1.00	0.99
16	40.72	4.47	0.91	0.99	1.00	0.91	16	40.72	296.17	0.91	1.00	0.91	0.00	16	40.72	296.17	0.98	1.00	1.00	0.99
17	44.40	4.53	0.92	0.99	1.00	0.93	17	44.40	289.18	0.92	1.00	0.93	0.00	17	44.40	289.18	0.96	1.00	1.00	1.00
18	47.32	4.64	0.94	1.00	1.00	0.95	18	47.32	292.89	0.94	1.00	0.95	0.00	18	47.32	292.89	0.97	1.00	1.00	1.00
19	50.49	4.64	0.94	1.00	1.00	0.95	19	50.49	293.45	0.94	1.00	0.95	0.00	19	50.49	293.45	0.97	1.00	1.00	1.00
20	54.11	4.81	0.98	1.00	1.00	0.97	20	54.11	294.63	0.98	1.00	0.97	0.00	20	54.11	294.63	0.98	1.00	1.00	1.00
21	57.26	4.81	0.98	1.00	1.00	0.97	21	57.26	294.28	0.98	1.00	0.97	0.00	21	57.26	294.28	0.98	1.00	1.00	1.00
22	60.78	4.76	0.97	1.00	1.00	0.98	22	60.78	292.47	0.97	1.00	0.98	0.00	22	60.78	292.47	0.97	1.00	1.00	1.00
23	63.80	4.79	0.97	1.00	1.00	0.98	23	63.80	294.00	0.97	1.00	0.98	0.00	23	63.80	294.00	0.98	1.00	1.00	1.00
24	67.44	4.74	0.96	1.00	1.00	0.99	24	67.44	294.42	0.96	1.00	0.99	0.00	24	67.44	294.42	0.98	1.00	1.00	1.00
25	70.43	4.75	0.96	1.00	1.00	0.99	25	70.43	296.31	0.96	1.00	0.99	0.00	25	70.43	296.31	0.98	1.00	1.00	1.00

Table C.3 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Nickel (Ni²⁺) transport through lateritic soil under pH 5 with initial concentration 4.93 mmol/L for Pb²⁺ and 5.13 mmol/L for Ni²⁺ (Continued)

26	73.65	4.83	0.98	1.00	1.00	0.99	26	73.65	296.45	0.98	1.00	1.00	1.00
27	75.77	4.97	1.01	1.00	1.00	0.99	27	75.77	296.19	0.98	1.00	1.00	1.00
28	80.19	4.28	0.87	1.00	0.99	0.97	28	80.19	301.29	1.00	1.00	1.00	0.98
29	81.63	3.06	0.62	1.00	0.52	0.44	29	81.63	204.32	0.68	0.98	0.75	0.47
30	81.93	2.46	0.50	1.00	0.49	0.41	30	81.93	161.44	0.54	0.97	0.71	0.44
31	82.14	2.14	0.43	0.99	0.46	0.38	31	82.14	134.57	0.45	0.96	0.67	0.42
32	82.60	1.65	0.34	0.99	0.41	0.35	32	82.60	95.09	0.32	0.94	0.61	0.38
33	82.87	1.46	0.30	0.99	0.39	0.33	33	82.87	85.28	0.28	0.92	0.58	0.36
34	83.63	0.64	0.13	0.88	0.27	0.24	34	83.63	40.75	0.14	0.64	0.38	0.27
35	86.66	0.57	0.12	0.71	0.21	0.21	35	86.66	28.86	0.10	0.39	0.28	0.23
36	90.16	0.62	0.13	0.50	0.18	0.19	36	90.16	19.04	0.06	0.20	0.22	0.19
37	92.69	0.53	0.11	0.36	0.16	0.17	37	92.69	15.75	0.05	0.11	0.18	0.17
38	95.64	0.47	0.10	0.24	0.14	0.16	38	95.64	14.05	0.05	0.05	0.15	0.15
39	98.50	0.41	0.08	0.14	0.12	0.14	39	98.50	12.68	0.04	0.02	0.12	0.13
40	100.82	0.40	0.08	0.13	0.11	0.13	40	100.82	12.71	0.04	0.01	0.10	0.13

Table C.4 Observed, predicted and simulated breakthrough curves of Lead (Pb2+) and Nickel (Ni2+) transport through lateritic soil under pH 5 with initial concentration 4.88 mmol/L for Pb2+ and 10.06 mmol/L for Ni2+

Sample #	Observed data Pb				Fitted			Observed data Ni				Fitted		
	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	
1	0.48	0.00	0.00	0.00	0.00	0.00	1	0.48	0.00	0.00	0.00	0.00	0.00	
2	2.17	0.00	0.00	0.01	0.00	0.00	2	2.17	0.00	0.00	0.05	0.00	0.00	
3	4.60	0.00	0.00	0.08	0.00	0.00	3	4.60	0.29	0.03	0.30	0.00	0.01	
4	7.03	0.00	0.00	0.23	0.00	0.00	4	5.82	1.88	0.19	0.45	0.00	0.09	
5	9.45	0.10	0.02	0.41	0.00	0.00	5	7.03	3.45	0.34	0.58	0.23	0.37	
6	11.88	0.96	0.20	0.57	0.00	0.14	6	8.24	4.65	0.46	0.69	0.89	0.57	
7	14.30	1.88	0.39	0.70	0.00	0.53	7	9.45	6.66	0.66	0.77	0.99	0.68	
8	16.74	2.51	0.51	0.80	0.00	0.61	8	10.67	7.22	0.72	0.84	1.00	0.75	
9	19.16	3.15	0.65	0.86	0.63	0.67	9	13.09	8.69	0.86	0.92	1.00	0.85	
10	21.58	3.56	0.73	0.91	1.00	0.72	10	15.53	9.30	0.92	0.96	1.00	0.91	
11	25.23	3.91	0.80	0.95	1.00	0.78	11	16.74	9.43	0.94	0.97	1.00	0.93	
12	27.65	4.20	0.86	0.97	1.00	0.81	12	19.16	9.63	0.96	0.98	1.00	0.96	
13	30.08	4.28	0.88	0.98	1.00	0.84	13	21.58	9.74	0.97	0.99	1.00	0.97	
14	32.50	4.35	0.89	0.99	1.00	0.86	14	25.23	9.97	0.99	1.00	1.00	0.99	
15	37.33	4.56	0.93	0.99	1.00	0.90	15	27.65	9.82	0.98	1.00	1.00	0.99	
16	39.75	4.55	0.93	1.00	1.00	0.92	16	30.08	9.78	0.97	1.00	1.00	1.00	
17	42.17	4.59	0.94	1.00	1.00	0.93	17	32.50	9.89	0.98	1.00	1.00	1.00	
18	44.57	4.67	0.96	1.00	1.00	0.94	18	37.33	9.88	0.98	1.00	1.00	1.00	
19	48.14	4.68	0.96	1.00	1.00	0.95	19	39.75	9.85	0.98	1.00	1.00	1.00	
20	50.51	4.60	0.94	1.00	1.00	0.96	20	42.17	9.91	0.98	1.00	1.00	1.00	
21	52.92	4.70	0.96	1.00	1.00	0.97	21	44.57	10.09	1.00	1.00	1.00	1.00	
22	55.35	4.61	0.94	1.00	1.00	0.97	22	48.14	10.02	1.00	1.00	1.00	1.00	
23	57.77	4.38	0.90	1.00	1.00	0.98	23	50.51	9.91	0.99	1.00	1.00	1.00	
24	58.25	4.74	0.97	1.00	0.93	0.90	24	52.92	9.98	0.99	1.00	1.00	1.00	

Table C.4 Observed, predicted and simulated breakthrough curves of Lead (Pb2+) and Nickel (Ni2+) transport through lateritic soil under pH 5 with initial concentration 4.88 mmol/L for Pb2+ and 10.06 mmol/L for Ni2+ (Continued)

25	58.73	4.70	0.96	1.00	0.74	0.70	25	55.35	9.81	0.98	1.00	1.00	1.00
26	59.22	3.88	0.80	1.00	0.60	0.57	26	57.77	7.87	0.86	1.00	1.00	1.00
27	59.46	3.15	0.65	1.00	0.55	0.52	27	58.25	8.82	0.97	1.00	0.96	0.94
28	59.70	2.47	0.51	1.00	0.51	0.48	28	59.22	8.58	0.94	0.99	0.68	0.63
29	59.95	1.99	0.41	1.00	0.47	0.45	29	59.46	7.92	0.87	0.98	0.62	0.58
30	60.19	1.64	0.34	0.99	0.44	0.42	30	59.70	6.60	0.72	0.97	0.58	0.54
31	62.58	0.84	0.17	0.92	0.29	0.29	31	59.95	4.11	0.45	0.96	0.54	0.51
32	64.98	0.56	0.11	0.77	0.23	0.24	32	60.19	3.38	0.37	0.95	0.50	0.48
33	67.37	0.47	0.10	0.59	0.19	0.21	33	62.58	1.59	0.17	0.70	0.31	0.33
34	70.94	0.37	0.08	0.36	0.16	0.18	34	64.98	1.09	0.12	0.41	0.23	0.27
35	73.31	0.37	0.08	0.25	0.14	0.17	35	67.37	0.90	0.10	0.23	0.18	0.23
36	75.67	0.32	0.06	0.17	0.13	0.16	36	70.94	0.76	0.08	0.08	0.14	0.18
37	77.98	0.32	0.07	0.11	0.12	0.15	37	73.31	0.72	0.08	0.04	0.12	0.16
38	80.33	0.31	0.06	0.08	0.11	0.14	38	75.67	0.68	0.07	0.02	0.11	0.14
39	82.65	0.32	0.06	0.05	0.11	0.14	39	77.98	0.71	0.08	0.01	0.09	0.13
40	84.92	0.25	0.05	0.03	0.10	0.13	40	80.33	0.65	0.07	0.01	0.08	0.12
41	87.09	0.28	0.06	0.02	0.09	0.12	41	82.65	0.59	0.06	0.00	0.07	0.10
42	89.23	0.32	0.07	0.01	0.09	0.12	42	84.92	0.61	0.07	0.00	0.07	0.09
43	0.00	0.00	0.00	0.00	0.00	0.00	43	87.09	0.58	0.06	0.00	0.06	0.09
44	0.00	0.00	0.00	0.00	0.00	0.00	44	89.23	0.53	0.06	0.00	0.05	0.08

Table C.5 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Zinc (Zn²⁺) transport through lateritic soil under pH 5 with initial concentration 5.26 mmol/L for Pb²⁺ and 2.91 mmol/L for Zn²⁺

Sample #	Observed data Pb				Fitted			Observed data Zn				Fitted		
	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	
1	1.77	0.01	0.00	0.00	0.00	0.00	1	1.77	0.02	0.01	0.04	0.00	0.00	
2	4.19	0.02	0.00	0.07	0.00	0.00	2	3.95	0.11	0.04	0.31	0.00	0.00	
3	6.31	0.17	0.03	0.22	0.00	0.00	3	6.09	0.57	0.19	0.62	0.00	0.14	
4	8.53	0.57	0.11	0.41	0.00	0.01	4	8.30	1.05	0.36	0.81	0.38	0.50	
5	10.53	1.22	0.23	0.56	0.00	0.19	5	10.53	1.59	0.55	0.91	1.00	0.63	
6	12.75	1.82	0.35	0.69	0.00	0.44	6	12.75	2.02	0.69	0.96	1.00	0.72	
7	14.94	2.47	0.47	0.79	0.00	0.55	7	14.94	2.31	0.79	0.98	1.00	0.79	
8	17.15	2.97	0.56	0.86	0.58	0.63	8	17.15	2.46	0.84	0.99	1.00	0.84	
9	19.35	3.34	0.63	0.91	1.00	0.70	9	19.35	2.83	0.97	1.00	1.00	0.88	
10	22.49	4.26	0.81	0.95	1.00	0.77	10	22.49	2.83	0.97	1.00	1.00	0.92	
11	24.77	4.51	0.86	0.97	1.00	0.81	11	24.77	2.78	0.96	1.00	1.00	0.94	
12	27.02	4.49	0.85	0.98	1.00	0.85	12	27.02	2.79	0.96	1.00	1.00	0.96	
13	29.29	4.75	0.90	0.99	1.00	0.87	13	29.29	2.80	0.96	1.00	1.00	0.97	
14	31.57	4.96	0.94	0.99	1.00	0.90	14	31.57	2.81	0.97	1.00	1.00	0.97	
15	33.85	5.07	0.96	0.99	1.00	0.92	15	33.85	2.82	0.97	1.00	1.00	0.98	
16	36.15	5.21	0.99	1.00	1.00	0.93	16	36.15	2.83	0.97	1.00	1.00	0.99	
17	38.43	5.19	0.99	1.00	1.00	0.94	17	38.43	2.85	0.98	1.00	1.00	0.99	
18	40.72	5.32	1.01	1.00	1.00	0.95	18	40.72	2.87	0.99	1.00	1.00	0.99	
19	44.14	5.06	0.96	1.00	1.00	0.97	19	44.14	2.85	0.98	1.00	1.00	0.99	
20	46.43	4.91	0.93	1.00	1.00	0.97	20	46.43	3.00	1.03	1.00	1.00	1.00	
21	48.74	5.11	0.97	1.00	1.00	0.98	21	48.74	2.89	0.99	1.00	1.00	1.00	
22	51.05	5.17	0.98	1.00	1.00	0.98	22	51.05	2.93	1.01	1.00	1.00	1.00	
23	53.38	4.99	0.95	1.00	1.00	0.98	23	53.38	2.91	1.00	1.00	1.00	1.00	
24	55.69	5.11	0.97	1.00	1.00	0.99	24	55.69	2.87	0.99	1.00	1.00	1.00	

Table C.5 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Zinc (Zn²⁺) transport through lateritic soil under pH 5 with initial concentration 5.26 mmol/L for Pb²⁺ and 2.91 mmol/L for Zn²⁺ (Continued)

25	58.00	4.88	0.93	1.00	1.00	0.99	25	58.00	2.92	1.00	1.00	1.00	1.00
26	60.31	5.16	0.98	1.00	1.00	0.99	26	60.31	2.81	0.97	1.00	1.00	1.00
27	62.64	4.94	0.94	1.00	1.00	0.99	27	62.64	2.71	0.93	1.00	1.00	1.00
28	64.97	5.27	1.00	1.00	1.00	0.99	28	64.97	2.89	0.99	1.00	1.00	1.00
29	65.20	5.17	0.98	1.00	1.00	0.99	29	65.20	2.88	0.99	1.00	1.00	1.00
30	65.43	5.07	0.96	1.00	0.99	0.98	30	65.43	2.85	0.98	1.00	0.99	0.98
31	65.67	5.01	0.95	1.00	0.93	0.89	31	65.67	2.79	0.96	1.00	0.92	0.90
32	65.90	4.95	0.94	1.00	0.84	0.78	32	65.90	2.81	0.97	1.00	0.82	0.79
33	66.13	4.59	0.87	1.00	0.75	0.68	33	66.13	2.58	0.89	1.00	0.72	0.69
34	66.37	4.04	0.77	1.00	0.68	0.60	34	66.37	2.19	0.75	0.99	0.64	0.61
35	66.60	2.98	0.57	1.00	0.62	0.54	35	66.60	1.72	0.59	0.99	0.58	0.55
36	66.83	2.32	0.44	1.00	0.57	0.49	36	66.83	1.31	0.45	0.98	0.53	0.50
37	67.30	1.86	0.35	1.00	0.49	0.43	37	67.30	0.99	0.34	0.95	0.45	0.44
38	68.00	1.15	0.22	0.99	0.41	0.37	38	68.00	0.77	0.27	0.88	0.38	0.38
39	68.47	1.09	0.21	0.98	0.38	0.34	39	68.47	0.66	0.23	0.82	0.34	0.35
40	70.79	0.98	0.19	0.86	0.27	0.27	40	70.79	0.41	0.14	0.49	0.24	0.27
41	73.09	0.83	0.16	0.69	0.22	0.24	41	73.09	0.34	0.12	0.25	0.19	0.24
42	75.39	0.71	0.14	0.51	0.19	0.22	42	75.39	0.29	0.10	0.12	0.16	0.21
43	77.51	0.67	0.13	0.36	0.17	0.20	43	77.51	0.27	0.09	0.06	0.14	0.19
44	79.50	0.59	0.11	0.24	0.15	0.19	44	79.50	0.24	0.08	0.03	0.12	0.17
45	81.39	0.53	0.10	0.17	0.14	0.17	45	81.39	0.23	0.08	0.01	0.11	0.16
46	83.13	0.48	0.09	0.11	0.13	0.16	46	83.13	0.20	0.07	0.01	0.10	0.15
47	85.40	0.41	0.08	0.06	0.12	0.15	47	85.40	0.19	0.06	0.00	0.09	0.13
48	90.53	0.37	0.07	0.04	0.11	0.14	48	90.53	0.16	0.06	0.00	0.08	0.12

Table C.6 Observed, predicted and simulated breakthrough curves of Lead (Pb2+) and Zinc (Zn2+) transport through lateritic soil under pH 5 with initial concentration 5.22 mmol/L for Pb2+ and 5.36 mmol/L for Zn2+

Sample #	Observed data Pb			Fitted			Observed data Zn			Fitted			
	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0
1	0.23	0.00	0.00	0.00	0.00	0.00	1	0.23	0.00	0.00	0.00	0.00	0.00
2	1.79	0.00	0.00	0.00	0.00	0.00	2	1.79	0.12	0.02	0.11	0.00	0.00
3	4.00	0.00	0.00	0.06	0.00	0.00	3	4.00	0.20	0.04	0.57	0.00	0.00
4	6.32	0.04	0.01	0.21	0.00	0.00	4	6.32	1.08	0.20	0.84	0.00	0.17
5	8.56	0.48	0.09	0.39	0.00	0.00	5	8.56	2.31	0.43	0.95	0.00	0.61
6	10.93	1.01	0.19	0.56	0.00	0.14	6	10.93	3.75	0.70	0.98	0.73	0.74
7	13.28	1.93	0.37	0.69	0.00	0.48	7	13.28	4.53	0.84	0.99	1.00	0.82
8	15.52	2.63	0.50	0.79	0.52	0.59	8	15.52	4.99	0.93	1.00	1.00	0.88
9	17.94	3.43	0.66	0.86	1.00	0.68	9	17.94	5.17	0.96	1.00	1.00	0.92
10	20.28	3.86	0.74	0.91	1.00	0.75	10	20.28	5.29	0.99	1.00	1.00	0.94
11	23.83	4.31	0.83	0.95	1.00	0.82	11	23.83	5.27	0.98	1.00	1.00	0.97
12	26.12	4.57	0.88	0.97	1.00	0.86	12	26.12	5.25	0.98	1.00	1.00	0.98
13	28.39	4.64	0.89	0.98	1.00	0.89	13	28.39	5.23	0.97	1.00	1.00	0.99
14	30.65	4.91	0.94	0.99	1.00	0.91	14	30.65	5.30	0.99	1.00	1.00	0.99
15	32.89	5.12	0.98	0.99	1.00	0.93	15	32.89	5.30	0.99	1.00	1.00	0.99
16	35.19	5.00	0.96	0.99	1.00	0.94	16	35.19	5.21	0.97	1.00	1.00	1.00
17	37.44	5.06	0.97	1.00	1.00	0.96	17	37.44	5.26	0.98	1.00	1.00	1.00
18	39.80	5.09	0.98	1.00	1.00	0.97	18	39.80	5.25	0.98	1.00	1.00	1.00
19	42.08	5.13	0.98	1.00	1.00	0.97	19	42.08	5.33	0.99	1.00	1.00	1.00
20	45.52	5.22	1.00	1.00	1.00	0.98	20	45.52	5.38	1.00	1.00	1.00	1.00
21	47.67	5.26	1.01	1.00	1.00	0.98	21	47.67	5.34	1.00	1.00	1.00	1.00
22	49.93	4.96	0.95	1.00	1.00	0.99	22	49.93	5.21	0.97	1.00	1.00	1.00
23	52.21	5.33	1.02	1.00	1.00	0.99	23	52.21	5.63	1.05	1.00	1.00	1.00
24	54.40	5.34	1.02	1.00	1.00	0.99	24	54.40	5.53	1.03	1.00	1.00	1.00

Table C.6 Observed, predicted and simulated breakthrough curves of Lead (Pb2+) and Zinc (Zn2+) transport through lateritic soil under pH 5 with initial concentration 5.22 mmol/L for Pb2+ and 5.36 mmol/L for Zn2+ (Continued)

25	55.09	5.04	0.97	1.00	1.00	0.99	0.99	25	55.09	5.43	1.01	1.00	1.00	1.00	1.00
26	55.31	6.22	1.19	1.00	0.99	0.98	0.98	26	55.31	6.23	1.16	1.00	1.00	0.99	0.99
27	55.55	5.04	0.97	1.00	0.89	0.90	0.90	27	55.55	5.13	0.96	1.00	1.00	0.92	0.90
28	55.78	4.92	0.94	1.00	0.76	0.78	0.78	28	55.78	5.08	0.95	0.99	0.99	0.80	0.77
29	56.02	4.49	0.86	1.00	0.65	0.67	0.67	29	56.02	4.59	0.86	0.99	0.99	0.69	0.66
30	56.25	3.66	0.70	1.00	0.57	0.59	0.59	30	56.25	3.66	0.68	0.97	0.97	0.61	0.58
31	56.48	3.01	0.58	1.00	0.50	0.53	0.53	31	56.48	2.88	0.54	0.95	0.95	0.54	0.52
32	56.71	1.84	0.35	1.00	0.46	0.48	0.48	32	56.71	2.17	0.40	0.92	0.92	0.49	0.47
33	57.40	1.75	0.33	0.99	0.36	0.39	0.39	33	57.40	1.29	0.24	0.80	0.80	0.39	0.38
34	57.86	1.54	0.30	0.99	0.32	0.36	0.36	34	57.86	1.12	0.21	0.71	0.71	0.35	0.35
35	59.06	1.21	0.23	0.95	0.26	0.30	0.30	35	59.06	0.91	0.17	0.48	0.48	0.28	0.29
36	61.38	0.87	0.17	0.81	0.19	0.25	0.25	36	61.38	0.61	0.11	0.18	0.18	0.21	0.23
37	63.67	0.69	0.13	0.62	0.16	0.22	0.22	37	63.67	0.46	0.09	0.06	0.06	0.17	0.20
38	67.00	0.56	0.11	0.38	0.13	0.19	0.19	38	67.00	0.40	0.07	0.01	0.01	0.14	0.16
39	69.16	0.52	0.10	0.26	0.12	0.18	0.18	39	69.16	0.37	0.07	0.00	0.00	0.12	0.15
40	71.32	0.47	0.09	0.18	0.11	0.16	0.16	40	71.32	0.35	0.06	0.00	0.00	0.11	0.13
41	73.51	0.44	0.08	0.12	0.10	0.15	0.15	41	73.51	0.32	0.06	0.00	0.00	0.10	0.12
42	75.56	0.40	0.08	0.08	0.09	0.14	0.14	42	75.56	0.30	0.06	0.00	0.00	0.09	0.11
43	79.91	0.47	0.09	0.03	0.08	0.13	0.13	43	79.91	0.31	0.06	0.00	0.00	0.08	0.10
44	81.96	0.34	0.06	0.02	0.08	0.12	0.12	44	81.96	0.28	0.05	0.00	0.00	0.08	0.09
45	84.07	0.35	0.07	0.01	0.07	0.12	0.12	45	84.07	0.28	0.05	0.00	0.00	0.07	0.08

Table C.7 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Zinc (Zn²⁺) transport through lateritic soil under pH 5 with initial concentration 5.41 mmol/L for Pb²⁺ and 9.67 mmol/L for Zn²⁺

Sample #	Observed data Pb				Fitted				Observed data Zn				Fitted			
	Pore volume	Conc.(mM)	C/C0	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	
1	0.26	0.00	0.00	0.00	0.00	0.00	0.00	1	0.26	0.09	0.01	0.01	0.01	0.00	0.00	
2	2.47	0.00	0.00	0.04	0.00	0.00	0.00	2	1.36	0.12	0.01	0.01	0.07	0.00	0.00	
3	5.96	0.16	0.03	0.28	0.00	0.00	0.00	3	2.47	0.20	0.02	0.02	0.22	0.00	0.01	
4	7.79	0.68	0.13	0.43	0.00	0.00	0.00	4	5.96	3.87	0.40	0.40	0.69	0.06	0.44	
5	11.57	1.44	0.27	0.68	0.00	0.00	0.24	5	7.79	5.68	0.59	0.59	0.82	0.64	0.63	
6	13.62	2.80	0.52	0.77	0.00	0.00	0.58	6	8.95	6.14	0.63	0.63	0.88	0.89	0.71	
7	16.35	3.30	0.61	0.86	0.61	0.61	0.70	7	11.57	6.85	0.71	0.71	0.95	0.99	0.81	
8	18.16	3.87	0.72	0.92	1.00	1.00	0.77	8	13.62	8.12	0.84	0.84	0.97	1.00	0.86	
9	20.53	4.60	0.85	0.95	1.00	1.00	0.82	9	16.35	8.43	0.87	0.87	0.99	1.00	0.91	
10	22.88	4.84	0.90	0.97	1.00	1.00	0.86	10	18.16	9.35	0.97	0.97	1.00	1.00	0.94	
11	25.49	4.95	0.92	0.98	1.00	1.00	0.89	11	20.53	10.31	1.07	1.07	1.00	1.00	0.96	
12	28.00	4.99	0.92	0.99	1.00	1.00	0.92	12	22.88	9.99	1.03	1.03	1.00	1.00	0.98	
13	30.51	5.25	0.97	0.99	1.00	1.00	0.94	13	25.49	10.18	1.05	1.05	1.00	1.00	0.99	
14	33.06	5.02	0.93	1.00	1.00	1.00	0.95	14	28.00	10.10	1.04	1.04	1.00	1.00	0.99	
15	35.62	5.31	0.98	1.00	1.00	1.00	0.96	15	30.51	10.04	1.04	1.04	1.00	1.00	0.99	
16	38.27	5.37	0.99	1.00	1.00	1.00	0.97	16	33.06	9.46	0.98	0.98	1.00	1.00	0.99	
17	41.21	5.66	1.05	1.00	1.00	1.00	0.98	17	35.62	9.57	0.99	0.99	1.00	1.00	1.00	
18	41.98	5.39	1.00	1.00	1.00	1.00	0.98	18	38.27	9.82	1.02	1.02	1.00	1.00	1.00	
19	42.17	5.19	0.96	1.00	1.00	1.00	0.98	19	41.21	9.32	0.96	0.96	1.00	1.00	1.00	
20	42.37	5.13	0.95	1.00	0.98	0.98	0.96	20	41.98	9.47	0.98	0.98	1.00	1.00	1.00	
21	42.57	5.08	0.94	1.00	0.92	0.92	0.90	21	42.17	8.85	0.92	0.92	1.00	1.00	1.00	
22	42.78	4.92	0.91	1.00	0.83	0.81	0.81	22	42.37	9.52	0.98	0.98	1.00	1.00	0.98	
23	42.97	4.94	0.91	1.00	0.75	0.73	0.73	23	42.57	9.25	0.96	0.96	1.00	0.97	0.94	
24	43.18	4.49	0.83	1.00	0.68	0.66	0.66	24	42.78	9.66	1.00	1.00	0.99	0.92	0.88	

Table C.7 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Zinc (Zn²⁺) transport through lateritic soil under pH 5 with initial concentration 5.41 mmol/L for Pb²⁺ and 9.67 mmol/L for Zn²⁺ (Continued)

25	43.58	3.21	0.59	0.99	0.57	0.56	25	42.97	9.47	0.98	0.98	0.86	0.82
26	43.98	1.95	0.36	0.98	0.50	0.49	26	43.18	9.05	0.94	0.97	0.81	0.76
27	45.16	1.35	0.25	0.91	0.37	0.37	27	43.58	6.35	0.66	0.93	0.70	0.66
28	47.05	0.86	0.16	0.69	0.27	0.28	28	43.98	4.45	0.46	0.86	0.62	0.58
29	48.89	0.64	0.12	0.47	0.22	0.24	29	45.16	1.85	0.19	0.61	0.45	0.44
30	50.71	0.49	0.09	0.30	0.19	0.21	30	47.05	1.08	0.11	0.28	0.31	0.32
31	52.66	0.38	0.07	0.18	0.16	0.19	31	48.89	0.86	0.09	0.11	0.23	0.26
32	54.64	0.30	0.06	0.11	0.15	0.17	32	50.71	0.70	0.07	0.04	0.18	0.21
33	57.01	0.28	0.05	0.06	0.13	0.16	33	52.66	0.61	0.06	0.02	0.15	0.18
34	0.00	0.00	0.00	0.00	0.00	0.00	34	54.64	0.56	0.06	0.01	0.12	0.15
35	0.00	0.00	0.00	0.00	0.00	0.00	35	57.01	0.48	0.05	0.00	0.10	0.12
36	0.00	0.00	0.00	0.00	0.00	0.00	36	59.55	0.46	0.05	0.00	0.08	0.10

Table C.8 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Zinc (Zn²⁺) transport through lateritic soil under pH 5 with initial concentration 5.41 mmol/L for Pb²⁺ and 9.67 mmol/L for Zn²⁺ (Duplicated)

Sample #	Observed data Pb				Fitted				Observed data Zn				Fitted			
	Pore volume	Conc.(mM)	C/C0	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	
1	0.26	0.00	0.00	0.00	0.00	0.00	0.00	1	0.26	0.02	0.00	0.00	0.00	0.00	0.00	
2	1.30	0.00	0.00	0.00	0.00	0.00	0.00	2	1.30	0.02	0.00	0.00	0.08	0.00	0.00	
3	2.62	0.00	0.00	0.00	0.04	0.00	0.00	3	2.62	0.24	0.02	0.40	0.00	0.00	0.00	
4	5.25	0.00	0.00	0.00	0.27	0.00	0.00	4	5.25	1.46	0.15	0.85	0.00	0.00	0.12	
5	7.81	0.29	0.06	0.54	0.54	0.00	0.00	5	7.81	4.20	0.44	0.97	0.44	0.44	0.54	
6	10.34	0.88	0.20	0.74	0.74	0.00	0.12	6	10.34	6.09	0.64	0.99	1.00	1.00	0.68	
7	12.72	1.48	0.33	0.86	0.86	0.00	0.44	7	12.72	7.46	0.79	1.00	1.00	1.00	0.78	
8	15.05	2.05	0.47	0.93	0.93	0.00	0.56	8	15.05	7.63	0.80	1.00	1.00	1.00	0.84	
9	17.28	2.50	0.58	0.96	0.96	0.57	0.63	9	17.28	9.06	0.96	1.00	1.00	1.00	0.89	
10	19.49	2.86	0.66	0.98	0.98	1.00	0.70	10	19.49	9.31	0.98	1.00	1.00	1.00	0.92	
11	21.64	3.15	0.73	0.99	0.99	1.00	0.75	11	21.64	8.67	0.91	1.00	1.00	1.00	0.94	
12	24.91	3.54	0.82	1.00	1.00	1.00	0.82	12	24.91	9.94	1.05	1.00	1.00	1.00	0.97	
13	27.09	3.72	0.86	1.00	1.00	1.00	0.85	13	27.09	9.77	1.03	1.00	1.00	1.00	0.98	
14	29.24	3.88	0.90	1.00	1.00	1.00	0.88	14	29.24	10.07	1.06	1.00	1.00	1.00	0.98	
15	31.40	4.00	0.92	1.00	1.00	1.00	0.90	15	31.40	9.56	1.01	1.00	1.00	1.00	0.99	
16	35.36	4.31	1.00	1.00	1.00	1.00	0.92	16	35.36	9.15	0.97	1.00	1.00	1.00	0.99	
17	37.68	4.26	0.98	1.00	1.00	1.00	0.93	17	37.68	9.51	1.00	1.00	1.00	1.00	0.99	
18	40.01	4.36	1.01	1.00	1.00	1.00	0.94	18	40.01	9.36	0.99	1.00	1.00	1.00	1.00	
19	42.45	4.56	1.05	1.00	1.00	1.00	0.95	19	42.45	9.33	0.98	1.00	1.00	1.00	1.00	
20	44.72	4.49	1.04	1.00	1.00	1.00	0.96	20	44.72	9.35	0.99	1.00	1.00	1.00	1.00	
21	46.31	4.40	1.02	0.99	0.99	0.79	0.72	21	46.31	9.21	0.97	0.95	0.75	0.72	0.72	
22	46.76	3.36	0.77	0.98	0.98	0.66	0.58	22	46.76	6.95	0.73	0.85	0.60	0.60	0.58	
23	47.21	2.19	0.51	0.96	0.96	0.56	0.49	23	47.21	4.38	0.46	0.71	0.50	0.50	0.48	
24	47.68	1.57	0.36	0.93	0.93	0.49	0.43	24	47.68	3.12	0.33	0.57	0.43	0.43	0.42	

Table C.8 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Zinc (Zn²⁺) transport through lateritic soil under pH 5 with initial concentration 5.41 mmol/L for Pb²⁺ and 9.67 mmol/L for Zn²⁺ (Duplicated) (Continued)

25	48.13	1.23	0.28	0.89	0.44	0.39	25	48.13	2.35	0.25	0.45	0.38	0.38
26	48.57	1.05	0.24	0.85	0.40	0.36	26	48.57	1.96	0.21	0.34	0.34	0.35
27	49.02	0.89	0.21	0.80	0.37	0.34	27	49.02	1.79	0.19	0.26	0.31	0.33
28	50.42	0.66	0.15	0.63	0.30	0.29	28	50.42	1.44	0.15	0.10	0.25	0.28
29	54.19	0.30	0.07	0.27	0.22	0.23	29	54.19	0.92	0.10	0.01	0.16	0.21
30	62.58	0.11	0.03	0.03	0.14	0.17	30	62.58	0.62	0.07	0.00	0.10	0.14
31	68.29	0.08	0.02	0.01	0.12	0.15	31	68.29	0.54	0.06	0.00	0.07	0.11
32	73.77	0.06	0.01	0.00	0.10	0.13	32	73.77	0.52	0.05	0.00	0.06	0.09
33	79.23	0.05	0.01	0.00	0.09	0.12	33	79.23	0.48	0.05	0.00	0.05	0.07

Table C.9 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Manganese (Mn²⁺) transport through lateritic soil under pH 5 with initial concentration 5.31 mmol/L for Pb²⁺ and 2.73 mmol/L for Mn²⁺

Sample #	Observed data Pb			Fitted			Observed data Mn			Fitted			
	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0
1	0.85	0.00	0.00	0.00	0.00	0.00	1	0.85	0.00	0.00	0.00	0.00	0.00
2	2.27	0.00	0.00	0.00	0.00	0.00	2	2.27	0.01	0.00	0.04	0.00	0.00
3	5.11	0.00	0.00	0.06	0.00	0.00	3	5.11	0.16	0.06	0.34	0.00	0.00
4	7.93	0.33	0.06	0.20	0.00	0.00	4	7.93	0.84	0.31	0.65	0.00	0.29
5	10.79	0.86	0.16	0.39	0.00	0.00	5	10.79	1.26	0.46	0.84	0.48	0.59
6	13.66	1.30	0.24	0.55	0.00	0.21	6	13.66	1.62	0.59	0.93	1.00	0.67
7	16.51	2.02	0.38	0.69	0.00	0.47	7	16.51	1.89	0.69	0.97	1.00	0.73
8	19.36	2.56	0.48	0.79	0.00	0.58	8	19.36	2.18	0.80	0.98	1.00	0.78
9	22.22	3.04	0.57	0.86	0.58	0.66	9	22.22	2.34	0.86	0.99	1.00	0.83
10	25.08	3.49	0.66	0.90	1.00	0.72	10	25.08	2.34	0.86	1.00	1.00	0.86
11	29.36	4.52	0.85	0.95	1.00	0.80	11	27.94	2.44	0.89	1.00	1.00	0.89
12	32.21	4.84	0.91	0.97	1.00	0.84	12	29.36	2.47	0.91	1.00	1.00	0.90
13	35.06	4.86	0.91	0.98	1.00	0.87	13	32.21	2.61	0.96	1.00	1.00	0.92
14	37.68	4.95	0.93	0.98	1.00	0.89	14	35.06	2.61	0.96	1.00	1.00	0.93
15	40.51	4.87	0.92	0.99	1.00	0.91	15	37.68	2.62	0.96	1.00	1.00	0.95
16	43.34	5.18	0.98	0.99	1.00	0.93	16	40.51	2.60	0.95	1.00	1.00	0.96
17	46.10	5.14	0.97	1.00	1.00	0.94	17	43.34	2.62	0.96	1.00	1.00	0.96
18	48.87	5.26	0.99	1.00	1.00	0.95	18	46.10	2.62	0.96	1.00	1.00	0.97
19	51.68	5.30	1.00	1.00	1.00	0.96	19	48.87	2.61	0.96	1.00	1.00	0.98
20	54.50	5.35	1.01	1.00	1.00	0.97	20	51.68	2.65	0.97	1.00	1.00	0.98
21	58.66	4.60	0.87	1.00	1.00	0.98	21	55.89	2.63	0.96	1.00	1.00	0.99
22	61.43	5.33	1.00	1.00	1.00	0.98	22	58.66	2.66	0.98	1.00	1.00	0.99
23	64.18	5.24	0.99	1.00	1.00	0.99	23	61.43	2.77	1.01	1.00	1.00	0.99
24	66.95	5.08	0.96	1.00	1.00	0.99	24	64.18	2.76	1.01	1.00	1.00	0.99

Table C.9 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Manganese (Mn²⁺) transport through lateritic soil under pH 5 with initial concentration 5.31 mmol/L for Pb²⁺ and 2.73 mmol/L for Mn²⁺ (Continued)

25	69.77	5.23	0.98	1.00	1.00	0.99	25	66.95	2.66	0.98	1.00	1.00	0.99
26	72.59	5.16	0.97	1.00	1.00	0.99	26	69.77	2.74	1.01	1.00	1.00	1.00
27	75.34	5.09	0.96	1.00	1.00	0.99	27	72.59	2.66	0.97	1.00	1.00	1.00
28	78.15	5.11	0.96	1.00	1.00	0.99	28	75.34	2.68	0.98	1.00	1.00	1.00
29	81.56	5.11	0.96	1.00	1.00	1.00	29	78.15	2.66	0.97	1.00	1.00	1.00
30	81.84	5.06	0.95	1.00	0.99	0.98	30	81.56	2.68	0.98	1.00	1.00	1.00
31	82.13	5.00	0.94	1.00	0.90	0.88	31	81.84	2.63	0.96	1.00	0.98	0.97
32	82.41	4.93	0.93	1.00	0.78	0.76	32	82.13	2.57	0.94	1.00	0.91	0.87
33	82.69	4.43	0.83	1.00	0.68	0.66	33	82.41	2.55	0.93	1.00	0.80	0.75
34	82.96	3.85	0.72	1.00	0.61	0.58	34	82.69	2.34	0.86	1.00	0.70	0.66
35	83.24	3.15	0.59	1.00	0.55	0.52	35	82.96	2.02	0.74	0.99	0.63	0.58
36	83.51	2.10	0.40	1.00	0.50	0.48	36	83.24	1.59	0.58	0.99	0.56	0.53
37	83.79	1.64	0.31	1.00	0.46	0.44	37	83.51	1.26	0.46	0.97	0.51	0.48
38	85.22	1.16	0.22	0.98	0.34	0.34	38	83.79	0.98	0.36	0.96	0.47	0.45
39	88.08	0.71	0.13	0.88	0.24	0.27	39	85.22	0.54	0.20	0.83	0.35	0.34
40	90.93	0.69	0.13	0.71	0.20	0.23	40	88.08	0.33	0.12	0.48	0.24	0.26
41	93.78	0.57	0.11	0.53	0.17	0.21	41	90.93	0.28	0.10	0.24	0.19	0.22
42	96.63	0.57	0.11	0.37	0.15	0.19	42	93.78	0.23	0.08	0.11	0.16	0.20
43	99.46	0.49	0.09	0.26	0.13	0.18	43	96.63	0.22	0.08	0.05	0.14	0.18
44	102.29	0.42	0.08	0.18	0.12	0.17	44	99.46	0.19	0.07	0.02	0.12	0.16
45	105.11	0.42	0.08	0.12	0.11	0.16	45	102.29	0.17	0.06	0.01	0.11	0.15
46	109.32	0.39	0.07	0.06	0.10	0.14	46	105.11	0.16	0.06	0.00	0.10	0.14
47	112.11	0.39	0.07	0.04	0.10	0.14	47	109.32	0.14	0.05	0.00	0.09	0.13
48	116.28	0.20	0.04	0.02	0.09	0.13	48	112.11	0.13	0.05	0.00	0.08	0.12
							49	114.89	0.07	0.03	0.00	0.07	0.11

Table C.10 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Manganese (Mn²⁺) transport through lateritic soil under pH 5 with initial concentration 5.14 mmol/L for Pb²⁺ and 4.47 mmol/L for Mn²⁺

Sample #	Observed data Pb				Fitted				Observed data Mn				Fitted			
	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Sample #	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0			
1	0.98	0.00	0.00	0.00	0.00	0.00	1	0.98	0.00	0.00	0.00	0.00	0.00			
2	4.46	0.01	0.00	0.06	0.00	0.00	2	1.92	0.01	0.00	0.03	0.00	0.00			
3	6.93	0.16	0.03	0.22	0.00	0.00	3	4.46	0.26	0.06	0.31	0.00	0.00			
4	9.33	0.59	0.11	0.41	0.00	0.00	4	6.93	1.26	0.28	0.63	0.00	0.26			
5	11.74	0.97	0.19	0.58	0.00	0.01	5	9.33	2.14	0.48	0.83	0.00	0.63			
6	14.23	2.04	0.40	0.72	0.05	0.39	6	11.74	2.89	0.65	0.92	0.67	0.75			
7	16.70	2.69	0.52	0.81	0.90	0.62	7	14.23	3.53	0.79	0.97	1.00	0.83			
8	19.19	3.18	0.62	0.88	0.99	0.71	8	16.70	4.12	0.92	0.98	1.00	0.88			
9	21.66	3.55	0.69	0.92	1.00	0.79	9	19.19	4.19	0.94	0.99	1.00	0.92			
10	25.42	4.62	0.90	0.96	1.00	0.86	10	21.66	4.35	0.97	1.00	1.00	0.94			
11	27.93	4.78	0.93	0.97	1.00	0.90	11	25.42	4.37	0.98	1.00	1.00	0.96			
12	30.47	5.03	0.98	0.98	1.00	0.92	12	27.93	4.38	0.98	1.00	1.00	0.97			
13	33.02	5.00	0.97	0.99	1.00	0.94	13	30.47	4.31	0.96	1.00	1.00	0.98			
14	35.56	5.22	1.02	0.99	1.00	0.96	14	33.02	4.41	0.99	1.00	1.00	0.98			
15	38.13	5.08	0.99	1.00	1.00	0.97	15	35.56	4.33	0.97	1.00	1.00	0.99			
16	40.69	5.25	1.02	1.00	1.00	0.97	16	38.13	4.42	0.99	1.00	1.00	0.99			
17	43.27	5.28	1.03	1.00	1.00	0.98	17	40.69	4.40	0.98	1.00	1.00	0.99			
18	45.80	5.26	1.02	1.00	1.00	0.99	18	43.27	4.54	1.02	1.00	1.00	1.00			
19	48.38	5.29	1.03	1.00	1.00	0.99	19	45.80	4.55	1.02	1.00	1.00	1.00			
20	49.67	5.34	1.04	1.00	1.00	0.99	20	48.38	4.49	1.00	1.00	1.00	1.00			
21	52.26	5.16	1.00	1.00	1.00	0.99	21	49.67	4.52	1.01	1.00	1.00	1.00			
22	54.88	5.13	1.00	1.00	1.00	0.99	22	52.26	4.49	1.00	1.00	1.00	1.00			
23	57.52	5.15	1.00	1.00	1.00	1.00	23	54.88	4.68	1.05	1.00	1.00	1.00			
24	60.18	5.05	0.98	1.00	1.00	1.00	24	57.52	4.82	1.08	1.00	1.00	1.00			

Table C.10 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Manganese (Mn²⁺) transport through lateritic soil under pH 5 with initial concentration 5.14 mmol/L for Pb²⁺ and 4.47 mmol/L for Mn²⁺ (Continued)

25	62.89	5.14	1.00	1.00	1.00	1.00	25	60.18	4.73	1.06	1.00	1.00	1.00
26	65.64	5.09	0.99	1.00	1.00	1.00	26	62.89	4.69	1.05	1.00	1.00	1.00
27	68.40	5.10	0.99	1.00	1.00	1.00	27	65.64	4.51	1.01	1.00	1.00	1.00
28	71.10	4.87	0.95	1.00	1.00	1.00	28	68.40	4.42	0.99	1.00	1.00	1.00
29	73.78	4.88	0.95	1.00	1.00	1.00	29	71.10	4.43	0.99	1.00	1.00	1.00
30	74.05	5.03	0.98	1.00	1.00	1.00	30	73.78	4.25	0.95	1.00	1.00	1.00
31	74.32	5.18	1.01	1.00	1.00	1.00	31	74.05	4.28	0.96	1.00	1.00	1.00
32	74.59	5.24	1.02	1.00	0.98	0.93	32	74.32	4.39	0.98	1.00	0.99	0.98
33	74.86	4.20	0.82	1.00	0.96	0.83	33	74.59	4.38	0.98	1.00	0.94	0.91
34	75.12	4.70	0.92	1.00	0.91	0.74	34	74.86	4.27	0.95	1.00	0.85	0.82
35	75.39	4.67	0.91	1.00	0.87	0.67	35	75.12	4.23	0.95	1.00	0.76	0.73
36	75.66	3.63	0.71	1.00	0.82	0.60	36	75.39	3.89	0.87	0.99	0.69	0.66
37	75.94	2.56	0.50	1.00	0.77	0.56	37	75.66	3.11	0.70	0.98	0.62	0.60
38	76.20	2.43	0.47	1.00	0.73	0.52	38	75.94	2.44	0.54	0.97	0.57	0.55
39	76.46	1.95	0.38	1.00	0.69	0.48	39	76.20	1.90	0.42	0.96	0.52	0.51
40	79.00	0.92	0.18	0.92	0.46	0.33	40	76.46	1.54	0.34	0.94	0.49	0.48
41	81.58	0.69	0.13	0.74	0.35	0.27	41	79.00	0.64	0.14	0.61	0.30	0.32
42	84.09	0.60	0.12	0.55	0.29	0.23	42	81.58	0.44	0.10	0.32	0.23	0.26
43	86.55	0.56	0.11	0.39	0.25	0.21	43	84.09	0.40	0.09	0.15	0.19	0.22
44	90.28	0.48	0.09	0.21	0.20	0.18	44	86.55	0.35	0.08	0.07	0.16	0.19
45	92.81	0.44	0.09	0.14	0.18	0.17	45	90.28	0.30	0.07	0.02	0.13	0.16
46	95.31	0.46	0.09	0.09	0.17	0.16	46	92.81	0.27	0.06	0.01	0.12	0.14
47	97.75	0.41	0.08	0.06	0.15	0.15	47	95.31	0.25	0.06	0.00	0.11	0.13
48	100.20	0.40	0.08	0.04	0.14	0.14	48	97.75	0.24	0.05	0.00	0.10	0.12
49	102.60	0.36	0.07	0.02	0.13	0.13	49	100.20	0.22	0.05	0.00	0.09	0.11
50	107.30	0.36	0.07	0.01	0.11	0.12	50	102.60	0.20	0.05	0.00	0.09	0.10
51	109.78	0.26	0.05	0.01	0.10	0.11	51	107.30	0.17	0.04	0.00	0.08	0.08
52							52	109.78	0.13	0.03	0.00	0.07	0.08

Table C.11 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Manganese (Mn²⁺) transport through lateritic soil under pH 5 with initial concentration 4.85 mmol/L for Pb²⁺ and 8.66 mmol/L for Mn²⁺

Sample #	Observed data Pb			Fitted			Observed data Mn			Fitted		
	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0	Pore volume	Conc.(mM)	C/C0	Linear C/C0	Langmuir C/C0	TSM C/C0
1	0.57	0.00	0.00	0.00	0.00	0.00	0.57	0.00	0.00	0.01	0.00	0.00
2	2.41	0.00	0.00	0.02	0.00	0.00	2.41	0.00	0.00	0.24	0.00	0.00
3	4.92	0.00	0.00	0.14	0.00	0.00	4.92	69.69	0.15	0.71	0.15	0.09
4	7.48	0.46	0.09	0.35	0.00	0.00	7.48	188.58	0.40	0.91	1.00	0.54
5	9.95	1.09	0.21	0.55	0.00	0.16	9.95	271.14	0.57	0.97	1.00	0.65
6	12.37	1.60	0.31	0.71	0.00	0.44	12.37	326.40	0.69	0.99	1.00	0.73
7	14.90	2.17	0.42	0.82	0.44	0.57	14.90	373.80	0.79	1.00	1.00	0.80
8	17.42	2.79	0.54	0.89	1.00	0.67	17.42	412.80	0.87	1.00	1.00	0.85
9	20.54	3.43	0.67	0.93	1.00	0.74	20.54	435.00	0.91	1.00	1.00	0.88
10	23.48	3.85	0.75	0.96	1.00	0.80	23.48	445.50	0.94	1.00	1.00	0.91
11	25.94	4.11	0.80	0.97	1.00	0.85	25.94	444.00	0.93	1.00	1.00	0.93
12	27.15	4.10	0.80	0.98	1.00	0.87	27.15	435.00	0.91	1.00	1.00	0.94
13	29.76	4.44	0.87	0.99	1.00	0.90	29.76	447.60	0.94	1.00	1.00	0.96
14	32.31	4.62	0.90	0.99	1.00	0.92	32.31	456.90	0.96	1.00	1.00	0.97
15	34.88	4.79	0.94	1.00	1.00	0.94	34.88	462.90	0.97	1.00	1.00	0.97
16	37.52	4.71	0.92	1.00	1.00	0.95	37.52	458.10	0.96	1.00	1.00	0.98
17	40.14	4.94	0.96	1.00	1.00	0.96	40.14	466.80	0.98	1.00	1.00	0.99
18	42.78	5.08	0.99	1.00	1.00	0.97	42.78	466.20	0.98	1.00	1.00	0.99
19	45.43	5.22	1.02	1.00	1.00	0.98	45.43	485.70	1.02	1.00	1.00	0.99
20	48.04	5.16	1.01	1.00	1.00	0.98	48.04	493.20	1.04	1.00	1.00	0.99
21	51.95	5.16	1.01	1.00	1.00	0.99	51.95	484.80	1.02	1.00	1.00	1.00
22	54.82	5.14	1.00	1.00	1.00	0.99	54.82	488.40	1.03	1.00	1.00	1.00
23	57.38	5.24	1.02	1.00	1.00	0.99	57.38	486.30	1.02	1.00	1.00	1.00
24	59.96	5.15	1.01	1.00	1.00	1.00	59.96	492.30	1.03	1.00	1.00	1.00

Table C.11 Observed, predicted and simulated breakthrough curves of Lead (Pb²⁺) and Manganese (Mn²⁺) transport through lateritic soil under pH 5 with initial concentration 4.85 mmol/L for Pb²⁺ and 8.66 mmol/L for Mn²⁺ (Continued)

25	62.56	5.18	1.01	1.00	1.00	1.00	25	62.56	496.20	1.04	1.00	1.00	1.00
26	65.17	5.17	1.01	1.00	1.00	1.00	26	65.17	488.40	1.03	1.00	1.00	1.00
27	67.77	5.05	0.99	1.00	1.00	1.00	27	67.77	489.60	1.07	1.00	1.00	1.00
28	70.37	5.21	1.02	1.00	1.00	1.00	28	70.37	488.40	1.07	1.00	1.00	1.00
29	72.72	5.26	1.03	1.00	1.00	1.00	29	72.72	472.50	1.04	1.00	1.00	1.00
30	75.34	5.04	0.98	1.00	1.00	1.00	30	75.34	482.10	1.06	1.00	1.00	1.00
31	77.42	5.03	0.98	1.00	1.00	1.00	31	77.42	484.80	1.06	1.00	1.00	1.00
32	78.11	4.91	0.96	1.00	0.90	0.90	32	78.11	478.80	1.05	1.00	0.87	0.84
33	78.86	4.79	0.94	1.00	0.68	0.68	33	78.86	389.10	0.85	0.98	0.62	0.58
34	79.23	3.68	0.72	1.00	0.61	0.61	34	79.23	298.95	0.66	0.96	0.54	0.51
35	79.52	2.82	0.55	1.00	0.54	0.54	35	79.52	218.76	0.48	0.93	0.47	0.45
36	79.77	2.38	0.46	1.00	0.49	0.49	36	79.77	174.75	0.38	0.89	0.43	0.41
37	80.03	1.98	0.39	0.99	0.46	0.46	37	80.03	157.41	0.35	0.85	0.39	0.38
38	80.29	1.75	0.34	0.99	0.42	0.42	38	80.29	131.58	0.29	0.81	0.36	0.35
39	80.56	1.55	0.30	0.99	0.40	0.40	39	80.56	106.17	0.23	0.75	0.33	0.33
40	81.08	1.26	0.25	0.97	0.35	0.35	40	81.08	88.17	0.19	0.65	0.29	0.30
41	83.72	0.69	0.13	0.82	0.24	0.24	41	83.72	47.49	0.10	0.24	0.19	0.23
42	86.30	0.53	0.10	0.60	0.19	0.19	42	86.30	35.40	0.08	0.07	0.14	0.20
43	88.98	0.54	0.11	0.41	0.16	0.16	43	88.98	31.74	0.07	0.02	0.11	0.17
44	91.63	0.40	0.08	0.26	0.14	0.14	44	91.63	26.95	0.06	0.01	0.09	0.16
45	94.28	0.39	0.08	0.17	0.13	0.13	45	94.28	33.03	0.07	0.00	0.08	0.14
46	96.91	0.23	0.04	0.10	0.12	0.12	46	96.91	22.46	0.05	0.00	0.07	0.13
47	100.77	0.21	0.04	0.05	0.10	0.10	47	100.77	19.59	0.04	0.00	0.06	0.12
48	103.24	0.27	0.05	0.03	0.10	0.10	48	103.24	19.70	0.04	0.00	0.05	0.11
49	105.63	0.27	0.05	0.02	0.09	0.09	49	105.63	24.59	0.05	0.00	0.05	0.10
50	107.85	0.23	0.04	0.01	0.08	0.08	50	107.85	18.38	0.04	0.00	0.04	0.09
51	110.02	0.22	0.04	0.01	0.08	0.08	51	110.02	17.26	0.04	0.00	0.04	0.09
52	111.96	0.24	0.05	0.00	0.08	0.08	52	111.96	17.16	0.04	0.00	0.04	0.08

APPENDIX D *t*-test results for the average SSEs obtaining from the chemical nonequilibrium two-site model and convection-dispersion model with Langmuir model of single metal, binary, and multi-metal systems

Table D.1 *t*-test for SSE of single metal system at pH 4

	Test Value = 0.139					
	t	df	Sig. (2-tailed)	Mean Difference	95% Confidence Interval of the Difference	
					Lower	Upper
EQUIL	-11.689	4	.000	-.0984	-.1218	-.0750

Table D.2 *t*-test for SSE of single metal system at pH 5

	Test Value = 0.0408					
	t	df	Sig. (2-tailed)	Mean Difference	95% Confidence Interval of the Difference	
					Lower	Upper
EQUIL	5.046	4	.007	.1398	.0629	.2167

Table D.3 *t*-test for SSE of Pb^{2+} in Pb^{2+} - Ni^{2+} system

	Test Value = 0.03075					
	t	df	Sig. (2-tailed)	Mean Difference	95% Confidence Interval of the Difference	
					Lower	Upper
EQUIL	7.078	3	.006	.0843	.0464	.1221

Table D.4 *t*-test for SSE of Ni^{2+} in Pb^{2+} - Ni^{2+} system

	Test Value = 0.035					
	t	df	Sig. (2-tailed)	Mean Difference	95% Confidence Interval of the Difference	
					Lower	Upper
EQUIL	5.800	3	.010	.0725	.0327	.1123

Table D.5 *t*-test for SSE of Pb^{2+} in Pb^{2+} - Zn^{2+} system

	Test Value = 0.046					
	t	df	Sig. (2-tailed)	Mean Difference	95% Confidence Interval of the Difference	
					Lower	Upper
EQUIL	3.963	3	.029	.0815	.0161	.1469

Table D.6 *t*-test for SSE of Zn^{2+} in Pb^{2+} - Zn^{2+} system

	Test Value = 0.04					
	t	df	Sig. (2-tailed)	Mean Difference	95% Confidence Interval of the Difference	
					Lower	Upper
EQUIL	3.806	3	.032	.0325	.0053	.0597

Table D.7 *t*-test for SSE of Pb^{2+} in Pb^{2+} - Mn^{2+} system

	Test Value = 0.03933					
	t	df	Sig. (2-tailed)	Mean Difference	95% Confidence Interval of the Difference	
					Lower	Upper
EQUIL	12.240	2	.007	.0707	.0458	.0955

Table D.8 *t*-test for SSE of Mn^{2+} in Pb^{2+} - Mn^{2+} system

	Test Value = 0.0333					
	t	df	Sig. (2-tailed)	Mean Difference	95% Confidence Interval of the Difference	
					Lower	Upper
EQUIL	5.673	2	.030	.0500	.0121	.0880

Table D.9 *t*-test for SSE of multi-metal system

	Test Value = 0.04					
	t	df	Sig. (2-tailed)	Mean Difference	95% Confidence Interval of the Difference	
					Lower	Upper
EQUIL	5.166	3	.014	.0325	.0125	.0525

BIOGRAPHY

Srilert Chotpantarat was born on 6 January 1977, in the Songkhla province. He received a bachelor's degree in Civil Engineering in 1997 from the Faculty of Engineering, Chulalongkorn University and a master's degree in Water Resources Engineering in 2002 from the Faculty of Engineering, Chulalongkorn University. He was awarded a scholarship by the Geology Department, Faculty of Science, Chulalongkorn University in 2001. He enrolled in the International Post-graduate Program in Environmental Management, National Center of Excellence for Environmental and Hazardous Waste Management, Chulalongkorn University in 2004 and completed his Doctor of Philosophy in Environmental Management in 2008.



