

REFERENCES

1. **Castilho, L. R., Mitchell, D. A., Freire, D. M.**: Production of polyhydroxyalkanoates (PHAs) from waste materials and by-products by submerged and solid-state fermentation, *Bioresour Technol*, **100**, 5996-6009 (2009).
2. **Cavalheiro, J. M. B. T., de Almeida, M. C. M. D., Grandfils, C., da Fonseca, M. M. R.**: Poly(3-hydroxybutyrate) production by *Cupriavidus necator* using waste glycerol, *Process Biochem*, **44**, 509-515 (2009).
3. **Chee, J. Y., Tan, Y., Samian, M. R., Sudesh, K.**: Isolation and characterization of a *Burkholderia* sp. USM (JCM15050) capable of producing polyhydroxyalkanoate (PHA) from triglycerides, fatty acids and glycerols, *J Polym Environ*, **18**, 584-592 (2010).
4. **Sudesh, K., Abe, H., Doi, Y.**: Synthesis, structure and properties of polyhydroxyalkanoates: biological polyesters, *Prog Polym Sci*, **25**, 1503-1555 (2000).
5. **Harding, K. G., Dennis, J. S., von Blottnitz, H., Harrison, S. T.**: Environmental analysis of plastic production processes: comparing petroleum-based polypropylene and polyethylene with biologically-based poly-beta-hydroxybutyric acid using life cycle analysis, *J Biotechnol*, **130**, 57-66 (2007).
6. **Jung, Y. K., Lee, S. Y., Tam, T. T.**: Towards systems metabolic engineering of PHA producers, plastic from bacteria, *14*, 63-84 (2010).
7. **Matsusaki, H., Abe, H., Doi, Y.**: Biosynthesis and properties of poly(3-hydroxybutyrate-co-3-hydroxyalkanoates) by recombinant strains of *Pseudomonas* sp. 61-3, *Biomacromolecules*, **1**, 17-22 (2000).
8. **Abe, H., Doi, Y., Fukushima, T., Eya, H.**: Biosynthesis from gluconate of a random copolyester consisting of 3-hydroxybutyrate and medium-chain-length 3-hydroxyalkanoates by *Pseudomonas* sp. 61-3, *Int J Biol Macromol*, **16**, 115-119 (1994).

- 9.**Doi, Y., Kitamura, S., Abe, H.:** Microbial synthesis and characterization of poly(3-hydroxybutyrate-co-3-hydroxyhexanoate), *Macromolecules*, **28**, 4822-4828 (1995).
- 10.**Lee, S. H., Oh, D. H., Ahn, W. S., Lee, Y., Choi, J., Lee, S. Y.:** Production of poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) by high-cell-density cultivation of *Aeromonas hydrophila*, *Biotechnol Bioeng*, **67**, 240-244 (2000).
- 11.**Janda, J. M., Abbott, S. L.:** The genus *Aeromonas*: taxonomy, pathogenicity, and infection, *Clin Microbiol Rev*, **23**, 35-73 (2010).
- 12.**Tsuge, T.:** Metabolic improvements and use of inexpensive carbon sources in microbial production of polyhydroxyalkanoates, *J Biosci Bioeng*, **94**, 579-584 (2002).
- 13.**Fukui, T., Doi, Y.:** Cloning and analysis of the poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) biosynthesis genes of *Aeromonas caviae*, *J Bacteriol*, **179**, 4821-4830 (1997).
- 14.**Antonio, R. V., Steinbuchel, A., Rehm, B. H.:** Analysis of in vivo substrate specificity of the PHA synthase from *Ralstonia eutropha*: formation of novel copolymers in recombinant *Escherichia coli*, *FEMS Microbiol Lett*, **182**, 111-117 (2000).
- 15.**Nomura, C. T., Taguchi, K., Taguchi, S., Doi, Y.:** Coexpression of genetically engineered 3-ketoacyl-ACP synthase III (fabH) and polyhydroxyalkanoate synthase (phaC) genes leads to short-chain-length-medium-chain-length polyhydroxyalkanoate copolymer production from glucose in *Escherichia coli* JM109, *Appl Environ Microbiol*, **70**, 999-1007 (2004).
- 16.**Zou, X. H., Chen, G. Q.:** Metabolic engineering for microbial production and applications of copolymers consisting of 3-hydroxybutyrate and medium-chain-length 3-hydroxyalkanoates, *Macromol Biosci*, **7**, 174-182 (2007).
- 17.**Choi, J., Lee, S. Y.:** Process analysis and economic evaluation for Poly(3-hydroxybutyrate) production by fermentation, *Bioprocess Eng* **17**, 335-342 (1997).

18. **Yazdani, S. S., Gonzalez, R.**: Anaerobic fermentation of glycerol: a path to economic viability for the biofuels industry, *Curr Opin Biotechnol*, **18**, 213-219 (2007).
19. **Phithakrotchanakoon, C., Champreda, V., Aiba, S., Pootanakit, K., Tanapongpipat, S.**: Engineered *Escherichia coli* for Short-Chain-Length Medium-Chain-Length polyhydroxyalkanoate copolymer biosynthesis from glycerol and dodecanoate, *Biosci Biotechnol Biochem*, **77**, 1262-1268 (2013).
20. **Lemoigne, M.**: Products of dehydration and of polymerization of 3-hydroxybutyric acid, *Bull Soc Chim Biol*, **8**, 770-782 (1926).
21. **Zhu, C., Chiu, S., Nakas, J. P., Nomura, T. N.**: Bioplastics from waste glycerol derived from biodiesel industry, *J Appl Polym Sci*, **130**, 1-13 (2013).
22. **Dawes, E. A., Senior, P. J.**: The role and regulation of energy reserve polymers in microorganisms, *Adv Microb Physiol*, **10**, 135-266 (1973).
23. **Findlay, R. H., White, D. C.**: Polymeric beta-hydroxyalkanoates from environmental samples and *Bacillus megaterium*, *Appl Environ Microbiol*, **45**, 71-78 (1983).
24. **Wallen, L. L., Rohwedder, W. K.**: Poly-beta-hydroxyalkanoate from activated sludge, *Environ Sci Technol*, **8**, 576-579 (1974).
25. **de Smet, M. J., Eggink, G., Witholt, B., Kingma, J., Wynberg, H.**: Characterization of intracellular inclusions formed by *Pseudomonas oleovorans* during growth on octane, *J Bacteriol*, **154**, 870-878 (1983).
26. **Lageveen, R. G., Huisman, G. W., Preusting, H., Ketelaar, P., Eggink, G., Witholt, B.**: Formation of polyesters by *Pseudomonas oleovorans*: effect of substrates on formation and composition of poly-(R)-3-hydroxyalkanoates and poly-(R)-3-hydroxyalenoates, *Appl Environ Microbiol*, **54**, 2924-2932 (1988).
27. **Slater, S. C., Voige, W. H., Dennis, D. E.**: Cloning and expression in *Escherichia coli* of the *Alcaligenes eutrophus* H16 poly-beta-hydroxybutyrate biosynthetic pathway, *J Bacteriol*, **170**, 4431-4436 (1988).

28. **Schubert, P., Steinbuchel, A., Schlegel, H. G.**: Cloning of the *Alcaligenes eutrophus* genes for synthesis of poly-beta-hydroxybutyric acid (PHB) and synthesis of PHB in *Escherichia coli*, *J Bacteriol*, **170**, 5837-5847 (1988).
29. **Rehm, B. H.**: Polyester synthases: natural catalysts for plastics, *Biochem J*, **376**, 15-33 (2003).
30. **Verlinden, R. A., Hill, D. J., Kenward, M. A., Williams, C. D., Radecka, I.**: Bacterial synthesis of biodegradable polyhydroxyalkanoates, *J Appl Microbiol*, **102**, 1437-1449 (2007).
31. **Asrar, J., Valentin, H. E., Berger, P. A., Tran, M., Padgett, S. R., Garbow, J. R.**: Biosynthesis and properties of poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) polymers, *Biomacromolecules*, **3**, 1006-1012 (2002).
32. **Wang, Y. W., Yang, F., Wu, Q., Cheng, Y. C., Yu, P. H. F., Chen, J., Chen, G. Q.**: Effect of composition of poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) on growth of fibroblast and osteoblast, *Biomaterials*, **26**, 755-761 (2005).
33. **Madison, L. L., Huisman, G. W.**: Metabolic engineering of poly(3-hydroxyalkanoates): from DNA to plastic, *Microbiol Mol Biol Rev*, **63**, 21-53 (1999).
34. **Rehm, B. H.**: Bacterial polymers: biosynthesis, modifications and applications, *Nat Rev Microbiol*, **8**, 578-592 (2010).
35. **Potter, M., Madkour, M. H., Mayer, F., Steinbuchel, A.**: Regulation of phasin expression and polyhydroxyalkanoate (PHA) granule formation in *Ralstonia eutropha* H16, *Microbiology*, **148**, 2413-2426 (2002).
36. **Dawes, E. A., Senior, P. J.**: The role and regulation of energy reserve polymers in micro-organisms, *Adv Microb Physiol*, **10**, 135-266 (1973).
37. **Jendrossek, D., Handrick, R.**: Microbial degradation of polyhydroxyalkanoates, *Annu Rev Microbiol*, **56**, 403-432 (2002).
38. **Anderson, A. J., Dawes, E. A.**: Occurrence, metabolism, metabolic role, and industrial uses of bacterial polyhydroxyalkanoates, *Microbiol Rev*, **54**, 450-472 (1990).
39. **Steinbüchel, A., Valentin, H. E.**: Diversity of bacterial polyhydroxyalkanoic acids, *FEMS Microbiol Lett*, **128**, 219-228 (1995).

40. **Kessler, B., Witholt, B.**: Factors involved in the regulatory network of polyhydroxyalkanoate metabolism, *J Biotechnol*, **86**, 97-104 (2001).
41. **Park, S. J., Choi, J., Lee, S. Y.**: Engineering of *Escherichia coli* fatty acid metabolism for the production of polyhydroxyalkanoates, *Enzyme Microb Tech*, **36**, 579-588 (2005).
42. **Lennen, R. M., Pfleger, B. F.**: Engineering *Escherichia coli* to synthesize free fatty acids, *Trends Biotechnol*, **30**, 659-667 (2012).
43. **Rehm, B. H., Kruger, N., Steinbuchel, A.**: A new metabolic link between fatty acid de novo synthesis and polyhydroxyalkanoic acid synthesis. The PHAG gene from *Pseudomonas putida* KT2440 encodes a 3-hydroxyacyl-acyl carrier protein-coenzyme a transferase, *J Biol Chem*, **273**, 24044-24051 (1998).
44. **Valentin, H. E., Steinbuchel, A.**: Cloning and characterization of the *Methylobacterium extorquens* polyhydroxyalkanoic-acid-synthase structural gene, *Appl Microbiol Biotechnol*, **39**, 309-317 (1993).
45. **Tombolini, R., Povolo, S., Buson, A., Squartini, A., Nuti, M. P.**: Poly-beta-hydroxybutyrate (PHB) biosynthetic genes in *Rhizobium meliloti* 41, *Microbiology*, **141**, 2553-2559 (1995).
46. **Ramachander, T. V., Rawal, S. K.**: PHB synthase from *Streptomyces aureofaciens* NRRL 2209, *FEMS Microbiol Lett*, **242**, 13-18 (2005).
47. **Khanna, S., Srivastava, A. K.**: Recent advances in microbial polyhydroxyalkanoates, *Process Biochem*, **40**, 607-619 (2005).
48. **Reinecke, F., Steinbüchel, A.**: *Ralstonia eutropha* strain H16 as model organism for PHA metabolism and for biotechnological production of technically interesting biopolymers, *J Mol Microb Biotech*, **16**, 91-108 (2009).
49. **Pohlmann, A., Fricke, W. F., Reinecke, F., Kusian, B., Liesegang, H., Cramm, R., Eitinger, T., Ewering, C., Potter, M., Schwartz, E. et al.**: Genome sequence of the bioplastic-producing "Knallgas" bacterium *Ralstonia eutropha* H16, *Nat Biotechnol*, **24**, 1257-1262 (2006).
50. **Luengo, J. M., García, B., Sandoval, A., Naharro, G., Olivera, E. R.**: Bioplastics from microorganisms, *Curr Opin Microbiol*, **6**, 251-260 (2003).

51. **Kim, B. S., Chul Lee, S. C., Lee, S. Y., Chang, H. N., Chang, Y. K., Woo, S. L.:** Production of poly(3-hydroxybutyric acid) by fed-batch culture of *Alcaligenes eutrophus* with glucose concentration control, *Biotechnol Bioeng*, **43**, 892-898 (1994).
52. **Doi, Y., Segawa, A., Nakamura, S., Kunioka, M.:** Production of biodegradable copolyesters by *Alcaligenes eutrophus*, *Novel biodegradable microbial polymers*, **186**, 37-48 (1990).
53. **Du, G. C., Chen, J., Yu, J., Lun, S.:** Feeding strategy of propionic acid for production of poly(3-hydroxybutyrate-*co*-3-hydroxyvalerate) with *Ralstonia eutropha*, *Biochem Eng J*, **8**, 103-110 (2001).
54. **Green, P. R., Kemper, J., Schechtman, L., Guo, L., Satkowski, M., Fiedler, S., Steinbuchel, A., Rehm, B. H.:** Formation of short chain length/medium chain length polyhydroxyalkanoate copolymers by fatty acid beta-oxidation inhibited *Ralstonia eutropha*, *Biomacromolecules*, **3**, 208-213 (2002).
55. **Silva-Queiroz, S. R., Silva, L. F., Pradella, J. G. C., Pereira, E. M., Gomez, J. G. C.:** PHA(MCL) biosynthesis systems in *Pseudomonas aeruginosa* and *Pseudomonas putida* strains show differences on monomer specificities, *J Biotechnol*, **143**, 111-118 (2009).
56. **Huijberts, G. N. M., Eggink, G.:** Production of poly(3-hydroxyalkanoates) by *Pseudomonas putida* KT2442 in continuous cultures, *Appl Microbiol Biotechnol*, **46**, 233-239 (1996).
57. **Ramsay, B. A., Saracovan, I., Ramsay, J. A., Marchessault, R. H.:** Continuous production of long-side-chain poly-beta-hydroxyalkanoates by *Pseudomonas oleovorans*, *Appl Environ Microbiol*, **57**, 625-629 (1991).
58. **Haywood, G. W., Anderson, A. J., Ewing, D. F., Dawes, E. A.:** Accumulation of a polyhydroxyalkanoate containing primarily 3-hydroxydecanoate from simple carbohydrate substrates by *Pseudomonas* sp. strain NCIMB 40135, *Appl Environ Microbiol*, **56**, 3354-3359 (1991).
59. **Timm, A., Steinbüchel, A.:** Formation of polyesters consisting of medium-chain-length 3-hydroxyalkanoic acids from gluconate by *Pseudomonas*

- aeruginosa* and other fluorescent *pseudomonads*, Appl Environ Microbiol, **56**, 3360-3367 (1990).
60. **Witholt, B., Kessler, B.**: Perspectives of medium chain length poly(hydroxyalkanoates), a versatile set of bacterial bioplastics, Curr Opin Biotechnol, **10**, 279-285 (1999).
61. **Timmis, K. N.**: *Pseudomonas putida*: a cosmopolitan opportunist par excellence, Environ Microbiol, **4**, 779-781 (2002).
62. **Wang, Q., Nomura, C. T.**: Monitoring differences in gene expression levels and polyhydroxyalkanoate (PHA) production in *Pseudomonas putida* KT2440 grown on different carbon sources, J Biosci Bioeng, **110**, 653-659 (2010).
63. **Nelson, K. E., Weinel, C., Paulsen, I. T., Dodson, R. J., Hilbert, H., dos Santos, V. A. P. M., Fouts, D. E., Gill, S. R., Pop, M., Holmes, M. et al**: Complete genome sequence and comparative analysis of the metabolically versatile *Pseudomonas putida* KT2440, Environ Microbiol, **4**, 799-808 (2002).
64. **Huisman, G. W., Wonink, E., Meima, R., Kazemier, B., Terpstra, P., Witholt, B.**: Metabolism of poly(3-hydroxyalkanoates) (PHAs) by *Pseudomonas oleovorans*. Identification and sequences of genes and function of the encoded proteins in the synthesis and degradation of PHA, J Biol Chem, **266**, 2191-2198 (1991).
65. **Timm, A., Steinbuchel, A.**: Cloning and molecular analysis of the poly(3-hydroxyalkanoic acid) gene locus of *Pseudomonas aeruginosa* PAO1, Eur J Biochem, **209**, 15-30 (1992).
66. **Matsusaki, H., Manji, S., Taguchi, K., Kato, M., Fukui, T., Doi, Y.**: Cloning and molecular analysis of the Poly(3-hydroxybutyrate) and Poly(3-hydroxybutyrate-co-3-hydroxyalkanoate) biosynthesis genes in *Pseudomonas* sp. strain 61-3, J Bacteriol, **180**, 6459-6467 (1998).
67. **Kim, T. K., Shin, H. D., Seo, M. C., Lee, J. N., Lee, Y. H.**: Molecular structure of PCR cloned PHA synthase genes of *Pseudomonas putida* KT2440 and its utilization for medium-chain length polyhydroxyalkanoate production, J Microbiol Biotechnol, **13**, 182-190 (2003).

68. **Tsuge, T., Taguchi, K., Taguchi, S., Doi, Y.:** Molecular characterization and properties of (R)-specific enoyl-CoA hydratases from *Pseudomonas aeruginosa*: metabolic tools for synthesis of polyhydroxyalkanoates via fatty acid beta-oxidation, *Int J Biol Macromol*, **31**, 195-205 (2003).
69. **Sato, S., Kanazawa, H., Tsuge, T.:** Expression and characterization of (R)-specific enoyl coenzyme A hydratases making a channeling route to polyhydroxyalkanoate biosynthesis in *Pseudomonas putida*, *Appl Microbiol Biotechnol*, **90**, 951-959 (2011).
70. **Seshadri, R., Joseph, S. W., Chopra, A. K., Sha, J., Shaw, J., Graf, J., Haft, D., Wu, M., Ren, Q., Rosovitz, M. J. et al:** Genome sequence of *Aeromonas hydrophila* ATCC 7966T: jack of all trades, *J Bacteriol*, **188**, 8272-8282 (2006).
71. **Chen, G. Q., Zhang, G., Park, S. J., Lee, S. Y.:** Industrial scale production of poly(3-hydroxybutyrate-co-3-hydroxyhexanoate), *Appl Microbiol Biotechnol*, **57**, 50-55 (2001).
72. **Lu, X. Y., Wu, Q., Chen, G. Q.:** Production of poly(3-hydroxybutyrate-*co*-3-hydroxyhexanoate) with flexible 3-hydroxyhexanoate content in *Aeromonas hydrophila* CGMCC 0911, *Appl Microbiol Biotechnol*, **64**, 41–45 (2004).
73. **Kato, M., Bao, H. J., Kang C. K., Fukui, T., Doi, Y.:** Production of a novel copolyester of 3-hydroxybutyric acid and medium-chain-length 3-hydroxyalkanoic acids by *Pseudomonas* sp. 61-3 from sugars, *Appl Microbiol Biotechnol*, **45**, 363-370 (1996).
74. **Steinbüchel, A., Wiese, S.:** A *Pseudomonas* strain accumulating polyesters of 3-hydroxybutyric acid and medium-chain-length 3-hydroxyalkanoic acids, *Appl Microbiol Biotechnol*, **37**, 691-697 (1992).
75. **Liebergesell, M., Hustede, E., Timm, A., Steinbüchel, A., Fuller, R. C., Lenz, R. W., Schlegel, H. G.:** Formation of poly(3-hydroxyalkanoates) by phototrophic and chemolithotrophic bacteria, *Arch Microbiol*, **155**, 415-421 (1991).
76. **Haywood, G. W., Anderson, A. J., Williams, D. R., Dawes, E. A., Ewing, D. F.:** Accumulation of a poly(hydroxyalkanoate) copolymer containing

- primarily 3-hydroxyvalerate from simple carbohydrate substrates by *Rhodococcus* sp. NCIMB 40126, *Int J Biol Macromol*, **13**, 83-88 (1991).
77. **Fukui, T., Shiomi, N., Doi, Y.**: Expression and characterization of (*R*)-specific enoyl coenzyme A hydratase involved in polyhydroxyalkanoate biosynthesis by *Aeromonas caviae*, *J Bacteriol*, **180**, 667-673 (1998).
78. **Lee, S. Y.**: *E. coli* moves into the plastic age, *Nat Biotechnol*, **15**, 1997 (1997).
79. **Lee, S. Y., Choi, J. I.**: Production of microbial polyester by fermentation of recombinant microorganisms, *Adv Biochem Eng Biotechnol*, **71**, 183-207 (2001).
80. **Li, R., Zhang, H., Qi, Q.**: The production of polyhydroxyalkanoates in recombinant *Escherichia coli*, *Bioresour Technol*, **98**, 2313-2320 (2007).
81. **Aldor, I. S., Keasling, J. D.**: Process design for microbial plastic factories: metabolic engineering of polyhydroxyalkanoates, *Curr Opin Biotechnol*, **14**, 475-483 (2003).
82. **Miyake, M., Miyamoto, C., Schnackenberg, J., Kurane, R., Asada, Y.**: Phosphotransacetylase as a key factor in biological production of polyhydroxybutyrate, *Appl Biochem Biotechnol*, **84-86**, 1039-1044 (2000).
83. **Kabir, M. M., Shimizu, K.**: Fermentation characteristics and protein expression patterns in a recombinant *Escherichia coli* mutant lacking phosphoglucose isomerase for poly(3-hydroxybutyrate) production, *Appl Microbiol Biotechnol*, **62**, 244-255 (2003).
84. **Jung, Y. M., Lee, J. N., Shin, H. D., Lee, Y. H.**: Role of *tktA* gene in pentose phosphate pathway on odd-ball biosynthesis of poly-beta-hydroxybutyrate in transformant *Escherichia coli* harboring phbCAB operon, *J Biosci Bioeng*, **98**, 224-227 (2004).
85. **Fiedler, S., Steinbuchel, A., Rehm, B. H.**: The role of the fatty acid beta-oxidation multienzyme complex from *Pseudomonas oleovorans* in polyhydroxyalkanoate biosynthesis: molecular characterization of the *fadBA* operon from *P. oleovorans* and of the enoyl-CoA hydratase genes *phaJ* from *P. oleovorans* and *Pseudomonas putida*, *Arch Microbiol*, **178**, 149-160 (2002).

86. **Tsuge, T., Fukui, T., Matsusaki, H., Taguchi, S., Kobayashi, G., Ishizaki, A., Doi, Y.**: Molecular cloning of two (R)-specific enoyl-CoA hydratase genes from *Pseudomonas aeruginosa* and their use for polyhydroxyalkanoate synthesis, FEMS Microbiol Lett, **184**, 193-198 (2000).
87. **Ren, Q., Sierro, N., Witholt, B., Kessler, B.**: FabG, an NADPH-dependent 3-ketoacyl reductase of *Pseudomonas aeruginosa*, provides precursors for medium-chain-length poly-3-hydroxyalkanoate biosynthesis in *Escherichia coli*, J Bacteriol, **182**, 2978-2981 (2000).
88. **Taguchi, K., Aoyagi, Y., Matsusaki, H., Fukui, T., Doi, Y.**: Co-expression of 3-ketoacyl-ACP reductase and polyhydroxyalkanoate synthase genes induces PHA production in *Escherichia coli* HB101 strain, FEMS Microbiol Lett, **176**, 183-190 (1999).
89. **Hoffmann, N., Steinbüchel, A., Rehm, B. H.**: The *Pseudomonas aeruginosa* *phaG* gene product is involved in the synthesis of polyhydroxyalkanoic acid consisting of medium-chain-length constituents from non-related carbon sources, FEMS Microbiol Lett, **184**, 253-259 (2000).
90. **Ren, Q., Sierro, N., Kellerhals, M., Kessler, B., Witholt, B.**: Properties of engineered poly-3-hydroxyalkanoates produced in recombinant *Escherichia coli* strains, Appl Environ Microbiol, **66**, 1311-1320 (2000).
91. **Park, S. J., Park, J. P., Lee, S. Y.**: Metabolic engineering of *Escherichia coli* for the production of medium-chain-length polyhydroxyalkanoates rich in specific monomers, FEMS Microbiol Lett, **214**, 217-222 (2002).
92. **Qi, Q., Steinbuchel, A., Rehm, B. H.**: Metabolic routing towards polyhydroxyalkanoic acid synthesis in recombinant *Escherichia coli* (*fadR*): inhibition of fatty acid beta-oxidation by acrylic acid, FEMS Microbiol Lett, **167**, 89-94 (1998).
93. **Langenbach, S., Rehm, B. H., Steinbuchel, A.**: Functional expression of the PHA synthase gene *phaC1* from *Pseudomonas aeruginosa* in *Escherichia coli* results in poly(3-hydroxyalkanoate) synthesis, FEMS Microbiol Lett, **150**, 303-309 (1997).
94. **Park, S. J., Lee, S. Y.**: Identification and characterization of a new enoyl coenzyme A hydratase involved in biosynthesis of medium-chain-length

- polyhydroxyalkanoates in recombinant *Escherichia coli*, *J Bacteriol*, **185**, 5391-5397 (2003).
95. **Park, S. J., Ahn, W. S., Green, P. R., Lee, S. Y.**: Production of poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) by metabolically engineered *Escherichia coli* strains, *Biomacromolecules*, **2**, 248-254 (2001).
96. **Davis, R., Anilkumar, P. K., Chandrashekhar, A., Shamala, T. R.**: Biosynthesis of polyhydroxyalkanoates co-polymer in *E. coli* using genes from *Pseudomonas* and *Bacillus*, *Anton Leeuw Int J G*, **94**, 207-216 (2008).
97. **Tokiwa, Y., Calabia, B. P.**: Degradation of microbial polyesters, *Biotechnol Lett*, **26**, 1181-1189 (2004).
98. **Phithakrotchanakoon, C., Rudeekit, Y., Tanapongpipat, S., Leejakpai, T., Aiba, S., Noda, I., Champreda, V.**: Microbial degradation and physico-chemical alteration of polyhydroxyalkanoates by a thermophilic *Streptomyces* sp., *Biologia*, **64**, 246-251 (2009).
99. **Philip, S., Keshavarz, T., Roy, I.**: Polyhydroxyalkanoates: biodegradable polymers with a range of applications, *J Chem Technol Biotechnol*, **82**, 233-247 (2007).
100. **Mergaert, J., Anderson, C., Wouters, A., Swings, J.**: Microbial degradation of poly(3-hydroxybutyrate) and poly(3-hydroxybutyrate-co-3-hydroxyvalerate) in compost, *J Environ Polym Degr*, **2**, 177-183 (1994).
101. **Mergaert, J., Webb, A., Anderson, C., Wouters, A., Swings, J.**: Microbial degradation of poly(3-hydroxybutyrate) and poly(3-hydroxybutyrate-co-3-hydroxyvalerate) in soils, *Appl Environ Microbiol*, **59**, 3233-3238 (1993).
102. **Mergaert, J., Wouters, A., Anderson, C., Swings, J.**: In situ biodegradation of poly(3-hydroxybutyrate) and poly(3-hydroxybutyrate-co-3-hydroxyvalerate) in natural waters, *Can J Microbiol*, **41**, 154-159 (1995).
103. **Mukai, K., Doi, Y., Sema, Y., Tomita, K.**: Substrate specificities in hydrolysis of polyhydroxyalkanoates by microbial esterases, *Biotechnol Lett*, **15**, 601-604 (1993).

- 104.**Lobler, M., Sass, M., Michel, P., Hopt, U. T., Kunze, C., Schmitz, K. P.:**
Differential gene expression after implantation of biomaterials into rat
gastrointestine, *J Mater Sci Mater Med*, **10**, 797-799 (1999).
- 105.**Volova, T., Shishatskaya, G., Sevastianov, V., Efremov, S., Mogilnaya, O.:**
Results of biomedical investigations of PHB and PHB/PHV fibers,
Biochem Eng J, **16**, 125-133 (2003).
- 106.**Ye, C., Hu, P., Ma, M. X., Xiang, Y., Liu, R. G., Shang, X. W.:** PHB/PHBHHx
scaffolds and human adipose-derived stem cells for cartilage tissue
engineering, *Biomaterials*, **30**, 4401-4406 (2009).
- 107.**Chen, G. Q.:** Plastics completely synthesized by bacteria:
polyhydroxyalkanoates, *Microbiology Monographs (plastic from bacteria)*,
14, 17-37 (2010).
- 108.**Chen, G. Q., Wu, Q.:** The application of polyhydroxyalkanoates as tissue
engineering materials, *Biomaterials*, **26**, 6565-6578 (2005).
- 109.**Chatzifragkou, A., Papanikolaou, S.:** Effect of impurities in biodiesel-derived
waste glycerol on the performance and feasibility of biotechnological
processes, *Appl Microbiol Biotechnol*, **95**, 13-27 (2012).
- 110.**Yang, F., Hanna, M. A., Sun, R.:** Value-added uses for crude glycerol--a
byproduct of biodiesel production, *Biotechnol Biofuels*, **5**, 13 (2012).
- 111.**da Silva, G. P., Mack, M., Contiero, J.:** Glycerol: a promising and abundant
carbon source for industrial microbiology, *Biotechnol Adv*, **27**, 30-39
(2009).
- 112.**Peters, D.:** Raw materials, *Adv Biochem Eng Biotechnol*, **105**, 1-30 (2007).
- 113.**Murarka, A., Dharmadi, Y., Yazdani, S. S., Gonzalez, R.:** Fermentative
utilization of glycerol by *Escherichia coli* and its implications for the
production of fuels and chemicals, *Appl Environ Microbiol*, **74**, 1124-1135
(2008).
- 114.**Johnson, D. T., Taconi, K. A.:** The glycerin glut: Options for the value-added
conversion of crude glycerol resulting from biodiesel production, *Environ
Prog*, **26**, 338-348 (2007).

115. **Cerrate, S., Yan, F., Wang, Z., Coto, C., Sacaklı, P., Waldroup, P. W.:** Evaluation of glycerine from biodiesel production as a feed ingredient for broilers, *Int J Poult Sci*, **5**, 1001-1007 (2006).
116. **Dasaria, M. A., Kiatsimkula, P., Sutterlinb, W. R., Suppes, G. J.:** Low-pressure hydrogenolysis of glycerol to propylene glycol, *Appl Catal A: Gen*, **281**, 225-231 (2005).
117. **Mattam, A. J., Clomburg, J. M., Gonzalez, R., Yazdani, S. S.:** Fermentation of glycerol and production of valuable chemical and biofuel molecules, *Biotechnol Lett*, **35**, 831-842 (2013).
118. **Ito, T., Nakashimada, Y., Senba, K., Matsui, T., Nishio, N.:** Hydrogen and ethanol production from glycerol-containing wastes discharged after biodiesel manufacturing process, *J Biosci Bioeng*, **100**, 260-265 (2005).
119. **Kenny, S. T., Runic, J. N., Kaminsky, W., Woods, T., Babu, R. P., O'Connor, K. E.:** Development of a bioprocess to convert PET derived terephthalic acid and biodiesel derived glycerol to medium chain length polyhydroxyalkanoate, *Appl Microbiol Biotechnol*, **95**, 623-633 (2012).
120. **Moralejo-Garate, H., Mar'atusalihat, E., Kleerebezem, R., van Loosdrecht, M. C.:** Microbial community engineering for biopolymer production from glycerol, *Appl Microbiol Biotechnol*, **92**, 631-639 (2011).
121. **Kang, Z., Du, L. L., Kang, J. H., Wang, Y., Wang, Q., Liang, Q. F., Qi, Q. S.:** Production of succinate and polyhydroxyalkanoate from substrate mixture by metabolically engineered *Escherichia coli*, *Bioresour Technol*, **102**, 6600-6604 (2011).
122. **Kawata, Y., Aiba, S.:** Poly(3-hydroxybutyrate) production by isolated *Halomonas* sp. KM-1 using waste glycerol, *Biosci Biotechnol Biochem*, **74**, 175-177 (2010).
123. **Shrivastav, A., Mishra, S. K., Shethia, B., Pancha, I., Jain, D., Mishra, S.:** Isolation of promising bacterial strains from soil and marine environment for polyhydroxyalkanoates (PHAs) production utilizing Jatropha biodiesel byproduct, *Int J Biol Macromol*, **47**, 283-287 (2010).
124. **Zhu, C., Nomura, C. T., Perrotta, J. A., Stipanovic, A. J., Nakas, J. P.:** Production and characterization of poly-3-hydroxybutyrate from biodiesel-

- glycerol by *Burkholderia cepacia* ATCC 17759, *Biotechnol Prog*, **26**, 424-430 (2010).
125. **Andreessen, B., Lange, A. B., Robenek, H., Steinbuchel, A.:** Conversion of glycerol to poly(3-Hydroxypropionate) in recombinant *Escherichia coli*, *Appl Environ Microbiol*, **76**, 622-626 (2010).
126. **Ibrahim, M. H., Steinbuchel, A.:** *Zobellella denitrificans* strain MW1, a newly isolated bacterium suitable for poly(3-hydroxybutyrate) production from glycerol, *J Appl Microbiol*, **108**, 214-225 (2010).
127. **Ibrahim, M. H., Steinbuchel, A.:** Poly(3-hydroxybutyrate) production from glycerol by *Zobellella denitrificans* MW1 via high-cell-density fed-batch fermentation and simplified solvent extraction, *Appl Environ Microbiol*, **75**, 6222-6231 (2009).
128. **Nikel, P. I., Pettinari, M. J., Galvagno, M. A., Mendez, B. S.:** Poly(3-hydroxybutyrate) synthesis from glycerol by a recombinant *Escherichia coli* arcA mutant in fed-batch microaerobic cultures, *Appl Microbiol Biotechnol*, **77**, 1337-1343 (2008).
129. **Almeida, A., Nikel, P. I., Giordano, A. M., Pettinari, M. J.:** Effects of granule-associated protein PhaP on glycerol-dependent growth and polymer production in poly(3-hydroxybutyrate)-producing *Escherichia coli*, *Appl Environ Microbiol*, **73**, 7912-7916 (2007).
130. **Mothes, G., Schnorpfeil, C., Ackermann, J.-U.:** Production of PHB from crude glycerol, *Eng Life Sci*, **7**, 475-479 (2007).
131. **Koller, M., Bona, R., Braunegg, G., Hermann, C., Horvat, P., Kroutil, M., Martinz, J., Neto, J., Pereira, L., Varila, P.:** Production of polyhydroxyalkanoates from agricultural waste and surplus materials, *Biomacromolecules*, **6**, 561-565 (2005).
132. **Ashby, R. D., Solaiman, D. K. Y., A., F. T.:** Bacterial Poly(hydroxyalkanoate) Polymer Production from the Biodiesel Co-product Stream, *J Polym Environ*, **12**, 105-112 (2004).
133. **Mahishi, L. H., Tripathi, G., Rawal, S. K.:** Poly(3-hydroxybutyrate) (PHB) synthesis by recombinant *Escherichia coli* harbouring *Streptomyces*

- aureofaciens* PHB biosynthesis genes: Effect of various carbon and nitrogen sources, Microbiol Res, **158**, 19-27 (2003).
134. Aldor, I. S., Kim, S. W., Prather, K. L., Keasling, J. D.: Metabolic engineering of a novel propionate-independent pathway for the production of poly(3-hydroxybutyrate-co-3-hydroxyvalerate) in recombinant *Salmonella enterica* serovar *typhimurium*, Appl Environ Microbiol, **68**, 3848-3854 (2002).
135. Bormann, E. J., Roth, M.: The production of polyhydroxybutyrate by *Methylobacterium rhodesianum* and *Ralstonia eutropha* in media containing glycerol and casein hydrolysates, Biotechnol Lett, **21**, 1059-1063 (1999).
136. Sambrook, J., Fritsch, E. F., Maniatis, T.: Molecular Cloning: A Laboratory Manual. Cold Spring Harbor, New York,
137. Ostle, A. G., Holt, J. G.: Nile blue A as a fluorescent stain for poly-beta-hydroxybutyrate, Appl Environ Microbiol, **44**, 238-241 (1982).
138. Hesselmann, R. P., Fleischmann, T., Hany, R., Zehnder, A. J.: Determination of polyhydroxyalkanoates in activated sludge by ion chromatographic and enzymatic methods, J Microbiol Meth, **35**, 111-119 (1999).
139. Watanabe, Y., Ichinomiya, Y., Shimada, D., Saika, A., Abe, H., Taguchi, S., Tsuge, T.: Development and validation of an HPLC-based screening method to acquire polyhydroxyalkanoate synthase mutants with altered substrate specificity, J Biosci Bioeng, **113**, 286-292 (2012).
140. Sangkharak, K., Prasertsan, P.: Utilization of biodiesel waste as a feedstock for the production of polyhydroxybutyrate by *Cupriavidus necator*, Afr J Biotechnol, **10**, 17812-17824 (2011).
141. Sun, Z. Y., Ramsay, J. A., Guay, M., Ramsay, B. A.: Automated feeding strategies for high-cell-density fed-batch cultivation of *Pseudomonas putida* KT2440, Appl Microbiol Biotechnol, **71**, 423-431 (2006).
142. Fukui, T., Yokomizo, S., Kobayashi, G., Doi, Y.: Co-expression of polyhydroxyalkanoate synthase and (R)-enoyl-CoA hydratase genes of *Aeromonas caviae* establishes copolyester biosynthesis pathway in *Escherichia coli*, FEMS Microbiol Lett, **170**, 69-75 (1999).

143. **Sato, S., Ishii, N., Hamada, Y., Abe, H., Tsuge, T.:** Utilization of 2-alkenoic acids for biosynthesis of medium-chain-length polyhydroxyalkanoates in metabolically engineered *Escherichia coli* to construct a novel chemical recycling system, *Polymer Degradation and Stability*, **97**, 329-336 (2012).
144. **Hisano, T., Tsuge, T., Fukui, T., Iwata, T., Miki, K., Doi, Y.:** Crystal structure of the (*R*)-specific enoyl-CoA hydratase from *Aeromonas caviae* involved in polyhydroxyalkanoate biosynthesis, *J Biol Chem*, **278**, 617-624 (2003).
145. **Phithakrotchanakoon, C., Champreda, V., Aiba, S., Pootanakit, K., Tanapongpipat, S.:** Production of polyhydroxyalkanoates from crude glycerol using recombinant *Escherichia coli*, *J Polym Environ*, *in press* (2014).
146. **Shimamura, E., Kasuya, K., Kobayashi, G., Shiotani, T., Shima, Y., Doi, Y.:** Physical properties and biodegradability of microbial poly(3-hydroxybutyrate-*co*-3-hydroxyhexanoate), *Macromolecules*, **27**, 878-880 (1994).
147. **Watanabe, T., He, Y., Fukuchi, T., Inoue, Y.:** Comonomer compositional distribution and thermal characteristics of bacterially synthesized poly(3-hydroxybutyrate-*co*-3-hydroxyhexanoate)s, *Macromol Biosci*, **1**, 75-83 (2001).
148. **Meng, D. C., Shi, Z. Y., Wu, L. P., Zhou, Q., Wu, Q., Chen, J. C., Chen, G. Q.:** Production and characterization of poly(3-hydroxypropionate-*co*-4-hydroxybutyrate) with fully controllable structures by recombinant *Escherichia coli* containing an engineered pathway, *Metab Eng*, **14**, 317-324 (2012).
149. **Wang, Q., Yang, P., Xian, M., Yang, Y., Liu, C., Xue, Y., Zhao, G.:** Biosynthesis of poly(3-hydroxypropionate-*co*-3-hydroxybutyrate) with fully controllable structures from glycerol, *Bioresour Technol*, **142**, 741-744 (2013).
150. **Yamada, M., Matsumoto, K., Shimizu, K., Uramoto, S., Nakai, T., Shozui, F., Taguchi, S.:** Adjustable mutations in lactate (LA)-polymerizing enzyme

- for the microbial production of LA-based polyesters with tailor-made monomer composition, *Biomacromolecules*, **11**, 815-819 (2010).
151. **Sun, Z., Ramsay, J. A., Guay, M., Ramsay, B. A.**: Carbon-limited fed-batch production of medium-chain-length polyhydroxyalkanoates from nonanoic acid by *Pseudomonas putida* KT2440, *Appl Microbiol Biotechnol*, **74**, 69-77 (2007).
152. **Agus, J., Kahar, P., Abe, H., Doi, Y., Tsuge, T.**: Altered expression of polyhydroxyalkanoate synthase gene and its effect on poly[(R)-3-hydroxybutyrate] synthesis in recombinant *Escherichia coli*, *Polym Degrad Stabil*, **91**, 1645-1650 (2006).
153. **San, K. Y., Bennett, G. N., Berrios-Rivera, S. J., Vadali, R. V., Yang, Y. T., Horton, E., Rudolph, F. B., Sariyar, B., Blackwood, K.**: Metabolic engineering through cofactor manipulation and its effects on metabolic flux redistribution in *Escherichia coli*, *Metab Eng*, **4**, 182-192 (2002).
154. **Henderson, R. A., Jones, C. W.**: Poly-3-hydroxybutyrate production by washed cells of *Alcaligenes eutrophus*; purification, characterisation and potential regulatory role of citrate synthase, *Arch Microbiol*, **168**, 486-492 (1997).
155. **Valentin, H. E., Mitsky, T. A., Mahadeo, D. A., Tran, M., Gruys, K. J.**: Application of a propionyl coenzyme A synthetase for poly(3-hydroxypropionate-*co*-3-hydroxybutyrate) accumulation in recombinant *Escherichia coli*, *Appl Environ Microbiol*, **66**, 5253-5258 (2000).
156. **Wang, Q., Yang, P., Liu, C., Xue, Y., Xian, M., Zhao, G.**: Biosynthesis of poly(3-hydroxypropionate) from glycerol by recombinant *Escherichia coli*, *Bioresour Technol*, **131**, 548-551 (2013).
157. **Lu, X., Zhang, J., Wu, Q., Chen, G. Q.**: Enhanced production of poly(3-hydroxybutyrate-*co*-3-hydroxyhexanoate) via manipulating the fatty acid beta-oxidation pathway in *E. coli*, *FEMS Microbiol Lett*, **221**, 97-101 (2003).
158. **Kahar, P., Agus, J., Kikkawa, Y., Taguchi, K., Doi, Y., Takeharu Tsuge, T.**: Effective production and kinetic characterization of ultra-high-molecular-weight poly[(R)-3-hydroxybutyrate] in recombinant *Escherichia coli*, *Polym Degrad Stabil*, **87**, 161-169 (2005).

159. **Desbois, A. P., Smith, V. J.**: Antibacterial free fatty acids: activities, mechanisms of action and biotechnological potential, *Appl Microbiol Biotechnol*, **85**, 1629-1642 (2010).
160. **Sprong, R. C., Hulstein, M. F., Van der Meer, R.**: Bactericidal activities of milk lipids, *Antimicrob Agents Chemother*, **45**, 1298-1301 (2001).
161. **Haywood, G. W., Anderson, A. J., Dawes, E. D.**: The importance of PHB-synthase substrate specificity in polyhydroxyalkanoate synthesis by *Alcaligenes eutrophus*, *FEMS Microbiol Lett*, **57**, 1-6 (1989).
162. **Thompson, J. C., He, B. B.**: Characterization of crude glycerol from biodiesel production from multiple feedstocks, *Appl Eng Agric*, **22**, 261-265 (2006).
163. **Teeka, J., Imai, T., Cheng, X., Reungsang, A., Higuchi, T., Yamamoto, K., Sekine, M.**: Screening of PHA-producing bacteria using biodiesel-derived waste glycerol as a sole carbon source, *J Water Environ Tech*, **8**, 373-381 (2010).
164. **Lee, S. J., Kim, S. B., Kang, S. W., Han, S. O., Park, C., Kim, S. W.**: Effect of crude glycerol-derived inhibitors on ethanol production by *Enterobacter aerogenes*, *Bioprocess Biosyst Eng*, **35**, 85-92 (2012).
165. **Teeka, J., Imai, T., Reungsang, A., Cheng, X., Yuliani, E., Thiantanankul, J., Poomipuk, N., Yamaguchi, J., Jeenanong, A., Higuchi, T. et al**: Characterization of polyhydroxyalkanoates (PHAs) biosynthesis by isolated *Novosphingobium* sp. THA_AIK7 using crude glycerol, *J Ind Microbiol Biotechnol*, **39**, 749-758 (2012).
166. **Nomura, C. T., Tanaka, T., Gan, Z. H., Kuwabara, K., Abe, H., Takase, K., Taguchi, K., Doi, Y.**: Effective enhancement of short-chain-length-medium-chain-length polyhydroxyalkanoate copolymer production by coexpression of genetically engineered 3-ketoacyl-acyl-carrier-protein synthase III (fabH) and polyhydroxyalkanoate synthesis genes, *Biomacromolecules*, **5**, 1457-1464 (2004).
167. **Madden, L. A., Anderson, A. J., Shah, D. T., Asrar, J.**: Chain termination in polyhydroxyalkanoate synthesis: involvement of exogenous hydroxy-compounds as chain transfer agents, *Int J Biol Macromol*, **25**, 43-53 (1999).

- 168.**Noda, I., Green, P. R., Satkowski, M. M., Schechtman, L. A.**: Preparation and properties of a novel class of polyhydroxyalkanoate copolymers, *Biomacromolecules*, **6**, 580-586 (2005).
- 169.**Hablot, E., Bordes, P., Pollet, E., Ave'rous, L.**: Thermal and thermo-mechanical degradation of poly(3-hydroxybutyrate)-based multiphase systems, *Polym Degrad Stabil*, **93**, 413-421 (2008).
- 170.**Kopinke, F. D., Remmler, M., Mackenzie, K.**: Thermal decomposition of biodegradable polyesters—I: Poly(β -hydroxybutyric acid), *Polym Degrad Stabil*, **52**, 25–38 (1996).
- 171.**Foster, L. J. R.**: PEGylation and BioPEGylation of polyhydroxyalkanoates: synthesis, characterisation and applications, *Biopolymers*, 243-256 (2010).
- 172.**Chen, G. Q.**: A microbial polyhydroxyalkanoates (PHA) based bio- and materials industry, *Chem Soc Rev*, **38**, 2434-2446 (2009).

BIOGRAPHY

NAME	Miss Chitwadee Phithakrotchanakoon
DATE OF BIRTH	27 September 1982
PLACE OF BIRTH	Chonburi, Thailand
INSTITUTIONS ATTENDED	Chulalongkorn University, 2001-2004 Bachelor of Science (Biochemistry) (First Class Honors) Mahidol University, 2008-2014 Doctor of Philosophy (Molecular Genetics and Genetic Engineering)
RESEARCH GRANTS	Royal Golden Jubilee Scholarship (PHD/0268/2549)
HOME ADDRESS	316/25 Wachiraprapkarn Rd., Bangplasoi, Muang, Chonburi, Thailand 20000 Tel. +66(38)271758, +66(7)1001686 E-mail: chitwadeeae@gmail.com
PUBLICATION / PRESENTATION	<p>Phithakrotchanakoon C, Champreda V, Aiba S, Pootanakit K, Tanapongpipat S. (2013) Engineered <i>Escherichia coli</i> for Short-Chain-Length Medium-Chain-Length Polyhydroxyalkanoate copolymer biosynthesis from glycerol and dodecanoate. <i>Biosci Biotechnol Biochem.</i> 77(6): 1262-68.</p> <p>Phithakrotchanakoon C, Champreda V, Aiba S, Pootanakit K, Tanapongpipat S. (2014) Production of polyhydroxyalkanoates from crude glycerol</p>

using recombinant *Escherichia coli*. *J*

Polym Environ (in press).

Phithakrotchanakoon C, Champreda V,

Aiba S, Pootanakit K, Tanapongpipat S.

(2013, 5-7 April) Engineered *Escherichia coli* for biosynthesis of

polyhydroxyalkanoates bio-plastic from glycerol. Paper presented at The RGJ-Ph.D.

Congress XIV, Pattaya, Thailand

(outstanding poster presentation award)

Phithakrotchanakoon C, Champreda V,

Aiba S, Pootanakit K, Tanapongpipat S.

(2012, 25-29 Nov) Bioproduction of

polyhydroxyalkanoate bio-plastic from by-product of biodiesel by engineered

Escherichia coli. Paper presented at 13th

FAOBMB congress, Bangkok, Thailand

Phithakrotchanakoon C, Champreda V,

Aiba S, Pootanakit K, Tanapongpipat S.

(2011, 20-21 Dec) Bioproduction of Short-Chain-Length-co-Medium-Chain-Length-

Polyhydroxyalkanoates bioplastic from

crude glycerol by engineered *Escherichia coli*. Paper presented at The third Thai-

Japan bioplastics and biobased materials symposium, Bangkok, Thailand

Phithakrotchanakoon C, Champreda V,

Aiba S, Pootanakit K, Tanapongpipat S.

(2010, 20-22 Oct) Polyhydroxyalkanoate bioplastics production from crude glycerol

by recombinant *Escherichia coli*. Paper

presented at The Thai Society of Biotechnology 2010, Trang, Thailand

Phithakrotchanakoon C, Champreda V,

Aiba S, Pootanakit K, Tanapongpipat S.

(2009, 14 Dec) Polyhydroxyalkanoate bioplastic production from crude glycerol by recombinant *Escherichia coli* using metabolic engineering approach. Paper presented at The young scientist seminar bioresources: their potentials and applications, Bangkok, Thailand

Phithakrotchanakoon C, Champreda V,

Aiba S, Pootanakit K, Tanapongpipat S.

(2009, 2 Sep) Polyhydroxyalkanoate bioplastics production from crude glycerol by recombinant *Escherichia coli* using metabolic engineering approach. Paper presented at The First Thai-Japan bioplastic and biobased material symposium, Bangkok, Thailand