

Effect of Light Intensities and Atmospheric Gas Conditions on Biohydrogen Production of Microalgae Isolated from Fisheries Wastewater

Mujalin Pholchan^{1*}, Kittiwara Kaewseesuk¹, Srikanjana Klayraung¹ and Udomluck Sompong²

¹Program in Environmental Technology, Faculty of Science, Maejo University, Thailand

²Faculty of Fisheries Technology and Aquatic Resources, Faculty of Science, Maejo University, Thailand

ARTICLE INFO

Received: 10 Jan 2017
Received in revised:
22 May 2017
Accepted: 23 May 2017
Published online:
7 Jun 2017

DOI:

Keywords:

Biohydrogen/ Microalgae/
Fishery farm/ *Scenedesmus
accuminatus*/ *Arthrospira
platensis*

* Corresponding author:

E-mail: mujalin@mju.ac.th

ABSTRACT

Recently, the fishery farming industry has been developed rapidly due to increasing demand and consumption as well as the depletion of wild fish resources. Production processes in the industry usually generate large amounts of wastewater containing high nutrients, posing a threat to downstream water. However, phytoplankton removal techniques commonly used to counteract the threat, though appearing to have low efficiency, are timeconsuming and less sustainable. Microalgae are photosynthetic microorganisms that convert solar energy into hydrogen. Using the isolated algae from fish farms as a source of renewable energy production could be a promising choice for handling fisheries wastewater in a more efficient manner. However, hydrogen production processes from algae still need more studies as their efficiencies vary between algae species and growth factors. In this work, the efficiency of hydrogen production from *Scenedesmus accuminatus* and *Arthrospira platensis* harvested from fish farms under three different light intensity conditions and three atmospheric gas conditions was determined. The results showed that the best conditions for hydrogen production from both species included 24-h darkness and carbon dioxide addition. Under the atmospheric gas combination of 99% argon and 1% carbon dioxide, *Scenedesmus accuminatus* could produce hydrogen gas as high as 0.572 $\mu\text{mol H}_2/\text{mgCh h}$ within 12 h, while the highest hydrogen production (0.348 $\mu\text{mol H}_2/\text{mgCh h}$) obtained from *Arthrospira platensis* was found under the atmospheric gas mixture of 98% argon and 2% carbon dioxide. Interestingly, *Scenedesmus accuminatus* appeared to produce more hydrogen than *Arthrospira platensis* under the same conditions.

1. INTRODUCTION

Recently, the fish farming industry has grown rapidly due to population growth and the increasing demand for healthy food choices. The process of raising fish commercially, however, requires large amounts of fresh water, which then causes high amounts of wastewater after the harvesting process, leading to the rapid growth of phytoplankton due to the microscopic dimensions of microalgae (0.5-30 μm). Moreover, phytoplankton removal techniques used commonly used appear to be time consuming and unsustainable while having low efficiency (Uduman et al., 2009; Alfafara et al., 2002).

Therefore, turning what is considered a problem into a raw material for producing a high value-added product such as bioenergy could be a promising approach for effectively handling fisheries wastewater.

Microalgae are photosynthetic microorganisms that naturally convert solar energy into hydrogen via respiration under dark aerobic conditions and via fermentation under anaerobic conditions. Compared with conventional hydrogen production, biological hydrogen production process using fermentative microorganisms (Kosourov et al., 2005; Wang et al., 2003) photosynthetic bacteria, or algae involves less

energy and is considered to be more environmental friendly (Das and Veziroglu, 2001; Levin et al., 2004). Previous reports showed that photosynthetic microorganisms that were cyanobacteria and green algae are of interest in hydrogen production from solar energy and water (Dasgupta et al., 2010; Tamagnini et al., 2007; Maneeruttanarungroj et al., 2010; Eroglu and Melis, 2011).

The hydrogen production processes from algae still require more in-depth studies as their efficiency levels vary between algae species and growth factors. Light is known as the primary energy source for the algae growth and enables it to carry out metabolic processes including hydrogen production (Dubinsky et al., 1995; Markou and Georgakakis, 2011). Different algae species have different light harvesting antenna pigments, light saturation for photosynthesis characteristics, and levels of solar energy conversion efficiency (Eroglu and Melis, 2011). As light intensity increases, chlorophylls and some biomass components such as lipid, fatty acids, and protein contents generally decrease (Tedesco and Duerr, 1928), while starch and polysaccharide contents increase (Friedman et al., 1991). Besides, other cultivation factors such as atmospheric gas composition, carbon source, nutrients, pH, and temperature also have some effects on biohydrogen production.

This work aims to study the effect of two growth factors including light intensity and atmospheric gas condition on the hydrogen production by microalgae harvested from fisheries wastewater. The findings of this study hopefully provide an alternative sustainable choice for fisheries wastewater management by biohydrogen production.

2. METHODOLOGY

2.1 Morphological classification

Morphological classification of algal samples was based on their appearances under a light microscope. Morphologies of cyanobacteria were recorded using photomicrographs. Techniques as described by Desikachary (1959), Anagnostidis and Komarek (1988), Hoffmann (1988) and Hegewald (1990) were used for the classification using an Olympus CH30RF200 compound microscope.

2.2 Sample cultivation and isolation

Water samples were filtered with plankton net (mesh size 5 μm). Pick cell method was used for algal isolation (APHA, 1992). The unialgal culture was examined under a light microscope. The successful cultures were inoculated into flasks containing BG-11 medium (for *Scenedesmus* cultivation) or Zarrouck medium (for *Arthrospira* cultivation). Streak plate dilutions or whole filament isolations were conducted on agar-solidified plates using the same medium for culture purification. All tubes, flasks, and plates were incubated in photochambers at room temperature under fluorescent lamps ($\sim 3,000$ lux) (Castenholz, 1988).

2.3 Strain of microalgae and growth conditions

The dominant species of microalgae from one of the fish farms in Chiang Mai were isolated and identified as described above. Two dominant species, *Scenedesmus accuminatus* (green algae) and *Arthrospira platensis* (cyanobacteria), were selected for the study of H_2 production efficiency. The green algae were cultivated on BG-11 medium, while the cyanobacteria strain was grown on ZN medium under white fluorescence lamps ($80 \mu\text{E}/\text{m}^2 \text{ s}$) at room temperature (Table 1). Cells grown as above were inoculated with an initial optical density of 0.01 and grown for 1 week at room temperature under a light intensity of $80 \mu\text{E}/\text{m}^2 \text{ s}$ until the optical density reached the OD_{730} of 0.672 for *Arthrospira platensis* and the OD_{750} of 0.755 for *Scenedesmus accuminatus*. During cultivation, pH was maintained at 10 and 7 for *Arthrospira platensis* and *Scenedesmus accuminatus*, respectively. Subsequently, cells were harvested by centrifugation at 5000 rpm at 4°C for 15 min. before adding to the photoreactors.

2.4 Photobioreactors setup

To identify the efficiency of both microalgae on H_2 production under different light intensities and atmospheric gas conditions, the batch experiments were conducted in 500 mL gas-tight photoreactors (Figure 1). A 350 mL of cell suspensions was transferred to the photoreactors, which were then tightly capped and flushed with argon gas for 10 min. to eliminate atmospheric oxygen in the headspace. Initial pH for the growth of *Arthrospira platensis* and *Scenedesmus accuminatus* was at 10 and 7, respectively. The

reactors were incubated at room temperature under different treatment conditions, as shown in Table 2, and manually shaken before each sampling. The gas mixture was allowed to pass through 10% NaOH solution for selective absorption of carbon dioxide. Gas samples were collected every 12-h interval

using a sterile glass syringe for composition analysis. Moreover, the algae cells and chlorophyll concentrations were also measured before and at the end of each experiment according to Mackinney (1941). All the experimental sets were carried out as triplicates.

Table 1. Composition of BG-11 medium and ZN₀ medium for *Scenedesmus accuminatus* and *Arthrospira platensis*, respectively

| BG-11 (N-free medium) | | ZN ₀ (N-free medium) | |
|--|--------------|-------------------------------------|--------------|
| Nutrient | Amount (g/L) | Nutrient | Amount (g/L) |
| NaHCO ₃ | 16.80 | H ₃ BO ₃ | 2.850 |
| K ₂ HPO ₄ | 0.50 | MnCl ₂ 4H ₂ O | 1.810 |
| NaNO ₃ | 2.50 | ZnSO ₄ 7H ₂ O | 0.220 |
| K ₂ SO ₄ | 1.00 | CuSO ₄ 5H ₂ O | 0.080 |
| NaCl | 1.00 | MoO ₃ | 0.015 |
| MgSO ₄ 7H ₂ O | 0.20 | | |
| CaCl ₂ 2H ₂ O | 0.04 | | |
| FeSO ₄ 7H ₂ O | 0.20 | | |
| Na ₂ EDTA 2H ₂ O | 1.60 | | |




Figure 1. A schematic of the photoreactors for H₂ production from microalgae

2.5 Measurement of hydrogen production

The composition of gas was analyzed by gas chromatography (Agilent 7890A, USA) with TDC using a molecular sieve 5A, 60/80 column. Helium gas was used as the carrier gas under 3.43 psi and 50 mL/min. of flow rate. In analyzing the gas mixture through chromatography, the moles of H₂ present were calculated taking into account that the total pressure shown on the meter was the sum of the partial pressures contributed by the gas present in the headspace of the photobioreactor. According to Dalton's law of partial pressure, the partial pressure of each gas is presented by the following expression.

$$P_H = X_H \times P,$$

Where P_H is the partial pressure of H₂, X_H is the mole fraction in the gas sample obtained from GC, and P is the total pressure. Then, the number of moles of H₂ could be calculated using the ideal gas equation defined as.

$$P_H V = n_H RT$$

After molar fraction and concentration of hydrogen had been calculated, the hydrogen production rate was obtained and reported as a ratio of μmol of H₂ per mg chlorophyll *a* per h (μmol H₂/mgCh h).

2.6 Chlorophyll a concentration

Once the gas production ceased, biomass samples were taken for determination of chlorophyll *a* concentration. Biomass samples were filtered through a glass fiber filter (Whatman GF/F, retention > 0.7 μm), and pigments were extracted overnight at 4°C with 90% ethanol in the dark. After centrifugation, the absorbance of the supernatant was measured spectrophotometrically (UV-visible spectrophotometer, Spectronic Genesys 5, USA) at 665 nm, before and after acidification with HCl 2 M (Mackinney, 1941). The chlorophyll *a* concentration was calculated as:

$$\text{Chlorophyll a concentration} = 29.6 \times (A-B) \times v/V, \text{ } (\mu\text{g/L})$$

Where A was the absorbance value before acidification, B was the absorbance value after acidification, v was the sample volume (L) and V was the ethanol volume used (mL).

2.7 Statistical analysis

The effects of light intensities and atmospheric gas conditions on hydrogen production from both algae strains were statistically evaluated using a 2-way ANOVA. All tests were considered at a significance level of 0.05.

Table 2. Treatment conditions used in this study

| Light intensities | Atmospheric gas condition |
|---|------------------------------|
| Light condition (80 $\mu\text{E}/\text{m}^2\text{s}$), 24 h | 100% argon |
| | 99% argon + 1% CO_2 |
| | 98% argon + 2% CO_2 |
| Light condition (80 $\mu\text{E}/\text{m}^2\text{s}$), 12 h | 100% argon |
| | 99% argon + 1% CO_2 |
| | 98% argon + 2% CO_2 |
| Dark condition, 12 h | 100% argon |
| | 99% argon + 1% CO_2 |
| | 98% argon + 2% CO_2 |
| Dark condition, 24 h | 100% argon |
| | 99% argon + 1% CO_2 |
| | 98% argon + 2% CO_2 |

3. RESULTS AND DISCUSSION

3.1 Effect of light intensity on H_2 production

In this work, the impact of three different light conditions (24 h of light (80 $\mu\text{E}/\text{m}^2\text{s}$), light (12 h) (80 $\mu\text{E}/\text{m}^2\text{s}$) and dark (12 h) (0 $\mu\text{E}/\text{m}^2\text{s}$), and 24 h of dark (0 $\mu\text{E}/\text{m}^2\text{s}$) on hydrogen production from both algae strains were examined. There is no doubt that solar energy conversion efficiency as well as

hydrogen production are affected by light intensity. Normally, higher light intensity accelerates the growth of biomass and promotes higher hydrogen production (Maneeruttanarungroj et al., 2010). On the other hand, too much light intensity can also suppress hydrogen production due to high oxygen production (Meli et al., 2000). The results revealed that both *Arthrospira platensis* and *Scenedesmus accuminatus* showed greater hydrogen production under 24 h of the dark condition, which was observed from the first 12 h with the value of 0.348 $\mu\text{mol H}_2/\text{mgCh h}$ and 0.572 $\mu\text{mol H}_2/\text{mgCh h}$, respectively (Figure 2(a)). This is in agreement with a previous study by Papazi et al. (2014) demonstrating that the fastest rate and the maximum value of hydrogen production were achieved in the dark conditions. Under 24 h of light conditions, however, there was almost no detectable hydrogen production from either species (data not shown). This probably indicated the result of oxygen produced via photosynthesis that inhibited activities of hydrogenase and nitrogenase (Eroglu and Melis, 2011; Eroglu et al., 2010). The application of green algae for H_2 production must address the effect of oxygen inhibition. In order to prevent the suppression of H_2 production, Hallenbeck and Benemann (2002) recommended that oxygen concentration should be maintained at less than 0.1%.

During the diurnal cycle of 12 h light/ dark condition, however, the result shows that both algae could still produce hydrogen at up to 0.298 $\mu\text{mol H}_2/\text{mgCh h}$ and 0.047 $\mu\text{mol H}_2/\text{mgCh h}$ under the atmospheric gas condition consisting of 98% argon and 2% CO_2 for *Scenedesmus accuminatus* and *Arthrospira platensis*, respectively (Figure 2(b)). This diurnal light-dark cycle has been reported to help increase the yield of hydrogen production due to the improved stability of nitrogenase activity, which creates greater hydrogen production capacity (Meli et al., 2000; Eroglu et al., 2010). Under dark condition, the results (Figure 2(a) and 2(b)) show high levels of hydrogen observed in the treatments where the carbon source (CO_2) was applied to the atmospheric gas, especially the green algae (*Scenedesmus accuminatus*). This may be due to the ability of green algae to produce hydrogen by means of carbon fixation from an external carbon source under dark condition (Chen and Johns, 1996).




Figure 2. Effect of light intensity on hydrogen production of *Arthrospira platensis* and *Scenedesmus accuminatus* (a) under 24 h dark condition and (b) under 12 h light/12 h dark

3.2 Effect of atmospheric gas conditions on H₂ production

Three atmospheric gas conditions used in the study consisted of 1) 100% argon; 2) 99% argon and 1% CO₂; and 3) 98% argon and 2% CO₂. The results show that greater hydrogen production by both algae strains achieved in the treatments used CO₂ added into the atmospheric gas conditions (Figure 3(a), 3(b) and 3(c)). This indicates that the carbon source, one of the photosynthetically metabolic

intermediates, affected the efficiency of hydrogen production. Dutta et al., (2005) suggested that the carbon source affected nitrogenase and reversible hydrogenase activity in the hydrogen production process. Moreover, it was shown that *Arthrospira platensis* achieved the highest hydrogen production (0.348 $\mu\text{mol H}_2/\text{mgCh h}$) in the gas mixture of 98% argon and 2 CO₂ (Figure 3 (c)), while *Scenedesmus accuminatus* tended to produce greater hydrogen (0.572 $\mu\text{mol H}_2/\text{mgCh h}$) in the atmospheric gas

mixture of 99% argon and 1% CO₂ (Figure 3(b)). This is due to the microalgae being photoautotrophic organisms and the carbon source (both inorganic and organic forms) being one of the indispensable nutrients for their cultivation (Vonshak, 1990). Apparently, the carbon source affects their carbon metabolisms (photoautotrophy, heterotrophy, photoheterotrophy and mixotrophy), biomass productivity and biomass composition (Markou and Georgakakis, 2011). In cyanobacteria, there are two different enzymes capable of producing molecular hydrogen, nitrogenase, and bidirectional hydrogenase (Ananyev et al., 2008). It is tempting to speculate that this strain is capable of high levels of net hydrogen production, especially under nitrogen-fixing condition (in the dark) and even in the presence of hydrogenase (in the daytime). As increasing CO₂ is related to photosynthesis, it has been reported that cyanobacteria has the ability to grow and use CO₂ at a higher rate (De Morais and Costa, 2007), meaning that hydrogen production becomes higher where photosynthetic activity is higher. However, it is notable that biomass productivity could be enhanced by the addition of CO₂ but each alga species responds to different CO₂ concentrations (Ho et al., 2011; Florin et al., 2001; Suali and Sarbatly, 2012).

3.3 Effect of Species on H₂ production

To compare the efficiency of hydrogen production between both alga species, the results showed that *Scenedesmus accuminatus* had greater ability to produce higher levels of hydrogen than *Arthrospira platensis* in every condition. At low light intensity conditions, some have reported that *Scenedesmus accuminatus* is capable of both uptake and production of hydrogen after anaerobic adaptation (photoreduction of CO₂) or photohydrogen production (Kosourov et al., 2005; Florin et al., 2001), while *Arthrospira platensis* is an oxygenic photosynthetic bacteria that can only produce hydrogen by reversible hydrogenase through photosynthesis. At the end of experiments, significant pH changes were observed (from 10.0 to 7.5) in the photobioreactor inoculated with *Arthrospira platensis*. On the other hand, the pH value detected from the photobioreactor fed with *Scenedesmus accuminatus* increased slightly from 7.0 to 7.76, which was still in the suitable range for its growth condition. This could lead to lower biomass and hydrogen production as *Arthrospira platensis* is more sensitive to the sudden change in pH due to the formation of hydrogen carbonate. Also, a pH value lower than 9 is not suitable for the growth of *Arthrospira platensis*.




Figure 3. Effect of atmospheric gas conditions on H₂ production: (a) 100% argon; (b) 99% argon + 1% CO₂; and (c) 98% argon + 2% CO₂




Figure 3. Effect of atmospheric gas conditions on H₂ production: (a) 100% argon; (b) 99% argon + 1% CO₂; and (c) 98% argon + 2% CO₂ (cont.)

4. CONCLUSIONS

Light intensity and atmospheric gas condition had some effects on hydrogen production from microalgae. Both species could not produce any hydrogen under 24 h of light. The best condition for hydrogen production from *Scenedesmus accuminatus* was in the atmospheric gas condition with a combination of 99% argon and 1% CO₂ under the 24 h dark condition, while *Arthrospira platensis* achieved the greatest hydrogen production under the atmospheric gas condition of 98% argon and 2% CO₂ in darkness. In comparison, however, it was found that *Scenedesmus accuminatus* could produce more hydrogen gas than *Arthrospira platensis* under the same condition.

ACKNOWLEDGEMENTS

This work was financial supported by National Research Council of Thailand.

REFERENCES

- Alfajara CG, Nakano K, Nomura N, Igarashi T, Matsumura M. Operating and scale-up factors for the electrolytic removal of algae from eutrophied lakewater. *Journal of Chemical Technology and Biotechnology* 2002;77:871-6.
- Anagnostidis K, Komárek J. Modern approach to the classification system of cyanobacteria 3-*Oscillatoriales*. *Algological Studies/Archiv Hydrobiologies* 1988;50:327-472.

- Ananyev G, Carrieri D, Dismukes GC. Optimization of metabolic capacity and flux through environmental cues to maximize hydrogen production by the cyanobacterium "*Arthrospira (spirulina) maxima*". *Applied Environmental Microbiology* 2008;74:6102-13.
- American Public Health Association (APHA). *Standard Methods for Examination of Water and Wastewater*. 18th ed. Washington DC: American Public Health Association; 1992.
- Castenholz RW. Culturing methods for cyanobacteria and thermophilic cyanobacteria: special problems. In: Packer L, Glazer AN. editors. *Methods in Enzymology* Vol. 167. New York: Academic Press Inc; 1988. p. 68-100.
- Chen F, Johns MR. Heterotrophic growth of *Chlamydomonas reinhardtii* on acetate in chemostat culture. *Process Biochemistry* 1996;31:601-4.
- Das D, Veziroglu TN. Hydrogen production by biological processes a survey of literature. *International Journal of Hydrogen Energy* 2001;26:13-28.
- Dasgupta CN, Gilbert J, Lindblad P, Heidorn T, Borgvang SA, Skjanes K. Recent trends on the development of photobiological processes for the improvement of hydrogen production. *International Journal of Hydrogen Energy* 2010;35:10218-38.
- De Moraes MG, Costa JAV. Biofixation of carbon dioxide by *Spirulina* sp. and *Scenedesmus obliquus* cultivated in a three-stage serial tubular photoreactor. *Journal of Biotechnology* 2007;129:439-45.
- Desikachary T. *Cyanophyta*. New Delhi: Indian Council of Agricultural Research; 1959.
- Dubinsky Z, Matsukawa R, Karube I. Photobiological aspects of algal mass culture. *Journal of Marine Biotechnology* 1995;2:61-5.
- Dutta D, De D, Chaudhuri S, Bhattacharya SK. Hydrogen production by cyanobacteria. *Microbial Cell Factories* 2005;4:36-47.
- Eroglu E, Melis A. Photobiological hydrogen production: recent advances and state of the art. *Bioresource technology*. 2011;102:8403-13.
- Eroglu E, Gunduz U, Yucel M, Eroglu I. Photosynthetic bacterial growth and productivity under continuous illumination or diurnal cycles with olive mill wastewater as feedstock. *International Journal of Hydrogen Energy* 2010;35(11):5293-300.
- Florin L, Tsokoglou A, Happe T. A novel type of iron hydrogenase in the green alga *Scenedesmus obliquus* is linked to the photosynthetic electron transport chain. *Journal of Biological Chemistry* 2001;276:6125-32.
- Friedman O, Dubinsky Z, Arad (Malis) S. Effect of light intensity on growth and polysaccharide production in red and blue-green Rhodophyta unicells. *Bioresource Technology* 1991;38:105-10.
- Hallenbeck PC, Benemann JP. *Biological hydrogen; fundamentals and limiting processes*. *International Journal of Hydrogen Energy* 2002;27:1185-93.
- Hegewald E, Hindak F, Schnepf E. *Studies on the Genus Scenedesmus Meyen (Chlorophyceae, Chlorococcales) from South India, with Special Reference to the Cell Wall Ultrastructure*. Berlin: Gebruder Borntraeger; 1990.
- Hoffmann L. Criteria for the classification of blue-green algae (cyanobacteria) at the genus and at the species level *Algological Studies/Archiv Hydrobiologies* 1988;80:131-9.
- Ho S-H, Chen C-Y, Lee D-J, Chang J-S. Research review paper: perspectives on microalgae CO₂ mitigation and biodiesel production. *Bioresource Technology* 2011;101:8725-30.
- Kosourov S, Makarova V, Fedorov AS, Tsygankov A, Seibert M, Ghirardi ML. The effect of sulfur re-addition on H₂ photoproduction by sulfur-deprived green algae. *Photosynthesis Research* 2005;85(3):295-305.
- Levin DB, Pitt L, Love M. Biohydrogen production prospects and limitations to practical application. *International Journal of Hydrogen Energy* 2004;29:173-85.
- Maneeruttanarungroj C, Lindblad P, Incharoensakdi A. A newly isolated green alga, *Tetraspora* sp. CU2551, from Thailand with efficient hydrogen production. *International Journal of Hydrogen Energy* 2010;35:13193-9.
- Mackinney G. Absorption of light by Chlorophyll solutions. *Journal of Biological Chemistry* 1941;140:315-22.
- Markou G, Georgakakis D. Cultivation of filamentous cyanobacteria (blue-green algae) in agro-industrial wastes and wastewater: a review. *Applied Energy* 2011;88:3389-401.
- Meli A, Zhang L, Forestier M, Ghirardi ML, Seibert M. Sustained photobiological hydrogen gas production upon reversible inactivation of oxygen evolution in the green alga *Chlamydomonas reinhardtii*. *Plant Physiology* 2000;122:127-33.
- Papazi A, Gjindali AI, Kastanaki E, Assimakopoulos K, Stamatakis K, Kotzabasis K. Potassium deficiency, a smart cellular switch for sustained high yield hydrogen production by the green alga *Scenedesmus obliquus*. *International Journal of Hydrogen Energy* 2014;39:19452-64.
- Suali E, Sarbatly R. Conversion of microalgae to biofuel. *Renew. Renewable and Sustainable Energy Reviews* 2012;16:4316-42.
- Tamagnini P, Leitao E, Oliveira P, Ferreira D, Pinto F, Harris DJ. *Cyanobacteria hydrogenase: diversity, regulation and applications*. *FEMS Microbiology Review* 2007;31:692-720.

- Tedesco M, Duerr E. Light temperature and nitrogen starvation effects on the total lipid and fatty acid content and composition of *Spirulina platensis* UTEX. *Journal of Applied Phycology* 1928;1:201-9.
- Uduman N, Ying Q, Danquah MK, Forde GM, Hoadley A. Dewatering of microalgae cultures: a major bottleneck to algae-based fuel. *Renewable and Sustainable Energy* 2009;2:1-15.
- Vonshak A. Recent advances in microalgal biotechnology. *Biotechnology Advances* 1990;8:709-27.
- Wang CC, Chang CW, Chu CP, Lee DJ, Chang BV, Liao CS. Producing hydrogen from wastewater sludge by *Clostridium bifermentans*. *Journal of Biotechnology* 2003;102:83-92.