

Figure Content

Figure		Page
2.1	Sangyod rice	5
2.2	Composition of rice bran	7
2.3	Structure of Oryzanol	9
2.4	Structure of tocopherol and tocotrienol	10
2.5	Cold pressing process of rice bran oil in Southern Thailand	13
2.6	Oil in water nanoemulsion	15
4.1	Effect of the stabilization of rice bran by domestic heating on DPPH scavenging effect (%) and inhibition on linoleic acid peroxidation (%) of cold-pressed rice bran oil. The different index letters are statistically significantly different ($P < 0.05$)	40
4.2	Influence of cold-pressed rice bran oil concentration on the particle diameter, mean particle diameter and zeta potential of diluted cold-pressed rice bran oil nanoemulsions stabilized by 3% glyceryl monostearate (GMS)	56
4.3	Influence of glyceryl monostearate (GMS) concentration on the particle diameter (a), mean particle diameter and zeta potential (b) of diluted 30% cold-pressed rice bran oil nanoemulsions	56
4.4	Influence of cold-pressed rice bran oil concentration (a) and glyceryl monostearate concentration (b) on color of cold-pressed rice bran oil nanoemulsions	57

4.5	Influence of cold-pressed rice bran oil concentration (a) and glyceryl monostearate concentration (b) on phytochemical contents and antioxidant activity of cold-pressed rice bran oil nanoemulsions.	60
4.6	Changes of lipid oxidation, phytochemical contents and antioxidant activity of cold-pressed rice bran oil nanoemulsion during storage at 25 °C for 90 days	61
4.7	Influence of pH on physicochemical properties of cold-pressed rice bran oil nanoemulsions	63
4.8	Influence of salt, pH and heat on the creaming index of cold-pressed rice bran oil nanoemulsions	64
4.9	Influence of salt concentration on physicochemical properties of cold-pressed rice bran oil nanoemulsions	66
4.10	Influence of heat on physicochemical properties of cold-pressed rice bran oil nanoemulsions	68