

บรรณานุกรม (Bibliography)

- Balk, M., Weijma, J. and Stams, A.J. (2002). *Thermotoga lettingae* sp. nov., a novel thermophilic, methanol-degrading bacterium isolated from a thermophilic anaerobic reactor. *Int J Syst Evol Microbiol* 52:1361-1368.
- Bhandari, V. and Gupta, R.S. (2014). Molecular signatures for the phylum (class) Thermotogae and a proposal for its division into three orders (Thermotogales, Kosmotogales ord. nov. and Petrotogales ord. nov.) containing four families (Thermotogaceae, Fervidobacteriaceae fam. nov., Kosmotogaceae fam. nov. and Petrotogaceae fam. nov.) and a new genus *Pseudothermotoga* gen. nov. with five new combinations. *Antonie Van Leeuwenhoek* 105: 143 – 168.
- Cedrola, S.M., de Melo, A.C., Mazotto, A.M., Lins, U., Zingali, R.B., Rosado, A.S., Peixoto, R.S. and Vermelho, A.B. (2012). Keratinases and sulfide from *Bacillus subtilis* SLC to recycle feather waste. *World J Microbiol Biotechnol.* 28(3): 1259-1269.
- Cuecas, A., Portillo, M.C., Kanoksilapatham, W. and Gonzalez, J.M. (2014). Bacterial distribution along a 50 °C temperature gradient reveals a parceled out hot spring environment. *Microb Ecol* 68: 729 – 739.
- Friedrich, A.B. and Antranikian, G. (1996). Keratin degradation by *Fervidobacterium pennavorans*, a novel thermophilic anaerobic species of the order Thermotogales. *Appl Environ Microbiol* 62: 2875 – 2882.
- Fardeau, M.L., Ollivier, B., Patel, B.K.C., Magot, M., Thomas, P., Rimbault, A., Rocchiccioli, F., Garcia, J.L. (1997). *Thermotoga hypogea* sp. nov., a xylanolytic, thermophilic bacterium from an oil-producing well. *Int J Syst Bacteriol* 47: 1013-1019.
- Gegeckas, A., Gudiukaitė, R., Debski, J. and Citavicius, D. (2015). Keratinous waste decomposition and peptide production by keratinase from *Geobacillus stearothermophilus* AD-11. *Int J Biol Macromol.* 75: 158-165.

- Godde, C., Sahm, K., Brouns, S.J., Kluskens, L.D., van der Oost, J., de Vos, W.M. and Antranikian, G., (2005). Cloning and expression of islandisin, a new thermostable subtilisin from *Fervidobacterium islandicum*, in *Escherichia coli*, *Appl Environ Microbiol*. 71(7): 3951-3958.
- Huber, R., Langworthy, T.A., König, H., Thomm, M., Woese, .C.R. Sleytr, U.B., Stetter, K.O. (1986). *Thermotoga maritima* sp. nov. represents a new genus of unique extremely thermophilic eubacteria growing up to 90 °C. *Arch Microbiol* 144: 324-333.
- Itoh, T., Onishi, M., Kato, S., Iino, T., Sakamoto, M., Kudo, T., Takashina, T. and Ohkuma, M. (2016). *Athalassotoga saccharophila* gen. nov. sp. nov. isolated from an acidic terrestrial hot spring of Japan, and proposal of *Mesoaciditogales* ord. nov., *Mesoaciditogaceae* fam. nov. in the phylum Thermotogae. *Int J Syst Evol Microbiol* 66: 1045 – 1051.
- Jannasch, H.W., Huber, R., Belkin, S. and Stetter, K.O. (1988). *Thermotoga neapolitana* sp. nov. of the extremely thermophilic, eubacterial genus *Thermotoga*. *Arch Microbiol* 150: 103-104.
- Jeanthon, C., Reysenbach, A.L., L'Haridon, S., Gambacorta, A., Pace, N.R., Glénat, P. and Prieur, D. (1995). *Thermotoga subterranea* sp. nov., a new thermophilic bacterium isolated from a continental oil reservoir. *Arch Microbiol* 164: 91-97.
- Kanoksilapatham, W., Keawram, P. and Gonzalez, J.M. and Robb, F.T. (2015). Isolation, characterization, and survival strategies of *Thermotoga* sp. strain PD524, a hyperthermophile from a hot spring in Northern Thailand. *Extremophiles* 19: 853 – 861.
- Keawram, P. and Kanoksilapatham, W. (2013). Diversity of Hyperthermophilic Bacteria Belonging to Order Thermotogales Thriving in Three Hot Springs in Thailand: Resources of Genes Encoding Thermostable Enzymes. *Silpakorn University Science and Technology Journal*. 7(2): 9-16.
- Keawram, P., Pongsapukdee, V. and Kanoksilapatham, W. (2016). Isolation of *Thermotoga* spp. and *Fervidobacterium* spp., and Characterization of 16S rRNA Genes of Order

- Thermotogales: Unique Lineage of Hyperthermophiles Thriving in 3 Hot Springs in Thailand. *Silpakorn University Science and Technology Journal*. 10(1): 9-20.
- Kim, J.S., Kluskens, L.D. Vos, W.M., Huber, R. and Oost, J.V.D. (2004). Crystal structure of fervidolysin from *Fervidobacterium pennivorans*, a keratinolytic enzyme related to subtilisin. *J. Mol. Biol.* 335: 787-797.
- Kluskens, L.D., Voorhorst, W.G., Siezen, R.J., Schwerdtfeger, R.M., Antranikian, G., van der Oost, J. and de Vos, W.M. (2002). Molecular characterization of fervidolysin, a subtilisin-like serine protease from the thermophilic bacterium *Fervidobacterium pennivorans*. *Extremophiles* 6(3): 185-194.
- Korniłłowicz-Kowalska, T. and Bohacz, J. (2011). Biodegradation of keratin waste: Theory and practical aspects. *Waste Manag.* 31(8): 1689-1701. Review.
- Kublanov, I.V., Tsirol'nikov, K.B., Kaliberda, E.N., Rumsh, L.D., Haertle, T. and Bonch-Osmolovskaia, E.A., (2009). Keratinase of an anaerobic thermophilic bacterium *Thermoanaerobacter* sp. strain 1004-09 isolated from a hot spring in the Baikal Rift zone. *Mikrobiologiya*. 78(1): 79-88.
- Mori, K., Yamazoe, A., Hosoyama, A., Ohji, S., Fujita, N., Ishibashi, J., Kimura, H. and Suzuki, K. (2014). *Thermotoga profunda* sp. nov. and *Thermotoga caldifontis* sp. nov., anaerobic thermophilic bacteria isolated from terrestrial hot springs. *Int J Syst Evol Microbiol.* 64: 2128-2136.
- Nam, G.W., Lee, D.W., Lee, H.S., Lee, N.J., Kim, B.C., Choe, E.A., Hwang, J.K., Suhartono, M.T. and Pyun, Y.R. (2002). Native-feather degradation by *Fervidobacterium islandicum* AW-1, a newly isolated keratinase-producing thermophilic anaerobe. *Arch Microbiol.* 178: 538 – 547.
- Ravot, G., Magot, M., Fardeau, M.L., Patel, B.K.C., Prensier, G., Egan, A., Garcia, J.L. and Ollivier, B. (1995). *Thermotoga elfii* sp. nov., a novel thermophilic bacterium from an African oil-producing well. *Int J Syst Bacteriol.* 45: 308-314.
- Reysenbach, A.L., Liu, Y., Lindgren, A.R., Wagner, I.D., Sislak, C.D., Mets, A. and Schouten, S. (2013). *Mesoaciditoga lauensis* gen. nov., sp. nov., a moderately thermoacidophilic

- member of the order Thermotogales from a deep-sea hydrothermal vent. *Int J Syst Evol Microbiol.* 63: 4724 – 4729.
- Riessen, S. and Antranikian, G. (2001). Isolation of *Thermoanaerobacter keratinophilus* sp. nov., a novel thermophilic, anaerobic bacterium with keratinolytic activity. *Extremophiles.* 5(6) 399 – 408.
- Ryu, S.-I., Kim, J.-E., Huong, N.T., Woo, E.-J., Moon, S.K. and Lee, S.-B. (2010). Molecular cloning and characterization of trehalose synthase from *Thermotoga maritima* DSM3109: Syntheses of trehalose disaccharide analogues and NDP-glucoses. *Enzyme Microb Technol.* 47(6): 249 – 256.
- Siezen, R.J. and Leunissen, J.A. (1997). Subtilases: the superfamily of subtilisin-like serine proteases. *Protein Sci.* 6(3): 501-23. Review.
- Takahata, Y., Nishijima, M., Hoaki, T. and Maruyama, T. (2001). *Thermotoga petrophila* sp. nov. and *Thermotoga naphthophila* sp. nov., two hyperthermophilic bacteria from the Kubiki oil reservoir in Niigata, Japan. *Int J Syst Evol Microbiol.* 51: 1901 – 1909.
- Windberger, E., Huber, R., Trincone, A., Fricke, H. and Stetter, K.O. (1989). *Thermotoga thermarum* sp. nov. and *Thermotoga neapolitana* occurring in African continental solfataric springs. *Arch Microbiol.* 151:506-512.
- Wang, Y., Wang, X., Tang, R., Yu, S., Zheng, B. and Feng, Y. (2001). A novel thermostable cellulase from *Fervidobacterium nodosum*. *J Mol Catal B: Enzym* 66(3-4): 294 – 301.
- Winterhalter, C. and Liebl, W., (1995). Two Extremely Thermostable Xylanases of the Hyperthermophilic Bacterium *Thermotoga maritima* MSB8. *Appl Environ Microbiol* 61(5): 1810–1815.
- Wu, B., Shi, P., Li, J., Wang, Y., Meng, K., Bai, Y., Luo, H., Yang, P., Zhou, Z. and Yao, B. (2010). A new aminopeptidase from the keratin-degrading strain *Streptomyces fradiae* var. k11. *Appl Biochem Biotechnol* 163(3): 730-739.