

Contents

| | Page |
|--|-------------|
| Thai Abstract | i |
| English Abstract | iii |
| Executive Summary | v |
| Contents | x |
| List of Figures | xii |
| List of Tables | xvi |
| Chapter 1 Introduction | |
| 1.1 Overview | 1 |
| 1.2 Objectives | 3 |
| Chapter 2 Model of Turbulent Magnetic Field | |
| 2.1 Two-Component Magnetic Turbulence | 4 |
| 2.1.1 Slab Fluctuation | 5 |
| 2.1.2 Two Dimensional (2D) Fluctuations | 7 |
| 2.2 Turbulent Magnetic Field with Taylor Microscale | 12 |
| Chapter 3 Methodology | |
| 3.1 Generation of magnetic field | 15 |
| 3.1.1 Turbulence Magnetic Field | 15 |
| 3.1.2 Simple Gaussian 2D Field | 17 |
| 3.1.3 Turbulent Magnetic Field with Taylor Microscale | 17 |
| 3.2 Particle Simulations | 19 |
| 3.3 Magnetic Field Line Simulations | 20 |
| 3.4 Simulation and Analysis Method for Separation between Charged Particles and Field Lines | 20 |

Contents

| | Page |
|---|-------------|
| Chapter 4 Effect of Reduced Dimensionality of the Magnetic Field Fluctuations on the Cross-Field Motion of Charged Particles | |
| 4.1 Simulation Setup | 23 |
| 4.2 Results | 24 |
| 4.3 Conclusions and Discussion | 28 |
| Chapter 5 Separation of Charged Particles from Their Turbulent Magnetic Field Lines | |
| 5.1 Separation of Charged Particles and Field Lines in Gaussian 2D Field + Slab Turbulence | 30 |
| 5.2 Separation of Charged Particles and their field lines in 2D+Slab turbulence | 38 |
| Chapter 6 Technique for Measuring and Correcting the Taylor Microscale | |
| 6.1 Correlation Function and Structure Function | 44 |
| 6.2 Extrapolation Method | 48 |
| 6.3 Correction Ratio | 52 |
| 6.4 Applying the Technique to Spacecraft Data | 53 |
| 6.5 Conclusions | 56 |
| References | 57 |
| Output | 60 |
| Appendix | 62 |

List of Figure

| Figure | Page |
|--|------|
| <p>Figure 1.1 Structure of interplanetary magnetic field and solar wind with moving charged particle.</p> | 1 |
| <p>Figure 2.1 Illustration of the slab fluctuation, which depends only on the z coordinate. The arrows demonstrate the slab fluctuation \vec{b}_{slab}.</p> | 5 |
| <p>Figure 2.2 Example of a slab power spectrum.</p> | 6 |
| <p>Figure 2.3 Example of two trajectories of magnetic field lines in pure slab turbulence.</p> | 7 |
| <p>Figure 2.4 Schematic contour plot of $a(x, y)$. The solid arrows show the 2D field, \vec{b}_{2D}, and the dashed arrows show examples of the directions of $\vec{\nabla}a(x, y)$ for both positive and negative potential functions. The 2D field must lie along the equipotential lines of the potential function. For a positive potential function, the 2D field is in a counter-clockwise direction, while a 2D field having a negative potential function is in the clockwise direction.</p> | 8 |
| <p>Figure 2.5 Example of a trajectory of a magnetic field line in pure 2D turbulence.</p> | 9 |
| <p>Figure 2.6 Sample of magnetic field lines in 2D+slab magnetic field turbulence that start at the difference locations. The field lines near O-point are trapped along some distance before they diffuse while the field line near X-point quickly spread at the beginning (Chuychai et al. 2007).</p> | 10 |
| <p>Figure 2.7 Magnetic field line trajectories for a single 2D island, with $\vec{B} = B_0\hat{z} + \vec{b}_{2D}(x, y)$. The surface plot at bottom shows the potential function $a(x, y)$ of the 2D field (Chuychai et al. 2007).</p> | 11 |

List of Figures (Cont.)

| Figure | Page |
|---|------|
| Figure 2.8 Example of two magnetic field lines in a single 2D Gaussian island plus slab turbulence. The red field line started deep inside the island, whereas the blue field line was initially located outside the island. The surface plot at bottom shows the potential function $a(x, y)$ of the 2D field (Chuychai et al. 2007). | 12 |
| Figure 3.1 The power spectrum for a number of values of q in the dissipation range. | 18 |
| Figure 3.2 Illustration of a particle orbit, magnetic field line and guiding center (Wikee 2013). | 21 |
| Figure 3.3 The diagram of separation between the guiding center and magnetic field line (Wikee 2013). | 22 |
| Figure 4.1 The mean squared separation between guiding centers of 100 MeV particles starting at $r_0 = 0.5\lambda$ with their field lines in the pure 2D field. The value close to r_0^2 implies confinement to flux surface. | 25 |
| Figure 4.2 The mean squared separation between guiding centers of 10 MeV particles starting at $r_0 = 0.1\lambda$ with their field lines in the pure 2D field. | 26 |
| Figure 4.3 The mean squared separation between guiding centers of 100 MeV particles and their field lines in pure slab turbulence. The low value indicates confinement of the particle to remain near the same field line. | 27 |
| Figure 4.4 The mean squared separation spreading between guiding centers of 100 MeV particles and their field lines in Gaussian 2D field + slab turbulence. The continual increase indicates unconstrained cross-field motion in this fully three-dimensional case. | 28 |

List of Figures (Cont.)

| Figure | Page |
|---|------|
| Figure 5.1 Example of the trajectory of a charged particle in our model; red line demonstrates trajectory of magnetic field line, black line and blue line demonstrate trajectories of charged particle and its guiding centers, respectively. | 31 |
| Figure 5.2 The results of the separation of charged particles and their corresponding field lines in the log-log scale. | 32 |
| Figure 5.3 The profile of the 2D Gaussian magnetic field along the distance from the center of the flux tube. | 34 |
| Figure 5.4 Showing shape of drift speed of guiding center due to the gradient of the magnetic field, radius of curvature of the magnetic field line and the summation of the gradient drift and curvature drift in arbitrary units. | 35 |
| Figure 5.5 Showing the separation of the charged particles at a) inside the 2D island and b) outside the 2D island. | 36 |
| Figure 5.6 The mean squared perpendicular displacement and time in the final range. | 37 |
| Figure 5.7 Examples of 20 MeV particle trajectories (BLACK lines), their guiding centers (BLUE dots), and their initial field lines (RED lines) in a) pure slab turbulence and b) 10% 2D+ 90% slab turbulence. Note that in our simulations the time scale is in units of λ/c and the length scale is in units of λ , where λ is the slab coherence scale and c is the speed of light. | 38 |
| Figure 5.8 The results from the simulations of 100 MeV protons in a) pure slab and b) 2D+slab turbulence. The statistics in the plots show the spreading of particles from field lines as a function of time. | 39 |
| Figure 5.9 The results of the separation of charged particles and their corresponding field lines when we vary particle energy. | 40 |

List of Figures (Cont.)

| Figure | Page |
|--|------|
| <p>Figure 5.10 The results of the separation of charged particles and their corresponding field lines when we vary initial pitch angles.</p> | 41 |
| <p>Figure 5.11 The results of the separation of charged particles and their corresponding field lines when we vary particle 2D fraction.</p> | 42 |
| <p>Figure 5.12 The results of the separation of charged particles and their corresponding field lines when we vary magnetic field strength.</p> | 42 |
| <p>Figure 6.1 The structure function computed from the time series data for a number of values of q. The bottom curve is associated with a q value of -5 and the higher curves are determined with $q = -4, -3, -2,$ and $-1,$ respectively.</p> | 47 |
| <p>Figure 6.2 The correlation function near the origin. The top curve is determined from $q = -5$ and the next curves are calculated using $q = -4, -3, -2,$ and $-1,$ respectively.</p> | 48 |
| <p>Figure 6.3 Taylor microscale from parabolic fit of the correlation function near the origin for each τ_{fit} for a number of q. Axes are in units of τ_d.</p> | 50 |
| <p>Figure 6.4 Values of τ_{TS} determined by linear extrapolation from the data in Figure 6.3 at the origin (i.e., the intercept). Plotted are the extrapolated Taylor microscale values determined from a range of τ_{max}.</p> | 51 |
| <p>Figure 6.5 Correction ratio versus $1/ q$ for number of different Δt values.</p> | 53 |
| <p>Figure 6.6 The left column shows the plots before applying the correlation ratio to the Taylor microscale λ_T and the right column shows the Taylor microscale (λ'_T) after applying correlation ratio.</p> | 55 |

List of Tables

| Table | Page |
|---|------|
| Table 3.1 Showing index q which we vary for each case and their Taylor scales when we fix dissipation scale ($\tau_d = 2.5$ s). | 19 |
| Table 4.1 Saturation values for 100 MeV particles in the pure 2D field when we vary the location of the initial positions. | 26 |