

## An Upper Cretaceous ichthyodectiform fish from Kyushu, Japan

Yoshitaka Yabumoto<sup>1,\*</sup>, Paulo M. Brito<sup>2</sup> and Koji Hirose<sup>3</sup>

<sup>1</sup>Department of Natural History, Kitakyushu Museum of Natural History and Human History, 2-4-1 Higas-hida, Yahatahigashi-ku, Kitakyushu, Fukuoka 805-0071, Japan.

<sup>2</sup>Departamento de Zoologia, Universidade do Estado do Rio de Janeiro, Rua São Francisco Xavier, Rio de Janeiro, Brazil.

<sup>3</sup>Goshoura Cretaceous Museum, 4310-5 Goshoura, Goshoura, Amakusa, Kumamoto 866-0313, Japan

(Received 11 May 2017; accepted 19 May 2017)

**Abstract** - Ichthyodectiform fishes are an exclusively extinct basal teleost clade, with a temporal range from the Middle Jurassic (Bathonian) to Upper Cretaceous (Maastrichtian). The fish fossils here described were found from Maeshima islet in Kumamoto, Kyushu, Japan in an outcrop of the Hinoshima Formation of the Lower Subgroup of the Himenoura Group. Although the material is constituted by parts of the head, the middle of the body, the caudal skeleton and fin of several individuals, all specimens belong to the same species due to the same features found in the bodies, vertebrae and scales. This species belongs to the order Ichthyodectiformes in having the elongate and laterally compressed body, the posteriorly situated dorsal fin and uroneurals covering the lateral faces of ural centra and the first preuralcentrum.

**Keywords:** Cretaceous, Ichthyodectiformes, Fish, Japan, Goshoura

### 1. Introduction

Ichthyodectiform fishes are an exclusively extinct basal teleost clade, erected by Bardack and Sprinkle (1969), with a temporal range from the Middle Jurassic (Bathonian) to Upper Cretaceous (Maastrichtian) (Arratia, 2004; Cavin *et al.*, 2013). The monophyly of this order was established by Patterson and Rosen (1977) based on two synapomorphies: the presence of an ethmopalatine bone roofing the floor of the nasal capsule (a uniquely ichthyodectiform element in the ethmoid region), and uroneurals covering the lateral sides of the preural centra (Patterson and Rosen, 1977; Stewart, 1999; Leal and Brito, 2004).

Fishes of this order are known from marine and estuarine strata from Australia, North and South America, Africa, Antarctica, Europe and the Middle East (Bardack, 1965; Bardack and Sprinkle, 1969; Schaeffer and Patterson 1984; Stewart 1999; Taverne and Bronzi 1999; Arratia 2000; Taverne and Chanet 2000; Blanco and Cavin 2003; Arratia, 2004; Alvarado-Ortega 2004; Taverne 2008, 2010; Alvarado-Ortega and Brito 2010, Berrell *et al.*, 2014), as well as from freshwater deposits from eastern Asia (Chang, 1963; Liu, 1974; Chang and Chou, 1977; Yabumoto, 1994) and from South America (Brito, 2007; Brito and Yabumoto, 2011).

### 2. Material and method

The fish fossils here described were found at the construction site of the bridge between Goshoura Island and Maeshima islet in Kumamoto, Kyushu, Japan in an outcrop

of the Hinoshima Formation of the Lower Subgroup of the Himenoura Group. The age of the Hinoshima Formation is the Late Cretaceous (Santonian). The specimens are deposited in Goshoura Cretaceous Museum.

Any remains of bones were removed from the specimens with a needle under a microscope. The bone impressions of fossils were then coated with a very thin synthetic resin, surrounded with a low clay wall. Latex was poured onto the fossil impression in a vacuum. The latex was allowed to dry for about an hour and then the latex was peeled off from the fossil. The latex cast was scanned or photographed and drawn on a personal computer and observed under a microscope.

### 3. Systematic description and discussion

Although the material is constituted by parts of the head, the middle of the body and the caudal skeleton and fin of several individuals, all specimens belong to the same species due to the same features found in the bodies, vertebrae and scales. This species belongs to the order Ichthyodectiformes in having the elongate and laterally compressed body, the posteriorly situated dorsal fin and uroneurals covering the lateral faces of ural centra and the first preuralcentrum.

This species resembles *Heckelichthys vexillifer* from Cenomanian of Slovenia and Morocco and *Heckelichthys* sp. from uppermost Cenomanian to Lower Turonian of Vallecillo, north-eastern Mexico (Giersch, 2014) in having a similar preopercle and the short dorsal fin base. How-

\*Author for correspondence: yabumoto@kmmh.jp

ever, this species differs from *H. vexillifer* and *H. sp.* in having a longer snout, the small mouth with small conical teeth and the falcate dorsal fin, not a sail-like. This species is similar to *Heckelichthys microdon* from Turonian of Croatia in having almost the same position of the lower jaw articulation, a similar preopercle, but this species differs from *H. microdon* in having the longer snout, the smaller mouth with small conical teeth and the longer dorsal fin base. The age of this species is slightly younger than those of the species of the genus *Heckelichthys*.

### Acknowledgements

We wish to express our sincere gratitude to Yoshizumi Wakisima for his support for conducting of the excavation and Kazuto Tomiyama for his information about the find of fish fossils. We are also grateful to Hiroaki Ugai and some local people for their help at the excavation. We thank Franz Tatzreiter and Hiromi Kurosu for their preparation of the specimens. This work was supported by JSPS KAK-ENHI Grant Number JP2640506.

### References

- Arratia, G. 2000. Remarkable teleostean fishes from the Late Jurassic of southern Germany and their phylogenetic relationships. *Mitteilungen aus dem Museum für Naturkunde in Berlin, Geowissenschaftliche Reihe* 2, 137–179.
- Arratia, G. 2004. Mesozoic halecostomes and the early radiation of teleosts. In: G. Arratia and A. Tintori (Eds), *Mesozoic Fishes 3 – Systematics, Paleoenvironments and Biodiversity*. Dr Friedrich Pfeil, München, pp. 279–315.
- Alvarado-Ortega, J. 2004. Description and relationships of a new ichthyodectiform fish from the Tlayu´a Formation (Early Cretaceous: Albian), Puebla, Mexico. *Journal of Vertebrate Paleontology* 24, 802–813.
- Alvarado-Ortega, J. and Brito, P. M. 2010. A new ichthyodectiform (Actinopterygii, Teleostei) from the Lower Cretaceous Marizal Formation, north pp. 279–315 east Brazil. *Palaeontology* 53, 297–306.
- Bardack, D. 1965. Anatomy and evolution of chirocentrid fishes. *University of Kansas Paleontological Contributions*, 10, 1–88.
- Bardack, D. and Sprinkle, G. 1969. Morphology and relationships of saurocephalid fishes. *Fieldiana Geology* 16, 297–340.
- Blanco, A. and Cavin, L. 2003. New Teleostei from the Agua Nueva Formation (Turonian), Vallecillo (NE Mexico). *Palevol* 2, 299–306.
- Berrell, W. R., Alvarado-Ortega, J., Yabumoto, Y. and Salisbury, S. W. 2014. The first record of the ichthyodectiform fish *Cladocyclus* from eastern Gondwana: A new species from the Lower Cretaceous of Queensland, Australia. *Acta Palaeontologica Polonica* 59(4), 903–920.
- Brito, P. M. 2007. The Crato Formation fish fauna. In: Martill, D., G. BeChlY, and R. F. loVeridGe (eds.), *The Crato Fossil Beds of Brazil – Window into an Ancient World*. Cambridge University Press, Cambridge, pp. 429–443.
- Brito, P. M. and Yabumoto, Y. 2011. An updated review of the fish faunas from the Crato and Santana formations in Brazil, a close relationship to the Tethys fauna. *Bulletin of Kitakyushu Museum Natural History and Human History, Series A (Natural History)* 9, 107–136.
- Cavin, L., Forey, P. L., and Giersch, S. 2013. Osteology of *Eubiodyctes libanicus* (Pictet and Humbert, 1866) and some other ichthyodectiformes (Teleostei): phylogenetic implications. *Journal of Systematic Palaeontology* 11, 115–177.
- Chang, M. 1963. New materials of *Mesoclupea* from southeastern China and on the systematic position of the genus. *Vertebrata Pal Asiatica*, 7, 105–115 [in Chinese, with English abstract].
- Liu, C. –C. 1974. A new Cretaceous Teleost from Chuh-siung, Yunnan. *Vertebrata PalAsiatica* 12(4), 249–256.
- Leal, M. E. C. and Brito, P. M. 2004. The ichthyodectiform *Cladocyclus gardneri* (Actinopterygii: Teleostei) from the Crato and Santana Formations, Lower Cretaceous of Araripe Basin, North–Eastern Brazil. *Annales de Paléontologie* 90, 103–113.
- Chang, M. –M. and Chou, C. –C. 1977. On late Mesozoic fossil fishes from Zhejiang Province, China. *Acad. Sin., Instit. Palaeont. Palaeoanthro. Mem.* (12), 1–59.
- Giersch, S. 2014. Die Knochenfische der Oberkreidezeit in Nordostmexiko: Beschreibung, Systematik, Vergesellschaftung, Paläobiogeographie und Paläoökologie. PhD. Dissertation, Faculty of Natural Sciences and Mathematics, Heidelberg University, Germany.
- Patterson, C. and Rosen, D.E. 1977. Review of ichthyodectiform and other Mesozoic teleost fishes and the theory and practice of classifying fossils. *American Museum of Natural History, Bulletin* 158, 81–172.
- Schaeffer, B. and Patterson, C. 1984. Jurassic fishes from the Western United States, with comments on Jurassic fish distribution. *American Museum Novitates*, 2796, 1–86.
- Stewart, J. D. 1999. A new genus of Saurodontidae (Teleostei: Ichthyodectiformes) from Upper Cretaceous rocks of the Western Interior of North America. In: Arratia, A. and Schultze, H. –P. (Eds.) *Mesozoic Fishes 2 – Systematics and Fossil Record*. Dr Friedrich Pfeil, München, pp. 335–360.
- Taverne, L. and Bronzi, P. 1999. Les poissons cre´tace´s de Nardo.9°. Note complémentaire sur le Saurodontinae (Teleostei, Ichthyodectiformes): *Saurodon elongatus*, sp. nov. *Studi e Ricerche sui Giacimenti Terziari di Bolca* 8, 105–116.
- Taverne, L. and Chanet, B. 2000. *Faugichthys loryi* n. gen., n. sp. (Teleostei, Ichthyodectiformes) de l'Albien terminal (Cré´tace´ infé´rieur marin) et considé´rations sur la phylogé´nie des Ichthyodectidae. *Geodiversitas*, 22, 23–34.
- Taverne, L. 2008. Considerations about the Late Creta-

ceous genus *Chirocentrites* and erection of the new genus *Heckelichthys* (Teleostei, Ichthyodectiformes) – A new visit inside the ichthyodectid phylogeny. *Bulletin de l'Institut Royal des Sciences naturelles de Belgique* 78, 209–228.

Taverne, L. 2010. Les Ichthyodectidae (Teleostei, Ichthyodectiformes) des schistes bitumineux de l'Aptien (Crétacé inférieur) de Guinée Equatoriale et du Gabon. *Bulletin de l'Institut Royal des Sciences naturelles de Belgique* 80, 115–143.

Yabumoto, Y. 1994. Early Cretaceous Freshwater fish fauna in Kyushu, Japan. *Bulletin of the Kitakyushu Museum of Natural History* 13, 107–254.