

รายงานการวิจัย

เรื่อง

ชื่อโครงการวิจัย

(ภาษาไทย) ผลของรางจืดในการยับยั้งการอักเสบในการติดเชื้อพยาธิใบไม้ตับและกระบวนการก่อมะเร็งท่อน้ำดีในหนูแฮมสเตอร์

(ภาษาอังกฤษ) Inhibitory effect of *Thunbergia laurifolia* Linn in experimental opisthorchiasis and cholangiocarcinogenesis

ชื่อแผนงานวิจัย (ภาษาไทย) การพัฒนาสมุนไพรและแนวทางการรักษาแบบมุ่งเป้าระดับโมเลกุลเพื่อป้องกัน ควบคุม และรักษามะเร็งท่อน้ำดี

(ภาษาอังกฤษ) Herb development and guidance for molecular targeting for prevention, control and treatment of cholangiocarcinoma

1. ผู้รับผิดชอบและหน่วยงาน ประกอบด้วยหน่วยงานหลักและหน่วยงานสนับสนุน

คณะผู้วิจัยและสัดส่วนที่ทำงานวิจัย สัดส่วนที่รับผิดชอบ หน่วยงาน

1. หัวหน้าโครงการ นางสาวพรทิพย์ เหลือมหมื่นไวย 40 % ภาควิชาปรสิตวิทยา คณะแพทยศาสตร์
รับผิดชอบบริหารโครงการ ดำเนินการวิจัย วิเคราะห์ข้อมูล
- 1.2 ผู้ร่วมวิจัย นางสาวธิดารัตน์ บุญมาศ 35 % ภาควิชาปรสิตวิทยา คณะแพทยศาสตร์
รับผิดชอบบริหารโครงการ ดำเนินการวิจัย วิเคราะห์ข้อมูล และให้คำปรึกษา
- 1.3 ผู้ร่วมวิจัย นางสาววัชรินทร์ ลอยลม 15 % ภาควิชาชีวเคมี คณะแพทยศาสตร์
ให้คำปรึกษาด้านเซลล์
- 1.4 ผู้ร่วมวิจัย นางจริยา หาญวงวงศ์ 10 % ภาควิชาจุลชีววิทยา คณะแพทยศาสตร์
ให้คำปรึกษาด้านเซลล์ และสมุนไพร

หน่วยงานหลักที่รับผิดชอบ

ภาควิชาปรสิตวิทยา ชีวเคมี และ จุลชีววิทยา คณะแพทยศาสตร์ มหาวิทยาลัยขอนแก่น 123 ถนนมิตรภาพ อ. เมือง จ.ขอนแก่น 40002 โทรศัพท์ 043-348-387 โทรสาร 043-202-475

2. ประเภทการวิจัย : การวิจัยประยุกต์

3. สาขาวิชาการและกลุ่มวิชาที่ทำการวิจัย: สาขาวิทยาศาสตร์การแพทย์

4. คำสำคัญ (keywords) ของแผนงานวิจัย

Thunbergia laurifolia Linn, hamster, opisthorchiasis, cholangiocarcinoma,

5. Introduction

Thunbergia laurifolia, commonly known as laurel clock vine or blue trumpet vine, is native to India (Starr et al., 2003), and in the Indomalaya ecozone the species occurs from Indochina to Malaysia (Schonenberger, 1999). It is locally known as “kar tuau” in Malaysia and “Rang Chuet” in Thailand (Chan and Lim, 2006). In Malaysia, juice from crushed leaves of *T. laurifolia* are taken for menorrhagia, placed into the ear for deafness, and applied as a poultice for cuts and boils (Burkill, 1966). In Thailand, leaves are used as an antipyretic and for their detoxifying effects, e.g. as an antidote for poisons (Kanchanapoom et al., 2002). Several Thai herbal companies have started producing and exporting Rang Chuet tea (Chan and Lim, 2006). The tea has been claimed to counteract the harmful effects of drugs, alcohol and cigarettes.

Opisthorchis viverrini, a human liver fluke, is the primary risk factor for cholangiocarcinoma (CCA), a rare but highly fatal disease most prevalent in Southeast Asia, including Thailand (IARC, 1994). Humans are infected by ingestion of raw cyprinid fish, which are contaminated by the infective stage known as metacercaria. After *O. viverrini* metacercariae ingestion, the excysted juveniles migrate to the bile canal at the duodenum and grow to adulthood at the common bile duct or gallbladder, an area which is suitable for their survival. At the early stages of infection, liver changes are due to the inflammatory response (eosinophils, monocytes and neutrophils) around the juvenile flukes in the intrahepatic bile ducts. The severity of inflammation gradually increases and reaches a maximum at about 3 to 4 weeks post-infection, as evidenced by the accumulation of mononuclear cells and eosinophils which infiltrate the intrahepatic bile ducts. The virulence of the disease also depends on the number of parasites and the duration of infection, which involves the host's immune response such as cytokine expression and free radicals (Pinlaor et al., 2004). Chronic infection with *O. viverrini* for many years is associated with hepatobiliary diseases (Sripa, 2003), including the development of hepatobiliary cancer and CCA. Reducing the behavior of eating raw fish has proven to be difficult, so the infection rate of this parasite remains high. Therefore, reducing the pathogenesis from an *O. viverrini* infection may be one of the preferred options to reduce the risk of CCA development.

Our previous report (Boonjaraspinyo et al., 2009) showed that a traditional medicine, a mixture of turmeric and fingerroot, reduced the inflammatory cells surrounding the hepatic bile duct but could not be used for prevention of cholangiocarcinogenesis. Thus, the present study was performed to evaluate the efficacy of another Thai medicinal plant, *Thunbergia laurifolia* Linn., or Rang Chuet in Thai. *Thunbergia laurifolia* Linn. is widely used for detoxification from lead poisoning or other toxins (Chattaviriya et al., 2010; Tangpong and Satarug, 2010; Palipoch et al., 2011). Recently, several papers have reported that apigenin, one of the flavonoid compounds in Rang Chuet (Oonsivilai et al., 2007), has antioxidant (Lim and Chan, 2006; Oonsivilai et al., 2008) and anti-cancer properties (Ujiki et al., 2006; Ruela-de-Sousa et al., 2010).

The anti-inflammatory and antioxidant effects of Rang Chuet were studied in a Syrian hamster opisthorchiasis model, and in hamsters with *N*-nitrosodimethylamine administration. Histopathological changes were observed. Liver function tests for alanine transaminase (ALT) and alkaline phosphatase (ALP), and kidney function tests for blood urea nitrogen (BUN) and creatinine, were performed. Total antioxidant capacity was measured by FRAP assay.

6. Materials and methods

6.1. Parasite preparation

Parasites were prepared following the method in a previous report (Wonkchalee et al. 2011). In brief, *Opisthorchis viverrini* metacercariae were obtained from naturally infected cyprinid fish in an endemic area of Khon Kaen, northeast Thailand. Fresh fish were digested in 1% pepsin/HCl and incubated at 37 °C for 1 h, then filtered and precipitated with normal saline in a sedimentation jar. Afterwards the metacercariae – oval-shaped, with large, black excretory bladders – were identified under a dissecting microscope.

6.2. Rang Chuet preparation

Rang Chuet was acquired from a farm in Khon Kaen province, Thailand. Fresh and dried leaves were used; fresh leaves were minced in distilled water, and dried leaves were ground into a powder and

then kept until used. Rang Chuet powder was diluted with distilled water, and 0.5 mL (or 100 mg/kg/d) of the solution was used for oral treatment of the assigned groups.

6.3. FRAP assay

The ferric reducing antioxidant power (FRAP) of extracts was determined following the previous method of Chan and Lim (2006), with modifications. Samples had to be diluted because precipitation occurred upon color development. Dilutions of extracts (1 ml) were added to 2.5 ml phosphate buffer (0.2 M, pH 6.6) and 2.5 ml of potassium ferricyanide (1% w/v). Each mixture was incubated at 50 °C for 20 min. A total of 2.5 ml trichloroacetic acid solution (10% w/v) was added to the mixture to stop the reaction. The mixture was then separated into aliquots of 2.5 ml and diluted with 2.5 ml water. A total of 500 µl ferric chloride solution (0.1% w/v) was added, and the resulting solution was allowed to stand for 30 min until the color was fully developed. Absorbance measured at 700 nm in triplicate was used to normalize with the Fe^{2+} standard curve. Results of the FRAP assay were expressed as mmol Fe^{2+} /g.

6.4. Infection with *Opisthorchis viverrini*

Fifteen Syrian hamsters were each administered 50 *O. viverrini* metacercariae by oral intragastric intubation, as in a previous protocol (Wonkchalee et al., 2011), and then fed with the assigned herb.

6.5. Animal groups

Thirty Syrian hamsters were divided into six groups: 1) infected with *O. viverrini* alone (OV); 2) infected with *O. viverrini* and administered with fresh Rang Chuet (OVFRC); 3) infected with *O. viverrini* and administered with dried Rang Chuet (OVDRC); 4) administered with NDMA alone (NDMA); 5) administered with NDMA and fresh Rang Chuet (NDMAFRC); and 6) administered with NDMA and dried Rang Chuet (NDMADRC). Hamsters were sacrificed on day 30; whole liver tissues were collected for observation of histopathological changes, and sera were tested for liver and kidney function. The protocol was approved by the Animal Ethics Committee of the Faculty of Medicine, Khon Kaen University, Thailand (Ethical Clearance No. AEKKU23/2554).

6.6. *Light microscopic observation*

Hamster livers were fixed in 10% buffered formalin. After fixation, liver tissue was soaked with phosphate-buffered saline and then dehydrated through a series of various concentrations of ethyl alcohol. The tissue was then embedded in paraffin wax. Sections of 5 μ thickness were cut using a microtome, then stained with hematoxylin-eosin and observed under a light microscope, as in previous studies (Boonmars et al., 2009; Boonjaraspinyo et al., 2011). Photographs of each slide were taken at 10x and 40x magnification. The histological feature of liver biopsy and grading criteria levels are shown in Table 1.

6.7. *Biochemical estimation*

6.7.1. *Measurement of serum liver enzymes and kidney function*

Syrian hamster sera were obtained to determine liver and kidney damage; ALT, ALP, BUN and creatinine analysis was performed at the Chemistry Room, Community Laboratory, Faculty of Associated Medical Sciences, Khon Kaen University.

6.7.2. *Statistical analysis*

The data on histopathological changes and serum levels of ALT, ALP, BUN and creatinine were presented as means \pm SD. Statistics were analyzed using one-way ANOVA (SPSS version 13.0, USA). Values were considered statistically significant when $p < 0.05$.

7. **Results**

7.1. *Total antioxidant capacity*

The effect of the temperature of diluents on total antioxidant capacity was evaluated. Fresh and dried Rang Chuet solutions using room temperature water resulted in 10.36 and 95.12 mmol Fe^{2+} /g sample, respectively, while fresh and dried Rang Chuet solutions using boiling water showed 15.72 and 156.29 mmol Fe^{2+} /g sample, respectively. Therefore, boiled dried Rang Chuet solutions were used to treat animal models.

7.2. *Inhibitory effects of Rang Chuet on inflammation*

Observation of the gross pathology of the livers revealed few differences between groups with *O. viverrini* infection (OV), and those treated with Rang Chuet (OVFRC and OVDRC). Liver surfaces were smooth and shiny, with slightly opaque common bile ducts and straw-colored bile fluid. However, a reduction in thickening of the wall of the common bile duct was evident in both OVFRC and OVDRC groups (Fig. 1B, C).

Analysis of histopathological changes focused on the aggregation of inflammatory cells surrounding the hepatic bile ducts and liver tissue, and hepatic bile duct proliferation. Histopathological changes in the *O. viverrini*-infected groups (Fig. 1A–F) were similar to those observed in the NDMA groups (Fig. 1G–L). Aggregation of inflammatory cells surrounding the hepatic bile ducts was observed both in *O. viverrini*-infected groups (Fig. 1A, D) and in those receiving NDMA administration (Fig. 1G, J). A reduction of inflammatory cells surrounding the intrahepatic bile ducts was observed in the OVFRC, OVDRC, NDMAFRC and NDMAFRC groups (Fig. 1E, F, K, L). The degree of inflammatory aggregation is shown in Table 1.




Fig. 1 Gross and histopathological changes in the group of *O. viverrini* infection (OV; a, d), *O. viverrini* infection with fresh Rang Chuet solutions (OVFRC; b, e), *O. viverrini* infection with dried Rang Chuet solutions (OVDRC; c, f) and the group of administration NDMA (NDMA; g, j), administration NDMA with fresh Rang Chuet solutions (NDMAFRC; h, k), administration NDMA with dried Rang Chuet solutions (OVDRC; i, l). GB gall bladder, CBC common bile duct, P parasite. Asterisk indicates inflammation. Magnification×10

Table 1 Histological feature of liver biopsy and grading criteria levels in the group of OV infection and administration NDMA with or without fresh Rang Chuet solutions (FRC) and with or without dried Rang Chuet solutions (DRC) by microscopic observation

Histopathology	Criteria	OV	OVFRC	OVDRC	NDMA	NDMAFRC	NDMADRC
Goblet cell proliferation	0	0	0	0	0	0	0
	1	0	0	0	0	0	0
	2	0	0	0	0	0	0
	3	0	0	0	0	0	0
Portal inflammation	0	0	0	0	0	0	0
	1	0	25% (1)	40% (2)	33.4% (2)	50% (2)	60% (3)
	2	50% (3)	25% (1)	40% (2)	66.6% (4)	50% (2)	40% (2)
	3	50% (3)	50% (2)	20% (1)	0	0	0
Focal inflammation	0	0	0	0	0	0	0
	1	0	0	0	33.4% (2)	50% (2)	60% (3)
	2	33.4% (2)	50% (2)	60% (3)	66.6% (4)	50% (2)	40% (2)
	3	66.6% (4)	50% (2)	40% (2)	0	0	0
	4	0	0	0	0	0	0
Hepatic bile duct proliferation	0	0	0	0	100% (6)	100% (4)	100% (5)
	1	33.4% (2)	25% (1)	50% (2)	0	0	0
	2	66.4% (4)	75% (3)	50% (2)	0	0	0
	3	0	0	0	0	0	0

Goblet cell proliferation: [0=absent, 1=single or discrete group, 2=two or more group, 3=diffuse], portal inflammation: [0=minimal/no portal inflammation, 1=mild (sparking of inflammatory cells <1/3 of portal tract, 2=moderate (increase in inflammatory cells 1/3–2/3 of portal), 3=severe (dense packing >2/3 of portal tract)], focal inflammation: [0=none, 1=one focus or less per 10× objective, 2=two to four foci per 10× objective, 3=four to ten foci per 10× objective, 4=more than ten foci per 10× objective], hepatic bile duct proliferation: [0=absent, 1=focal, 2=scattering, 3=diffuse]

Table 2 Blood chemistry test tests serum levels of alanine transaminase (ALT), alkaline phosphatase (ALP), blood urea nitrogen (BUN), creatinine (Cr), levels in the group of *O. viverrini* infection alone (OV), *O. viverrini* infection with fresh Rang Chuet solutions (OVFRC), *O. viverrini* infection with dried Rang Chuet solutions (OVDRC) and administration NDMA alone (NDMA), administration NDMA with fresh Rang Chuet solutions (NDMAFRC), administration NDMA with dried Rang Chuet solutions (NDMADRC) compared normal control

Group	BUN (mg/dl) Mean±SEM	Creatinine (g/dl) Mean±SEM	ALT (U/l) Mean±SEM	ALP (U/l) Mean±SEM
Normal	22.0±4 ^a	0.3±0.1 ^a	90±15 ^a	172.5±47.5 ^a
OV	26.0±1.14 ^a	0.24±0.02 ^a	1,121.1±88.9 ^{c,d}	139.4±4.79 ^a
OVFRC	21.67±0.88 ^a	0.26±0.03 ^a	605.33±44.86 ^{b,c}	204.67±37.7 ^b
OVDRC	22.4±0.81 ^a	0.38±0.02 ^b	853.75±56.73 ^b	130.25±116.85 ^{a,b}
NDMA	18.6±0.6 ^b	0.22±0.02 ^a	1,238±154 ^d	99.20±3.97 ^a
NDMAFRC	17.4±0.68 ^b	0.26±0.02 ^a	1,016±77.57 ^c	142±18.03 ^a
NDMADRC	22.4±0.51 ^a	0.3±0.0 ^a	258.8±52.81 ^a	137.8±9.29 ^a

Mean with different superscripts are significantly different ($P < 0.05$)

7.3. Effects of Rang Chuet on liver enzyme and kidney function

Table 1 shows the activities of serum ALT, ALP, BUN and creatinine, which correlate with histopathological changes (Fig. 1). The liver and kidney serum markers (ALT, ALP, BUN and creatinine) in the Rang Chuet group remained within normal levels. Serum ALT levels increased about 3- to 10-fold after Syrian hamsters were administered with NDMA or infected with *O. viverrini*. There was a significant decrease in serum ALT in the groups of OVFRC ($p = 0.012$), OVDRC ($p = 0.001$) and NDMAFRC ($p = 0.000$) at 1 month. Serum BUN, ALP and creatinine levels in the Rang Chuet, NDMA, NDMADRC, DMAFRC, OV, OVFRC and OVDRC groups remained within normal levels.

8. Discussion

The present study clearly shows that Rang Chuet has anti-inflammatory and antioxidant properties which improve liver function in hamsters with liver fluke infection or after administration of NDMA. Pathological changes, as evidenced by a reduction of the inflammatory cells surrounding the hepatic bile ducts, were correlated with the results of the liver function tests (especially ALT) showing decreased liver cell damage. Higher antioxidant activity, as shown by FRAP analysis, was due to the reduction of inflammatory cells surrounding the hepatic bile duct and to the reduction of ALT levels.

Examination of pathological changes in the livers revealed the anti-inflammatory property of Rang Chuet in Syrian hamsters with inflammatory cells induced by *O. viverrini* infection or NDMA administration. This was in agreement with a previous report in which Rang Chuet administered at 5 g/kg to mice with induced paw edema demonstrated anti-inflammatory activity at 3 h ($p > 0.01$) and 6 h ($p > 0.001$) (Pongphasuk et al., 2005). Histopathological results, supported by liver function tests, showed that Rang Chuet has no toxic side effects; this was in line with previous reports where Rang Chuet

demonstrated no toxic effects in a mouse model at 8 g/kg/d for 30 d (Pongphasuk et al., 2005), and in a rat model at up to 2 g/kg/d for 6 months (Chivapat et al., 2009). The administration of NDMA was directly toxic to liver tissue, and induced the infiltration of inflammatory cells surrounding the biliary tree and hepatic tissue by day 30, similar to the findings of Boonmars et al. (2009). Moreover, NDMA administration in rats induced chronic inflammation and led to liver tumors (Peto et al., 1991), bile duct proliferation, and liver fibrosis (George et al., 2001). The anti-inflammatory property of Rang Chuet was clearly demonstrated by the reduction of inflammatory cells in hepatic tissue, leading to decreased serum ALT (Table 1) in all groups treated with a Rang Chuet solution. This result agrees with a previous study in which the extract compounds from Rang Chuet showed anti-inflammatory effects, as well as antioxidant effects as determined by FRAP assay (Oonsivilai et al., 2007). In addition, phenolic profiling of Rang Chuet water extract revealed the presence of apigenin and apigenin glucosides as well as phenolic acids such as caffeic acid, gallic acid and protocatechuic acid (Oonsivilai et al., 2007); these compounds are involved in its anti-inflammatory and anti-cancer properties. About eight steroid substances have also reportedly been found by thin layer chromatographic study of the extracts.

The histopathology of Syrian hamsters infected with *O. viverrini* was similar to previous reports (Boonmars et al., 2007; 2008; 2009). At 30 days post-infection, a peak of inflammatory cells (mononuclear cells and eosinophils) was observed surrounding the hepatic bile ducts, as well as epithelial hyperplasia, goblet cell metaplasia, adenomatous metaplasia, and thickened periductal fibrosis (Fig. 1), findings which correspond with increased serum ALT level. A slight decrease in inflammatory cells was observed, both in the case of *O. viverrini* infection and NDMA administration, when a Rang Chuet diet was administered. Although the results were similar for all Rang Chuet-treated groups, different inflammatory inducers may trigger different host immune response mechanisms. The metabolized product from NDMA is directly toxic to the liver, and subsequently generates an inflammatory cell response; whereas *O. viverrini* induces a host immune response, as evidenced by inflammatory cells surrounding the hepatic bile duct. Moreover, the present study shows that the antioxidant property was in agreement with previous reports, which found that Rang Chuet powder diluted with boiling water had a high antioxidant capacity with low toxicity (Wong et al., 2006; Oonsivilai et al., 2008).

The present study demonstrates the advantages of Rang Chuet as an antioxidant and anti-inflammatory in Syrian hamsters that were administered NDMA, as well as in those infected with *O. viverrini*, leading to a reduction in liver pathology and an improvement in liver function.

Acknowledgements

This work was granted by the Thailand research fund through RMU5480002 Khon Kaen University (KKU) and the Royal Golden Jubilee-Ph.D. program (Grant No. PHD/0280/2552) to Miss. Orasa Wonkchalee and Associate Professor Thidarut Boonmars. We thank the Department of Parasitology, Liver Fluke and Cholangiocarcinoma Research Center, the Animal Experimental Unit, Faculty of Medicine, Khon Kaen University for their support.

9. References

- Boonjaraspinyo S, Boonmars T, Aromdee C, Srisawangwong T, Kaewsamut B, Pinlaor S, Yongvanit P, Puapairoj A (2009) Turmeric reduces inflammatory cells in hamster opisthorchiasis. *Parasitol Res* 105:1459-1463.
- Boonmars T, Boonjaraspinyo S, Kaewsamut B (2009) Animal models for *Opisthorchis viverrini* infection *Parasitol Res* 104:701-703.
- Boonmars T, Srirach P, Kaewsamut B, Srisawangwong T, Pinlaor S, Pinlaor P, Yongvanit P, Sithithaworn P (2008) Apoptosis-related gene expression in hamster opisthorchiasis post praziquantel treatment. *Parasitol Res* 102:447-455.
- Boonmars T, Srisawangwong T, Srirach P, Kaewsamut B, Pinlaor S, Sithithaworn P (2007) Apoptosis-related gene expressions in hamsters re-infected with *Opisthorchis viverrini* and re-treated with praziquantel. *Parasitol Res* 102:57-62.
- Chandra K, Graham PJ, Ian RR, Graeme HM (2007) Anticancer properties of panduratin A isolated from *Boesenbergia pandurata* (Zingiberaceae). *J Nat Med* 61:131-137.
- Cuendet M, Oteham CP, Moon RC, Pezzuto JM (2006) Quinone reductase induction as a biomarker for cancer chemoprevention. *J Nat Prod* 69:460-463.

- Fahey JW, Stephenson KK (2002) Pinostrobin from honey and Thai ginger (*Boesenbergia pandurata*): a potent flavonoid inducer of mammalian phase 2 chemoprotective and antioxidant enzymes. *J Agric Food Chem* 50:7472-7476.
- George J, Rao KR, Stern R, Chandrakasan G (2001) Dimethylnitrosamine-induced liver injury in rats: the early deposition of collagen. *Toxicology* 156:129-138.
- IARC (1994) Infection with liver flukes (*Opisthorchis viverrini*, *Opisthorchis felinus* and *Clonorchis sinensis*). IARC monographs on the evaluation of carcinogenic risks to humans 61:121-175.
- Mahidol C, Tuntiwachwuttikul P, Reutrakul V, Taylor WC (1984) Constituents of *Boesenbergia pandurata* (syn. *Kaempferia pandurata*) III Isolation and synthesis of (\pm)-boesenbergin B. *Aust J Chem* 37:1739-1745.
- Murakami A, Jiwajinda S, Koshimizu K, Ohigashi H (1995) Screening for in vitro anti-tumor promoting activities of edible plants from Thailand. *Cancer Lett* 95:139-146.
- Pongphasuk N, Khunkitti W, Chitcharoenthum M. III WOCMAP Congress on Medicinal and Aromatic Plants - Volume 6: Traditional Medicine and Nutraceuticals. Palaniswamy UR, Gardner ZE, Craker LE eds (2005) ISHS Acta Horticulturae 680
- Peto R, Gray R, Brantom P, Grasso P (1991) Dose and time relationships for tumor induction in the liver and esophagus of 4080 inbred rats by chronic ingestion of N-nitrosodiethylamine or N-nitrosodimethylamine. *Cancer Res* 51:6452-6469.
- Pinlaor S, Hiraku Y, Ma N, Yongyanit P, Semba R, Oikawa S, Murata M, Sripa B, Sithithaworn P, Kawanishi S (2004) Mechanism of NO-mediated oxidative and nitrative DNA damage in hamsters infected with *Opisthorchis viverrini*: a model of inflammation-mediated carcinogenesis. *Nitric Oxide* 11:175-183.
- Shindo K, Kato M, Kinoshita A, Kobayashi A, Koike Y (2006) Analysis of antioxidant activities contained in the *Boesenbergia pandurata* Schult. Rhizome. *Biosci Biotechnol Biochem* 70:2281-2284.
- Sripa B (2003) Pathobiology of opisthorchiasis: an update. *Acta Tropica* 88, 209-220.

Tuchinda P, Reutrakul V, Claeson P, Pongprayoon U, Sematong T, Santisuk T, Taylor WC (2002) Anti-inflammatory cyclohexenyl chalcone derivatives in *Boesenbergia pandurata*. *Phytochemistry* 59:169-173.

Yanti, Anggakusuma, Gwon SH, Hwang JK (2009) *Kaempferia pandurata* Roxb. inhibits *Porphyromonas gingivalis* supernatant-induced matrix metalloproteinase-9 expression via signal transduction in human oral epidermoid cells. *J Ethnopharmacol* 123:315-324.

10. ผลลัพธ์ที่ได้จากโครงการวิจัย

1. ผลิตนักศึกษาระดับปริญญาเอกจำนวน 1 คน
2. ตีพิมพ์วารสารนานาชาติจำนวน 1 เรื่อง ดังเอกสารแนบ
 - 2.1 Anti-inflammatory, antioxidant and hepatoprotective effects of *Thunbergia laurifolia* Linn. on experimental opisthorchiasis. Wonkchalee O, Boonmars T, Aromdee C, Laummaunwai P, Khunkitti W, Vaeteewoottacharn K, Sriraj P, Aukkanimart R, Loilome W, Chamgramol Y, Pairojkul C, Wu Z, Juasook A, Sudsarn P. *Parasitol Res.* 2012 Jul;111(1):353-9.
3. นำเสนอผลงานวิจัยในการประชุมระดับชาติ 1 ครั้ง และนานาชาติ 1 ครั้ง ดังเอกสารแนบ

ภาคผนวก


Anti-inflammatory, antioxidant and hepatoprotective effects of Thunbergia laurifolia Linn. on experimental opisthorchiasis

Orasa Wonkchalee, Thidarut Boonmars, Chantana Aromdee, Porntip Laummaunwai, Watcharee Khunkitti, Kulthida Vaeteewoottacharn, Pranee

Parasitology Research
Founded as Zeitschrift für
Parasitenkunde

ISSN 0932-0113

Parasitol Res
DOI 10.1007/s00436-012-2846-5



 Springer

Anti-inflammatory, antioxidant and hepatoprotective effects of *Thunbergia laurifolia* Linn. on experimental opisthorchiasis

Orasa Wonkchalee · Thidarut Boonmars ·
Chantana Aromdee · Porntip Laummaunwai ·
Watcharee Khunkitti · Kulthida Vaeteewoottacharn ·
Pranee Sriraj · Ratchadawan Aukkanimart ·
Watchalin Loilome · Yaovaluk Chamgramol ·
Chawalit Pairojkul · Zhiliang Wu · Amornrat Juasook ·
Pakkayane Sudsarn

Received: 19 January 2012 / Accepted: 24 January 2012
© Springer-Verlag 2012

Abstract *Thunbergia laurifolia* Linn (Rang Chuet) possesses antioxidant and anti-inflammatory properties as well as anticancer activities. The aim of the present study was to evaluate the efficacy of *T. laurifolia* in reducing inflammation

from pathological changes in Syrian hamsters infected with the human liver fluke *Opisthorchis viverrini*. Hamster groups were also administered *N*-nitrosodimethylamine (NDMA) and treated with *T. laurifolia*. Light microscopic observation of histopathological changes, liver function tests for alanine transaminase (ALT) and alkaline phosphatase (ALP) and kidney function tests for blood urea nitrogen (BUN) and creatinine were performed. Antioxidant effects of both fresh and dried Rang Chuet solutions were observed. Analysis of the histopathological changes showed anti-inflammatory properties, both in the case of *O. viverrini* infection or with NDMA

O. Wonkchalee · T. Boonmars (✉) · P. Laummaunwai · P. Sriraj ·
R. Aukkanimart · A. Juasook · P. Sudsarn
Department of Parasitology, Faculty of Medicine,
Khon Kaen University,
Khon Kaen 40002, Thailand
e-mail: bthida@kku.ac.th



A combination of praziquantel and traditional plant, *Thunbergia laurifolia* Linn, improve hepatobiliary system in Syrian hamster opisthorchiasis and cholangiocarcinoma

Wichitarn Worachalee^{1,2,3}, Pongp Lammunwai¹, Thidarat Boonmorn^{1,2}, Chantana Arandee², Chanya Samnangwong¹, Kulnida Veerawuthacham^{2,4}, Wu Zhiqiang¹, Preeee Sitr^{1,2}, Ratchadawan Aukkanim^{1,2}, Ratchan Lakshe^{1,2}, Yanyan Chongrattana^{2,4}, Chawalit Pankaj^{2,5}, Anomrat Jussak^{1,2} and Pailaywanee Subhan^{1,2}

Department of Parasitology, Faculty of Medicine, Khon Kaen University, Khon Kaen, 40002, Thailand; ²Liver and Cholangiocarcinoma Research Center, Khon Kaen University, Khon Kaen, 40002, Thailand; ³The Royal Golden Jubilee Ph.D Programme of Thailand Research Fund (RGJ-TRF); Faculty of Pharmaceutical Science, Khon Kaen University, Khon Kaen, 40002, Thailand; ⁴Department of Microbiology; ⁵Department of Biochemistry, Faculty of Medicine, Khon Kaen University, Khon Kaen, 40002, Thailand; ⁶Department of Parasitology, Oita University Medicine, Graduate School, Gifu 5011794, Japan; ⁷Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen, 40002, Thailand

Email: goret@gmail.com

ABSTRACT

Praziquantel is the drug of choice for liver fluke infections, during which the parasites release a large amount of antigens, leading to an accumulation of inflammatory cells surrounding the hepato bile ducts which causes cell damage. *Thunbergia laurifolia* Linn. is a traditional plant which has antioxidant and anti-inflammatory properties in vitro and in previous studies on hamster opisthorchiasis. Therefore, in the present study we demonstrated the combined effect of *Thunbergia laurifolia* Linn. (RC) plus praziquantel (PZ) on hamster opisthorchiasis and cholangiocarcinoma (CCA). Hamsters were divided into six groups: i) infected with *Opisthorchis viverrini* plus praziquantel (OV_PZ); ii) infected with *O. viverrini* plus PZ and RC (OV_PZ_RC); iii) infected with *O. viverrini* plus N-methyl-N-nitrosodimethylamine (OV_NDMA); iv) infected with *O. viverrini* plus NDMA and RC (OV_NDMA_RC); v) infected with *O. viverrini* plus NDMA and PZ (OV_NDMA_PZ); and vi) infected with *O. viverrini* plus NDMA, PZ and RC (OV_NDMA_PZ_RC). Light microscopic observation of histopathological changes, liver function tests for alanine transaminase (ALT) and alkaline phosphatase (ALP), and kidney function tests for blood urea nitrogen (BUN) and creatinine were performed. The study found that RC extracts clearly improve the hepatobiliary system by reducing inflammatory cell aggregation and inhibiting CCA development, results which were correlated with serum ALT and ALP. These findings suggest that administration of *Thunbergia laurifolia* Linn. after praziquantel treatment can improve the hepatobiliary system and could reduce the risk of CCA development.

supported by The Royal Golden Jubilee Ph.D Programme and PHD 0260/2552, RMU5480002 and TRF Senior Research Scholar Grant no. RTA5580004





บทคัดย่อ

Pre-Congress Symposium 2012

"การวิจัยและพัฒนาเพื่อกำจัดพยาธิใบไม้ตับและมะเร็งท่อน้ำดี"

(Research and Development for eliminating the liver fluke and cholangiocarcinoma)

วันที่ 9 ตุลาคม 2555

ณ ห้องบรรยาย 2 คณะแพทยศาสตร์ มหาวิทยาลัยขอนแก่น

ศูนย์วิจัยพยาธิใบไม้ตับและมะเร็งท่อน้ำดี และภาควิชาชีวเคมี คณะแพทยศาสตร์

มหาวิทยาลัยขอนแก่น

LIVER FLUKE AND CHOLANGIOCARCINOMA RESEARCH CENTER
KHON KAEN UNIVERSITY

Effects of *Thunbergia laurifolia* Linn. on experimental opisthorchiasis

Nadchanan Workhawe^{1,2}, Thidarat Boonmars^{1,2}, Chantana Anomdee³, Pomtip Laummaurwai¹,
Welcharee Khunkit², Kulthida Vateewootacham⁴, Pranee Siriraj^{1,2}, Ratchadawan Aulikanimart^{1,2},
Witcherin Lolkime⁴, Yaovasilk Chamgramol⁵, Chawalit Petrokijul⁶, Zhikang Wu⁴, Amornrat Juasook^{1,2},
Pakkayamee Suddsam^{1,2}

¹Department of Parasitology, Faculty of Medicine, Khon Kaen University, ²Liver Fluke and Cholangiocarcinoma Research Center, ³Faculty of Pharmacology, ⁴Department of Biochemistry, Faculty of Medicine, Khon Kaen University, Thailand, ⁵Department of Parasitology, Gifu University Medicine Graduate School, Japan, ⁶Department of Pathology, Faculty of Medicine, Khon Kaen University, Thailand
E-mail: jradcha@gmail.com

Thunbergia laurifolia Linn. is locally known as Rang Chuet in Thailand. Rang Chuet properties are anti-oxidant and anti-inflammation including anti-cancer activities. Therefore, the present study is revealing the efficacy of *Thunbergia laurifolia* Linn. on reducing the inflammation through pathological changes. Syrian hamsters were divided into 6 groups, i) *Opisthorchis viverrini* infection(OV), ii) *N*-nitrosodimethylamine administration (NDMA), iii) OV plus fresh Rang Chuet, iv) NDMA plus fresh Rang Chuet, v) OV plus dried Rang Chuet vi) NDMA plus dried Rang Chuet. Gross and histopathology were observed under digital camera and light microscope respectively. Liver function tests for alanine transaminase (ALT) and alkaline phosphatase (ALP) and kidney function tests (blood urea nitrogen (BUN) and creatinine) were investigated. Results showed that both fresh and dried Rang Chuet had the antioxidant capacity correlated to histopathological changes which showed the anti-inflammatory property in the case of OV plus Rang Chuet and NDMA plus Rang Chuet by reducing the aggregation of inflammatory cells surrounding the hepatic bile ducts, correlates to serum ALT, ALP, BUN and creatinine levels in Syrian hamsters treated groups. The present study found that fresh and dried Rang Chuet is clearly reduces the inflammatory cells in both treated groups, infected and administered NDMA groups and correlated with the total anti-oxidant capacity. This finding suggests that *Thunbergia laurifolia* Linn has anti-oxidant and anti-inflammatory properties which may useful for application of *Thunbergia laurifolia* Linn on prevention of inflammatory process which is the risk factors of *O. viverrini* associated - cholangiocarcinoma.

