

## เอกสารอ้างอิง

1. Vatanasapt V, Sriamporn S, Vatanasapt P. Cancer control in Thailand. *Jpn J Clin Oncol* 2002;32(Suppl):S82-91.
2. Wanrum S, Limpaiboom T, Leelayuwat C, Yaenyao P, Laranratanakarn S, Jearanaikoon P. Development of method for integrated HPV16 detection in cervical carcinomas using multiplex PCR. *J Med Tech Phy Ther* 2007; 19: 6-15.
3. Moretta A, Biassoni R, Bottino C, Mingari MC, Moretta L. Natural cytotoxicity receptors that trigger human NK-cell-mediated cytotoxicity. *Immunology Today* 2000;21:228-34.
4. Garrido F, Cabrera T, Concha A, Glew S, Ruiz-Cabello F, et al. Natural history of HLA expression during tumour development. *Immunology Today* 1993;14:491-9.
5. Garrido F, Ruiz-Cabello F, Cabrera T, Perez-Villar JJ, Lopez-Botet M, Duggan-Keen M, et al. Implications for immunosurveillance of altered HLA class I phenotypes in human tumours. *Immunology Today* 1997;18:89-95.
6. Groh V, Rhinehart R, Randolph-Habecker J, Topp MS, Riddell SR, Spies T. Costimulation of CD8 $\alpha\beta$  T cells by NKG2D via engagement by MIC induced on virus-infected cells. *Nat Immunol* 2001;2:255-60.
7. Diefenbach A, Raulet DH. Strategies for target cell recognition by natural killer cells. *Immunol Rev* 2001;181:170-84.
8. Diefenbach A, Jensen ER, Jamieson AM, Raulet DH. Rae-1 and H60 ligands of the NKG2D receptor stimulate tumour immunity. *Nature* 2001;413:165-71.
9. Houchins JP, Yabe T, McSherry C, Bach FH. DNA sequence analysis of NKG2, a family of related cDNA clones encoding type II integral membrane proteins on human natural killer cells. *J Exp Med* 1991;173:1017-20.
10. Bauer S, Groh V, Wu J, Steinle A, Phillips JH, Lanier LL, et al. Activating of NK cells and T cells by NKG2D, a receptor for stress-inducible MICA. *Science* 1999;285:727-8.
11. Wu J, Song Y, Bakker ABH, Bauer S, Spies T, Lanier LL, et al. An activating immunoreceptor complex formed by NKG2D and DAP10. *Science* 1999; 285:730-2.

12. Wu J, Cherwinski H, Spies T, Phillips JH, Lanier LL. DAP10 and DAP12 form distinct, but functionally cooperative, receptor complexes in natural killer cells. *J Exp Med* 2000;192:1059-68.
13. Trinchieri G. The choices of a natural killer. *Nature Immunology* 2003;4:509-10.
14. Billadeau DD, Upshaw JL, Schoon RA, Dick CJ, Leibson PJ. NKG2D-DAP10 triggers human NK cell-mediated killing via a Syk-independent regulatory pathway. *Nat Immunol* 2003;4:557-64.
15. Alegre ML, Frauwirth KA, Thompson CB. T-cell regulation by CD28 and CTLA-4. *Nat Rev Immunol* 2001;1:220-8.
16. Cosman D, Mullberg J, Sutherland CL, Chin W, Armitage R, Fanslow W, et al. ULBPs, Novel MHC Class I-related molecules, bind to CMV glycoprotein UL16 and stimulate NK cytotoxicity through the NKG2D receptor. *Immunity* 2001;14:123-33.
17. Cerwenka A, Baron JL, Lanier LL. Ectopic expression of retinoic acid early inducible-1 gene (RAE-1) permits natural killer cell-mediated rejection of a MHC class I-bearing tumor in vivo. *Proc Natl Acad Sci USA* 2001;98:11521-6.
18. Watzl C. The NKG2D receptor and its ligands-recognition beyond the "missing self"? *Microbes and Infection* 2003;5:31-37.
19. Ravetch JV, Lanier LL. Immune inhibitory receptors. *Science* 2000; 290:84-9.
20. Sutherland CL, Chalupny NJ, Schooley K, VandenBos T, Kubin M, Cosman D. UL16-binding proteins, novel MHC class I-related proteins, bind to NKG2D and activate multiple signaling pathways in primary NK cells. *J Immunol* 2002;168: 671-9.
21. Matzinger P. The danger model: a renewed sense of self. *Science* 2002;296:301-5.
22. Jamieson AM, Diefenbach A, McMahon CW, Xiong N, Carlyle JR, Raulet DH. The Role of the NKG2D Immunoreceptor in Immune Cell Activation and Natural Killing. *Immunity* 2002;17:19-29.
23. Jiang K, Zhong B, Gilvary DL, Corliss BC, Hong-Geller E, Wei S, et al. Pivotal role of phosphoinositide-3 kinase in regulation of cytotoxicity in natural killer cells. *Nat Immunol* 2000;1:419-25.

24. Hayakawa Y, Kelly JM, Westwood JA, Darcy PK, Diefenbach A, Raulet D, et al. Cutting edge: tumor rejection mediated by NKG2D receptor-ligand interaction is dependent upon perforin. *J Immunol* 2002;169:5377-81.
25. Baharm S, Bresnahan M, Geraghty DE, Spies T. A second lineage of mammalian major histocompatibility complex class I genes. *Proc Natl Acad Sci* 1994; 91:6259-63.
26. Leelayuwat C, Townsend DC, Degli-Esposti MA, Abraham LJ, Dawkins RL. A new polymorphic and multicopy MHC gene family related to nonmammalian class I. *Immunogenetics* 1994;40:339-51.
27. Groh V, Bahram S, Bauer S, Herman A, Beauchamp M, Spies T. Cell stress-regulated human major histocompatibility complex class I gene expressed in gastrointestinal epithelium. *Proc Natl Acad Sci USA* 1996;93:12445-50.
28. Groh V, Rhinehart R, Secrist H, Bauer S, Grabstein KH, Spies T. Broad tumor-associated expression and recognition by tumor-derived  $\gamma\delta$  T cells of MICA and MICB. *Proc Natl Acad Sci USA* 1999;96:6879-84.
29. Chalupny NJ, Sutherland CL, Lawrence WA, Rein-Weston A, Cosman D. ULBP4 is a novel ligands for human NKG2D. *Biochem Biophys Res Commun* 2003;305:129-135.
30. Das H, Groh V, Kuijl C, Sugita M, Morita CT, Spies T, et al. MICA engagement by human V gamma2 V delta2 T cells enhances their antigen-dependent effector function. *Immunity* 2001;15:83-93.
31. Tieng V, Le Bouguence C, du Merle L, Bertheau P, Desreumaux P, Janin A, et al. Binding of Escherichia coli adhesin AfaE10 to CD55 triggers cell-surface expression of the MHC class I-related molecule MICA. *Proc Natl Acad Sci USA* 2002;99:2977-82.
32. Diefenbach A, Jamieson AM, Liu SD, Shastri N, Raulet DH. Ligands for the murine NKG2D receptor : expression by tumor cells and activation of NK cells and macrophages. *Nat Immunol* 2000;1:119-26.
33. Kurman RJ, Solomon D. The Bethesda system for reporting cervical/vaginal cytologic diagnoses: definitions, criteria, and explanatory notes for terminology and specimen adequacy. New York: Springer-Verlag; 1994.

34. Lu Z, Friess H, Graber HU, Guo X, Schilling M, Zimmermann A, et al. Presence of Two Signaling TGF- $\beta$  receptors in human pancreatic cancer correlates with advanced tumor stage. *Dig Dis Sci* 1997;42:2054-63.

35. Liu SS, Tsang BK, Cheung ANY, Xue WC, Cheng DKL, Ng TY, et al. Anti-apoptotic proteins, apoptotic and proliferative parameters and their prognostic significance in cervical carcinoma. *Eur J Cancer* 2001;37:1104-10.

36. Li Z, Groh V, Strong RK, Spies T. A single amino acid substitution causes loss of expression of a MICA allele. *Immunogenetics* 2000;51:246-48.

37. Gaudieri S, Leelayuwat C, Townend D, Mullberg J, Cosman D, Dawkins RL. Allelic and interlocus comparison of the PERB11 multigene family in the MHC. *Immunogenetics* 1997;45:209-16.

38. Fodil N, Laloux L, Wanner V, Pellet P, Hauptmann G, Mizuki N, et al. Allelic repertoire of the human MHC class I MICA gene. *Immunogenetics* 1996;44:351-7.

39. Yao Z, Volgger A, Helmberg W, Keller E, Fan LA, Chandanayingyong D, et al. Definition of new alleles of MIC-A using sequence-based typing. *Eur J Immunogenetics* 1999;26:225-32.

40. Visser CJT, Tilanus MGJ, Tatari Z, van der Zwan A, Bakker R, Rozemuller EH, et al. Sequencing-based typing of MICA reveals 33 alleles: a study on linkage with classical HLA genes. *Immunogenetics* 1999; 49 :561-566.

41. Dunn DS, Williamson JF, Cattley SK, Tay GK, Guadieri S, Leelayuwat C, et al. Co-evolution of HLA-B and PERB11.1 (MICA): Significance of independent triplet expansion within the transmembrane region of PERB11.1 (MICA). *J Mol Evol* 2000;50:359-65.

42. Groh V, Li YQ, Cioca D, Hunder NN, Wang W, Riddell SR, et al. Efficient cross-priming of tumor antigen-specific T cells by dendritic cells sensitized with diverse anti-MICA opsonized tumor cells. *Proc Natl Acad Sci USA*. 2005;102:6461-6.

43. Carbone E, Neri P, Mesuraca M, Fulciniti MT, Otsuki T, Pende D, et al. HLA class I, NKG2D, and natural cytotoxicity receptors regulate multiple myeloma cell recognition by natural killer cells. *Blood* 2005;105:251-8.

44. Tay GK, Hui J, Gaudier S, Schmitt-Egenolf M, Martinez OP, Leelayuwat C, et al. PERB11(MIC): a polymorphic major histocompatibility complex gene is expressed in skin and single nucleotide polymorphisms are associated with psoriasis. *Clin Exp Immunol* 2000;117:2-5.
45. Mizuki N, Ota M, Kimura M. Triplet repeat polymorphism in the transmembrane region of the MICA gene: strong association of six GCT repetition with Behcet disease. *Proc Natl Acad Sci USA* 1997;94:1298-303
46. Romphruk AV, Leelayuwat C, Barusrux S, Puapairoj C, Romphruk Ar, Urwijitaroon Y. DNA Typing of the HLA-A, -B and -C Genes : Possible MHC Class I Haplotypes in the Northeastern-Thais. *J Med Assoc Thai* 1997;80(Suppl 1):S13-19.
47. Ooi EE, Ren EC, Chan SH. Association between microsatellites within the human MHC and nasopharyngeal carcinoma. *Int J Cancer* 1997;74:229-32.