

เอกสารอ้างอิง

- กรมพัฒนาที่ดิน. *ความรู้เรื่องดินเค็มภาคตะวันออกเฉียงเหนือ โครงการพัฒนาดินเค็มภาคตะวันออกเฉียงเหนือ*. กรมพัฒนาที่ดิน กระทรวงเกษตรและสหกรณ์. กรุงเทพฯ. 2527. 159 หน้า.
- พรเทพ ถนณแก้ว และ ปิยะดา ชีระกุลพิศุทธิ์. *ผลกระทบของเกลือโซเดียมคลอไรด์ต่อลักษณะทางสรีรวิทยาของข้าวพันธุ์ขาวดอกมะลิ 105 และ กข.6* รายงานการวิจัยเสนอมหาวิทยาลัยขอนแก่น. 2544.
- สมศรี อรุณินท์. พืชทนเค็ม. *พัฒนาที่ดิน*. เม.ย. 2539; 26(287): 38-46.
- สรวงสุดา สืออ่อน. *ผลของเกลือโซเดียมคลอไรด์ต่อสรีรวิทยาของมะเขือเทศ*. วิทยานิพนธ์ปริญญาวิทยาศาสตรมหาบัณฑิต คณะวิทยาศาสตร์ มหาวิทยาลัยขอนแก่น. 2544.
- สำนักงานเศรษฐกิจการเกษตร. *การสัมมนาวิชาการข้าวภาคตะวันออกเฉียงเหนือ*. สถาบันวิจัยข้าว กรมวิชาการเกษตร กระทรวงเกษตรและสหกรณ์. หจก. ขอนแก่นการพิมพ์. ขอนแก่น. 2542.
- อาทิตย์ยา ฉิมรักแก้ว. *สรีรวิทยาและกายวิภาคศาสตร์ของข้าวทนเค็ม*. วิทยานิพนธ์ปริญญาวิทยาศาสตรมหาบัณฑิตมหาวิทยาลัยขอนแก่น. คณะวิทยาศาสตร์ มหาวิทยาลัยขอนแก่น. 2544.
- Ahn SW, Bonman JM, Brandon DM, Groth DE, Gunnell PS, Hibono H, et al. Compendium of rice diseases. In: Webster, Gunnell, editors. **The American Phytopathological Society**. USA: St. Paul, Minnesota; 1992.
- Alexandrova KS, Conger BV. Isolation of two somatic embryogenesis-related genes from orchardgrass (*Dactylis glomerata*). **Plant Sci** 2002; 162: 301-307.
- Allard F, Houde M, Krol M, Ivanov A, Huner NPA, Sarhan F. Betaine improves freezing tolerance in wheat. **Plant Cell Physiol** 1998; 39: 1194-1202.
- Anthraper A, Dubois JD. The effect of NaCl on growth, N₂ fixation (Acetylene reduction), and percentage total nitrogen in *Leucaena leucocephala* (Leguminosae) var. K-8. **Amer J Botany** 2003; 90(5): 683-692.
- Artitya C. **Physiology and anatomy of salt tolerant rice** [Master of science thesis in biology]. Khon Kaen: The Graduate School, Khon Kaen University; 2000. [in Thai].

- Bell HL, O'Leary JW. Effects of salinity on growth and cation accumulation of *Sporobolus virginicus* (Poaceae). *Amer J Botany* 2003; 90(10): 1416-1424.
- Bhumbla D, Abrol I. Saline and sodic soils : soils and rice. *International Rice Research Institute* Philippines: Manila; 1978.
- Blumwald B. Sodium transport and salt tolerance in plants. *Curr Opinion Cell Biol* 2000; 12: 431-434.
- Borrone WJ, Kuhn DN, Schnell RJ. Isolation, characterization, and development of WRKY genes as useful genetic markers in *Theobroma cacao*. *Theor App Genet* 2004; 109: 495-507.
- Botia P, Carvajal M, Cerdá A, Martinez V. Response of eight *Cucumis melo* cultivars to salinity during germination and early vegetative growth. *Agronomie* 1998; 18: 503-513.
- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dry binding. *Anal Biochem* 1976; 72: 248-254.
- Cano EA, Pérez-Alfocea F, Moreno V, Caro M, Bolarin MC. Evaluation of salt tolerance in cultivates and wild tomato species through *in vitro* shoot apex culture. *Plant Cell Tissue and Organ Culture* 1998; 53: 19-26.
- Chen CCS, Plant AL. Salt-induced protein synthesis in tomato roots: the role of ABA. *J Exp Botany* 1999; 50: 677-687.
- Chen GP, Ma WS, Huang ZJ, Xu T, Xue YB, Shen YZ. Isolation and characterization of TaGSK1 involved in wheat salt tolerance. *Plant Sci* 2003; 165(6): 1369-1375.
- Chourey K, Ramani S, Apte SK. Accumulation of LEA proteins in salt (NaCl) stressed young seedlings of rice (*Oryza sativa* L.) cultivar Bura Rata and their degeneration during recovery from salinity stress. *J Plant Physiol* 2003; 1-10.
- Çiçek N, Çakırlar H. The effect of salinity on some physiological parameters in two maize cultivars. *Bulg J Plant Physiol* 2002; 28(1-2): 66-74.
- Claes B, Dekeyser R, Villarroel R, Van den Bulcke M, Bauw G, Van Montagu M, et al. Characterization of a rice gene showing organ-specific expression in response to salt stress and drought. *Plant Cell* 1990; 2: 19-27.

- Cordovilla MDP, Ligeró F, Lluch C. Effect of salinity on growth, nodulation and nitrogen assimilation in nodules of faba bean (*Vicia faba* L.). **App Soil Ecol** 1999; 11: 1-7.
- Cortina C, Culiáñez-Macià FA. Tomato abiotic stress enhanced tolerance by trehalose biosynthesis. **Plant Sci** 2005; 169: 75-82.
- Cramer GR, Nowak RS. Supplemental manganese improve the relative growth, net assimilation and photosynthetic rate of salt-stressed barley. **Physiol Plant** 1992; 84: 600-605.
- Crowe JH, Hoekstra FA, Crowe LM. Arabidopsis. **Annu Rev Physiol** 1992; 54: 579-599.
- Cuartero J, Fernández-Muñoz R. Tomato and salinity. **Sci Hort** 1999; 78: 83-125.
- de Souza Filho GA, Ferreira BS, Dias JM, Queiroz KS, Branco AT, Bressan-Smith RE, et al. Accumulation of SALT protein in rice plants as response to environment stresses. **Plant Sci** 2003; 164: 623-628.
- Demiral T, Türkan I. Exogenous glycinebetain affects growth and proline accumulation and retards senescence in two rice cultivars under NaCl stress. **Envi Exp Botany** 2006; 56: 72-79.
- Dobey RS. Protein synthesis by plants under stressful conditions. In: Pessarakli M, editor. **Handbook of Plant and Crop Stress**. New York: Marcel Dekker, Inc; 1994. p. 277-299.
- Dubouzet JG, Sakuma Y, Ito Y, Kasuga M, Dubouzet EG, Miura S, et al. *OsDREB* genes in rice, *Oryza sativa* L., encode transcription activators that function in drought-, high salt- and cold-responsive gene expression. **Plant J** 2003; 33: 751-763.
- Duwayri M, Tran DV, Nguyen VN. Reflections on yield gaps in rice production: How to narrow the gaps. In: Papademetriou MK, Dent FJ, Herath EM, editors. **Food and agriculture organization of the united nations regional office for Asia and the Pacific**. Bangkok, Thailand; 2000. p. 26-45.
- Ehsanpour AA, Amini F. Effect of salt and drought stress on acid phosphatase activities in alfalfa (*Medicago sativa* L.) explants under in vitro culture. **African J Biotechnol** 2003; 2(5): 133-135.
- Espelund M, Saeboe-Larssen S, Hughes DW, Galau GA, Larsen F, Jakobsen KS. Late embryogenesis-abundant genes encoding proteins with different

- numbers of hydrophilic repeats are regulated differentially by abscisic acid and osmotic stress. *Plant J* 1992; 2: 241-252.
- Eulgem T, Rushton PJ, Robatzek S, Somssich IE. The WRKY superfamily of plant transcription factor. *Trends Plant Sci* 2000; 5(5): 199-206.
- Flowers TJ, Yeo AR. Breeding for salinity resistance in crop plants: where next? *Aust J Plant Physiol* 1995; 22: 875-884.
- Fukutaku Y, Yamada Y. Source of proline nitrogen in water-stressed soybean (*Glycine max*). Part II. Fate of ¹⁵N-labeled protein. *Plant Physiol* 1984; 61: 622-628.
- Garcia AB, Engler JDA, Claes B, Villarroel R, Montagu MV, Gerats T, et al. The expression of the salt-responsive gene *salT* from rice is regulated by hormonal and developmental cues. *Planta* 1998; 207: 172-180.
- Garg AK, Kim JK, Owens TG, Ranwala AP, Choi YD, Kochian LV, et al. Trehalose accumulation in rice plants confers high tolerance levels to different abiotic stresses. *PNAS* 2002; 99(25): 15898-15903.
- Ghassemi F, Jakeman AJ, Nix HA. **Salinisation of land and water resources: Human causes, extent, management and case studies**. UK: UNSW Press, Sydney, Australia and CAB International, Wallingford; 1995.
- Gisbert C, Rus AM, Bolarin MC, López-Coronado JM, Arrillaga I, Montesinos C, et al. The yeast *HAL1* gene improves salt tolerance of transgenic tomato. *Plant Physiol* 2000; 123: 393-402.
- Gregorio GB, Dharmawansa S, Mendoza RD. Screening rice for salinity tolerance. *IRRI* 1997; 22: 2-23.
- Grierson W. Beneficial aspects of stress on plants. In: Pessarakli M, editor. **Handbook of Plant and Crop Stress**. New York: Macel Dekker, Inc; 1994. p. 645-657.
- Gu Z, Wang J, Huang J, Zhang H. Cloning and characterization of a novel rice family encoding putative dual-specificity protein kinases, involved in plant responses to abiotic and biotic stresses. *Plant Sci* 2005; 169: 470-477.
- Guerta CQ, Kirk GJD. Tolerance of rice germplasm to salinity and other soil chemical stresses in tidal wetlands. *Field Crops Res* 2002; 76: 111-121.
- Guo XL, Cao YR, Cao ZY, Zhao YX, Zhang H. Molecular cloning and characterization of a stress-induced peroxiredoxin Q gene in halophyte *Suaeda salsa*. *Plant Sci* 2004; 167: 969-975.

- Hara K, Yagi M, Kusano T, Sano H. Rapid sytemic accumulation of transcripts encoding a tobacco WRKY transcription factor upon wounding. *Mol Gen Genet* 2000; 263: 30-37.
- Heuer B. Osmoregulatory role of proline in water- and salt-stressed plants. In: Pessarakli M, editor. *Handbook of Plant and Crop Stress*. New York: Macel Dekker, Inc; 1994. p. 363-382.
- Hien DT, Jacobs M, Angenon G, Hermans C, Thu TT, Son LV, et al. Proline accumulation and Δ^1 -pyrroline-5-carboxylate synthetase gene properties in three rice cultivars differing in salinity and drought tolerance. *Plant Sci* 2003; 165: 1059-1068.
- Hirano K, Teraoka T, Yamanaka H, Harashima A, Kunisaki A, Takahashi H, et al. Novel mannose-binding rice lectin composed of some isolectins and its relation to a stress-inducible *salT* gene. *Plant Cell Physiol* 2000; 41(3): 258-267.
- Hong B, Barg R, Ho THD. Developmental and organ-specific expression of an ABA- and stress-induced protein in barley. *Plant Mol Biol* 1992; 18: 663-674.
- Hong Z, Lakkineni K, Zhang Z, Verma DPS. Removal of feedback inhibition of Δ^1 -pyrroline-5-carboxylate synthetase results in increased proline accumulation and protection of plants from osmotic stress. *Plant Physiol* 2000; 122: 1129-1136.
- Hoyos ME, Zhang S. Calcium-independent activation of salicylic acid-induced protein kinase and a 40-kilodalton protein kinase by hyperosmotic stress. *Plant Physiol* 2000; 122: 1355-1363.
- Huang J, Duman JG. Cloning and characterization of a thermal hysteresis (antifreeze) protein with DNA-binding activity from winter bittersweet nightshade, *Solanum dulcamara*. *Plant Mol Biol* 2002; 48: 339-350.
- Hurkman WJ, Tanaka CK. The effects of salt on the pattern of protein synthesis in barley roots. *Plant Physiol* 1987; 83: 517-524.
- Hurkman WJ, Tao HP, Tanaka CK. Germin-like polypeptides increase in barley roots during salt stress. *Plant Physiol* 1991; 97: 366-374.
- Igarashi Y, Yoshiba Y, Sanada S, Yamaguchi-Shinozaki K, Wada K, Shinozaki K. Characterization of the gene for Δ^1 -pyrroline-5-carboxylate synthetase

- and correlation between the expression of the gene and salt tolerance in *Oryza sativa* L. **Plant Mol Biol** 1997; 33: 857-865.
- Iyer S, Caplan A. Products of proline catabolism can induced osmotically regulated genes in rice. **Plant Physiol** 1998; 116: 203-211.
- Jang IC, Oh SJ, Seo JS, Choi WB, Song SI, Kim CH, et al. Expression of a bifunctional fusion of the *Escherichia coli* gene for trehalose-6-phosphate synthase and trehalose-6-phosphate phosphatase in transgenic rice plants increases trehalose accumulation and abiotic stress tolerance without stunting growth. **Plant Physiol** 2003; 131: 516-524.
- Jain S, Nainawatee HS, Jain RK, Chowdhury JB. Salt-tolerance in *Brassica juncea* L. II. Salt-stress induced changes in polypeptide pattern of *in vitro* selected NaCl tolerance plant. **Euphytica** 1993; 65: 107-112.
- Jearakongman S, Rajatasereekul S, Naklang K, Romyen P, SkulKhu E, Jumpaket B, et al. Growth and grain yield of contrasting rice cultivars grown under different conditions of water availability. **Field Crops Res** 1995; 44: 139-150.
- Jeong MJ, Park SC, Byun MO. Improvement of salt tolerance in transgenic potato plants by glyceraldehydes-3-phosphate dehydrogenase gene transfer. **Mol Cells** 2001; 12(2): 185-189.
- Jha KP, Moorthy BTS, Rao KS. Rice. In: Rathore PS, editor. **Techniques and management of field crop production**. Agrobios (India): Jodhpur; 2005. p. 6-40.
- Kadpal RP, Rao NA. Alterations in the biosynthesis of proteins and nucleic acids in finger millet (*Eleusine coracana*) seedlings during water stress and the effect of proline on protein biosynthesis. **Plant Sci** 1985; 40: 73-79.
- Kavi Kishor PB, Hong Z, Miao G, Hu CAA, Verma DPS. Over expression of Δ^1 -pyrroline-5-carboxylate synthetase increases proline overproduction and confers osmtolerance in transgenic plants. **Plant Physiol** 1995; 108: 1387-1394.
- Kawasaki S, Borchert C, Deyholos M, Wang H, Brazille S, Kawai K, et al. Gene expression profiles during the initial phase of salt stress in rice. **Plant Cell** 2001; 13: 889-905.
- Khush GS. Green revolution: the way forward. **Nature Rev Genet** 2001; 2: 815-822.

- Kishitani S, Watanabe K, Yasuda S, Arakawa K, Takabe T. Accumulation of glycinebetaine during cold acclimation and freezing tolerance in leaves of winter and spring barley plants. *Plant cell Envi* 1994; 17: 89-95.
- Komatsu S, Konishi H, Shen S, Yang G. *Rice Proteomics: A step towards functional analysis of the rice genome* [online] 2003 [cited 2004 Dec 15]. Available from: <http://www.mcponline.org/cgi/content/full/2/1/2>
- Kumar SG, Reddy AM, Sudhakar C. NaCl effects on proline metabolism in two high yielding genotypes of mulberry (*Morus alba* L.) with contrasting salt tolerance. *Plant Sci* 2003; 165: 1245-1251.
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; 227: 680-685.
- Lanceras JC, Huang ZL, Naivikul O, Vanavichit A, Ruanjaichon V, Tragoonrung S. Mapping of genes for cooking and eating qualities in Thai jasmine rice (KDML105). *DNA Res* 2000; 7: 93-101.
- Lee KS, Choi WY, Ko JC, Kim TS, Gregorio GB. Salinity tolerance of japonica and indica rice (*Oryza sativa* L.) at the seedling stage. *Planta* 2003; 216: 1043-1046.
- Lee SC, Lee MY, Kim SJ, Jun SH, An G, Kim SR. Characterization of an Abiotic Stress-inducible Dehydrin Gene, *OsDhn1*, in Rice (*Oryza sativa* L.). *Mol Cells* 2004; 19(2): 212-218.
- Lefevre I, Gratia E, Lutts S. Discrimination between the ionic and osmotic components of salt stress in relation to free polyamine level in rice (*Oryza sativa*). *Plant Sci* 2001; 161: 943-952.
- Liu J, Zhu JK. Proline accumulation and salt-stress-induced gene expression in a salt hypersensitive mutant of Arabidopsis. *Plant Physiol* 1997; 114: 591-596.
- Lopez F, Vansuyt G, Fourcroy P, Case-Delbart F. Accumulation of a 22-kDa protein and its mRNA in the leaves of *Raphanus sativas* in response to salt stress or water stress. *Physiol Plant* 1994; 91: 605-614.
- Maclean JL, Dawe DC, Hardy B, Hettel GP, editors. *Rice Almanac: Source book for the most important economic activity on earth*. 3rd ed. Wallingford: CABI Publishing; 2002.
- McCue KF, Hanson AD. Drought and salt tolerance: towards understanding and application. *Trends in Biotechnol* 1990; 8: 358-362.

- Miller RW, Gardiner DT. **Soils In Our Environment**. 8th ed. New Jersey: Upper Saddle River, Prentice-Hall Inc; 1998.
- Misra AN, Sahu SM, Misra M, Singh P, Meera I, Das N, et al. Sodium chloride induced changes in leaf growth, and pigment and protein contents in two rice cultivars. **Biol Plant** 1997; 39: 257-262.
- Moghaieb REA, Saneoka H, Fujita K. Effect of salinity on osmotic adjustment, glycinebetaine accumulation and the betaine aldehyde dehydrogenase gene expression in two halophytic plants, *Salicornia europaea* and *Suaeda maritima*. **Plant Sci** 2004; 166: 1345-1349.
- Munns R. **The impact of salinity stress** [online] nd [cited 2004 Dec 15]. Available from: http://www.plantstress.com/Articles/salinity_i/salinity_i.htm
- Munns R. Comparative physiology of salt and water stress. **Plant cell Envi** 2002; 25: 239-250.
- Murray SS. **Searching for the origins of African rice domestication** [online] nd [cited 2005 May 20]. Available from: <http://antiquity.ac.uk/ProjGall/murray>.
- Narita Y, Taguchi H, Nakamura T, Ueda A, Shi W, Takabe T. Characterization of the salt-inducible methionine synthase from barley leaves. **Plant Sci** 2004; 167: 1009-1016
- Nikolopoulos D, Manetas Y. Compatible solutes and in vitro stability of *Salsola* soda enzymes: proline incompatibility. **Phytochemistry** 1991; 30: 411-413.
- Ohta M, Hayashi Y, Nakashima A, Hamada A, Tanaka A, Nakamura T, et al. Introduction of a Na^+/H^+ antiporter gene from *Atriplex gmelini* confers salt tolerance to rice. **FEBS Letters** 2002; 532: 279-282.
- Oosterbaan RJ. **Soil salinity** [online] 2003 [cited 2005 Feb 24]. Available from: <http://www.waterlog.info/salinity.pdf>
- Pengthaisong S. **Characterization and purification of NaCl-inductive proteins in plaunoi (Croton sublyratus Kurz.) callus** [Master of science thesis in biology]. Khon Kaen: The Graduate School, Khon Kaen University; 2001.
- Piao HL, Lim JH, Kim SJ, Cheong GW, Hwang I. Constitutive over-expression of AtGSK1 induces NaCl stress responses in the absence of NaCl stress and results in enhanced NaCl tolerance in Arabidopsis. **Plant J** 2001; 27(4): 305-314.

- Pnueli L, Hallak-Herr E, Rozenberg M, Cohen M, Goloubinoff P, Kaplan A, et al. Molecular and biochemical mechanisms associated with dormancy and drought tolerance in the desert legume *Retama raetam*. **Plant J** 2002; 31: 319-330.
- Ponnamperuma FN. Role of cultivar tolerance in increasing rice production on saline lands. In: Staple RC, Toenniessen GH, editors. **Salinity Tolerance in Plants: Strategies for Crop Improvement**. New York: John Wiley & Sons, Inc. 1984. p. 255-271.
- Powar SL, Mehta VB. Salt tolerance of some cultivars of rice (*Oryza sativa* L.) at germination stage. **Curr Agri Res** 1995; 19: 43-45.
- Prasad SR, Bagali PG, Hittalmani S, Shashidhar HE. Molecular mapping of quantitative trait loci associated with seedling tolerance to salt stress in rice (*Oryza sativa* L.). **Curr Sci** 2000; 78: 162-164.
- Ramagopal S. Salinity stress induced tissue-specific proteins in barley seedlings. **Plant Physiol** 1987; 84: 324-331.
- Rani UR, Reddy AR. Salt stress-responsive polypeptides in germinating seeds. **J Plant Physiol** 1994; 143: 250-253.
- Rhodes D, Hanson AD. Quaternary ammonium and tertiary sulfonium compounds in higher plants. **Annu Rev Plant Physiol Plant Mol Biol** 1993; 44: 357-384.
- Rhodes D, Pich PJ, Brunk DG, Ju GC, Rhodes JC, Pauly MH, et al. Development of two isogenic sweet corn hybrids differing for glycine betaine content. **Plant Physiol** 1989; 91: 1112-1121.
- Rizhsky L, Davletova S, Liang H, Mittler R. The zinc finger protein Zat12 is required for cytosolic ascorbate peroxidase 1 expression during oxidative stress in Arabidopsis. **J Biol Chem** 2004; 279: 11736-11743.
- Rizhsky L, Liang H, Mittler R. The combined effect of drought stress and heat shock on gene expression in tobacco. **Plant Physiol** 2002; 130: 1143-1151.
- Rohila JS, Jain RK, Wu R. Genetic improvement of basmati rice for salt and drought tolerance by regulated expression of a barley *Hva1* cDNA. **Plant Sci** 2002; 163: 525-532.

- Romero-Aranda R, Soria T, Cuartero J. Tomato plant-water uptake and plant-water relationships under saline growth conditions. *Plant sci* 2001; 160: 265-272.
- Roxas VP, Smith RKJ, Allen ER, Allen RD. Overexpression of glutathione S-transferase/glutathioneperoxidase enhances the growth of transgenic tobacco seedlings during stress. *Nat Biotechnol* 1997; 15: 988-991.
- Salekdeh GH, Siopongco J, Wade LJ, Ghareyazie B, Bennett J. A proteomic approach to analyzing drought- and salt-responsiveness in rice. *Field Crops Res* 2002; 76: 199-219.
- Sambrook J, Russell DW. *Molecular cloning: A laboratory manual*. 3rd edition. New York: Cold Spring Harbor Laboratory Press; 2001.
- Saneoka H, Nagasaka C, Hahn DT, Yang WJ, Premachandra GS, Joly RT, et al. Salt tolerance of glycinebetaine-deficient and containing maize lines. *Plant Physiol* 1995; 107: 631-638.
- Sanger F, Nicklen S, Coulson AR. DNA sequencing with chain-terminating inhibitor. *Proc Natl Acad Sci USA* 1977; 74: 5463-5467.
- Schobert BH, Tschesche H. Unusual solution properties of proline and its interaction with proteins. *Biochim Biophys Acta* 1978; 541: 270-277.
- Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, Oono Y, et al. Monitoring the expression profiles of 7,000 *Arabidopsis* genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant J* 2002; 31: 279-292.
- Serrano R. Salt tolerance in plants and microorganisms: toxicity targets and defense responses. *Int Rev Cytol* 1996; 1-52.
- Shannon MC, Rhoades JD, Draper JH, Scardaci SC, Spyres MD. Assessment of salt tolerance in rice cultivars in response to salinity problems in California. *Crop Sci* 1998; 38: 394-398.
- Shah SH, Tobita S, Shono M. Supplemental calcium regulates proline accumulation in NaCl-stressed suspension cultures of *Oryza sativa* L. at the level of mRNA translation. *Pakistan J Biol Sci* 2001; 4(6): 707-710.
- Shen YG, Zhang WK, He SJ, Zhang JS, Liu Q, Chen SY. An EREBP/AP2-type protein in *Triticum aestivum* was a DRE-binding transcription factor induced by cold, dehydration and ABA stress. *Theor App Genet* 2003; 106: 923-930.

- Shih CY, Kao CH. Induction of acid phosphatase in detached rice leaves under stress conditions. *Bot Bull Acad Sin* 1998; 39: 29-32.
- Shirata K, Takagishi H. Salt-induced accumulation of 26 and 27 kD proteins in cultured cells of rice plant. *Soil Sci Plant Nutr* 1990; 36: 153-157.
- Silva JV, Lacerda CF, Costa PHA, Filho JE, Filho EG, Prisco JT. Physiological responses of NaCl stressed cowpea plants grown in nutrient solution supplemented with CaCl₂. *Braz J Plant Physiol* 2003; 15(2): 99-105.
- Silveira JAG, Melo ARB, Viégas RA, Oliveira JTA. Salinity-induced effects on nitrogen assimilation related to growth in cowpea plants. *Envi Exp Botany* 2001; 46: 171-179.
- Singh NK, Handa AK, Hasegawa PM, Bressan RA. Proteins associated with adaptation of cultured tobacco cells to NaCl. *Plant Physiol* 1985; 79: 126-137.
- Smirnoff N, Cumbes QJ. Hydroxyl radical scavenging activity of compatible solutes. *Phytochemistry* 1989; 28: 1057-1060.
- Szabolcs I. Soils and salinization. In: Pessarakli M, editor. *Handbook of Plant and Crop Stress*. New York: Macel Dekker, Inc; 1994. p. 3-12.
- Taniyama T. Salt injuries and morphology. In: Matsuo T, Hoshikawa K, editors. *Science of the rice plant*. Morphology v.1 Tokyo: Nosan Gyoson Bunka Kyokai; 1993. p. 601-605.
- The genetic engineering newsletter. *Transgenic drought- and salt-tolerant plants* [online] 2004 [cited 2005 Aug 20]. Available from: http://www.oeko.de/gennews_engl.htm.
- The office of the gene technology regulator. The biological and ecology of rice (*Oryza sativa* L.) in Australia. *Australian government* 2005. p. 1-26.
- Tinjuangjun P, Loc NT, Gatehouse AMR, Gatehouse JA, Christou P. Enhanced insect resistance in Thai rice varieties generated by particle bombardment. *Mol Breeding* 2000; 6: 391-399.
- Turgeon BG, Garber RC, Yoder OC. Development of a fungal transformation system based on selection of sequence with promoter activity. *Mol Cell Biol* 1987; 7: 3297-3305.
- Uchida A, Jagendorf AT, Hibino T, Takabe T, Takabe T. Effects of hydrogen peroxide and nitric oxide on both salt and heat stress tolerance in rice. *Plant Sci* 2002; 163: 515-523.

- Ülker B, Somssich IE. WRKY transcription factors: from DNA binding towards biological function. *Curr Opin Plant Biol* 2004; 7: 491-498.
- Ungar IA. Effect of salinity on seed germination, growth, and ion accumulation of *Atriplex patula* (Chenopodiaceae). *Amer J Botany* 1996; 83(5): 604-607.
- Venekamp JH, Lamp JE, Koot JTM. Organic acids as sources of drought-induced proline synthesis in field bean plants, *Vicia faba* L. *J Plant Physiol* 1989; 133: 654-659.
- Vitoon P. *Thai Rice, The rice of freedom* [online] nd [cited 2005 July 4]. Available from: <http://www.panap.net/ricecampaign/docs/greenet.pdf> and www.greenetorganic.com
- Voetberg GS, Sharp RE. Growth of the maize primary root at low water potentials. Part III. Role of increased proline deposition in osmotic adjustment. *Plant Physiol* 1991; 96: 1125-1130.
- Wahome PK, Jesch HH, Pinker I. Effect of sodium chloride stress on *Rosa* plants growing in vitro. *Scientia Horticulturae* 2001; 90: 187-191.
- Wang YJ, Zhang ZG, He XJ, Zhou HL, Wen YX, Dai JX, et al. A rice transcription factor OsbHLH1 is involved in cold stress response. *Theor App Genet* 2003; 107: 1402-1409.
- Wanichananan P, Busayanggoon T, Kirdmanee C. Physiological index for selection of salt-tolerant rice (*Oryza sativa* L.). In: Jamratlertluck S, Jindamanee V, editors. *Proceeding of Conference on Functional Genomics of Rice and Seed Biotechnology*. BioThailand 2001 from Research to Market; 2001 Nov 8-9; Queen Sirikit National Convention Center. Bangkok, Thailand. p. 1-8.
- Watad AA, Reinhold L, Lerner HR. Comparison between a soluble NaCl-selected *Nicotiana* cell line and the wild type. *Plant Physiol* 1983; 73: 634-629.
- Weigel P, Weretilnyk EA, Hanson AD. Betaine aldehyde oxidation by spinach chloroplasts. *Plant Physiol* 1986; 82: 753-759.
- Winicov I, Bastola DR. Transgenic overexpression of the transcription factor *Alfin1* enhances expression of the endogenous *MsPRP2* gene in alfalfa and improves salinity tolerance of the plants. *Plant Physiol* 1999; 120: 473-480.

- Wongpornchai S, Dumri K, Jongkaewwattana S, Siri B. Effects of drying methods and storage time on the aroma and milling quality of rice (*Oryza sativa* L.) cv. Khao Dawk Mali 105. **Food Chemistry** 2004; 87: 407-414.
- Wu KL, Guo ZJ, Wang HH, Li J. The WRKY transcription factors in rice and Arabidopsis and their origins. **DNA Res** 2005; 12: 9-26.
- Xu D, Duan X, Wang B, Hong B, David Ho TH, Wu R. Expression of late embryogenesis abundant protein gene, *HVA1*, from barley confers tolerance to water deficit and salt stress in transgenic rice. **Plant Physiol** 1996; 110: 249-257.
- Xue ZY, Zhi DY, Xue GP, Zhang H, Zhao YX, Xia GM. Enhanced salt tolerance of transgenic wheat (*Triticum aestivum* L.) expressing a vacuolar Na^+/H^+ antiporter gene with improved grain yields in saline soils in the field and a reduced level of leaf Na^+ . **Plant Sci** 2004; 167: 849-859.
- Yamaguchi SK, Kasuga M, Liu Q, Nakashima K, Sakuma Y, Abe H, et al. Biological mechanisms of drought stress response. **JIRCAS Working Report** 2002. p. 1-8.
- Yen HE, Edwards GE, Grimes HD. Characterization of a salt-responsive 24-kilodalton glycoprotein in *Mesembryanthemum crystallinum*. **Plant Physiol** 1994; 105: 1179-1187.
- Yoshihashi T. Does Drought Condition Induce the Aroma Quality of Aromatic Rice?. **JIRCAS Newsletter** 2005; 43: 4.
- Zapata PJ, Serrano M, Pretel MT, Amorós A, Botella MÁ. Changes in ethylene evolution and polyamine profiles of seedlings of nine cultivars of *Lactuca sativa* L. in response to salt stress during germination. **Plant Sci** 2003; 164: 557-563.
- Zapata PJ, Serrano M, Pretel MT, Amorós A, Botella MÁ. Polyamines and ethylene changes during germination of different plant species under salinity. **Plant Sci** 2004; 167: 781-788.
- Zhang HX, Hodson JN, Williams JP, Blumwald E. Engineering salt-tolerant Brassica plants: Characterization of yield and seed oil quality in transgenic plants with increased vacuolar sodium accumulation. **PNAS** 2001; 98(22): 12832-12836.

- Zhang W, Peumans WJ, Barre A, Astoul CH, Rovira P, Rouge P, et al. Isolation and characterization of a jacalin-related mannose-binding lectin from salt-stressed rice (*Oryza sativa*) plants. *Planta* 2000; 210: 970-987.
- Zhang Y, Somantri IH, Tobita S, Nagamine T, Senboku T. Variation of salt tolerance at germination in cultivated rice (*Oryza sativa* L.) varieties. *Sabao J* 1996; 28(2): 15-23.
- Zhang Y, Wang L. The WRKY transcription factor superfamily: its origin in eukaryotes and expansion in plants. *BMC Evolutionary Biol* 2005; 5(1): 1-12.
- Zhao F, Guo S, Zhang H, Zhao Y. Expression of yeast *SOD2* in transgenic rice results in increased salt tolerance. *Plant Sci* 2006; 170: 216-224.
- Zhu B, Chen TH, Li PH. Expression of three osmotin-like protein genes in response to osmotic stress and fungal infection in potato. *Plant Mol Biol* 1995; 28: 17-26.
- Zhu B, Su J, Chang M, Verma DPS, Fan YL, Wu R. Overexpression of a Δ^1 -pyrroline-5-carboxylate synthetase gene and analysis of tolerance to water- and salt-stress in transgenic rice. *Plant Sci* 1998; 139: 41-48.
- Zörb C, Schmitt S, Neeb A, Karl S, Linder M, Schubert S. The biochemical reaction of maize (*Zea mays* L.) to salt stress is characterized by a mitigation of symptoms and not by a specific adaptation. *Plant Sci* 2004; 167: 91-100.