

## เอกสารอ้างอิง

- กสิณ สุวตะพันธ์. (1958). On the Nymphaeaceae of Thailand. ใน **Natural Bulletin of Siam Society**. (Vol.17).
- คำารพ รัตน์สุด. (2553). พันธุ์วิศวกรรมของพืช. คณะเกษตรศาสตร์และทรัพยากรธรรมชาติ มหาวิทยาลัยนเรศวร.
- คุณหญิงสุชาดา ศรีเพ็ญ, วีรญา บุญเตี้ย และวิสาखा เพียรสุภาพ. (2550). การจำแนกพันธุ์บัว. **The Proceeding of IWGS Annual Symposium**, 181 – 189.
- ณรงค์ โฉมเฉลา. (2550). บัวพื้นเมืองของไทย. **วารสารไม้ประดับ**. 32(2), พฤษภาคม – สิงหาคม.
- เรืออากาศเอกหญิงปริมลภ (วสุวัต) ชูเกียรติมัน. (2552). Review of the Waterlily in Thailand. **WGI Online**. 8(3). สืบค้นเมื่อ 19 กรกฎาคม 2554, จาก <http://www.watergardenersinternational.org/journal/4-3/primlarp/page1.html>.
- พระยาวิจิตรวินัยนทร. (2499). The Ornamental Plant of Thailand. **The Science Bulletin Issue** December, B.E.
- ยุวดี วัฒนสุนทร, สุมาลี เลี่ยมทอง และศุภวรรณ พรหมเพรา. (2553). **คุณค่าทางอาหารของเมล็ดบัวสาย จากอำเภอปากพนัง จังหวัดนครศรีธรรมราช**. สืบค้นเมื่อ 1 พฤศจิกายน 2554, จาก [http://www.scisoc.or.th/stt/36/sec\\_h/paper/STT36\\_H\\_H0017.pdf](http://www.scisoc.or.th/stt/36/sec_h/paper/STT36_H_H0017.pdf).
- สุรินทร์ ปิยะโชคณากุล. (2545). จีโนมและเครื่องหมายดีเอ็นเอ: ปฏิบัติการอาร์เอพีดีและเอเอฟแอลพี, สำนักพิมพ์มหาวิทยาลัยเกษตรศาสตร์, กรุงเทพฯ.
- สุรินทร์ ปิยะโชคณากุล. (2552). **เครื่องหมายดีเอ็นเอ: จากพื้นฐานสู่การประยุกต์**, กรุงเทพฯ: สำนักพิมพ์มหาวิทยาลัยเกษตรศาสตร์.
- Aaron, L., Loren, H.R., Robert, P. A., Nhan, D. and Zhu, G. (1990). A Method For Collecting Dried Plant Specimens For DNA And Isozyme Analyses, And The Results Of A Field Test In Xinjiang, China. **Annals of the Missouri Botanical Garden**, 77(4), 859-863.
- Agrawal, G.K., Pandey, R.N. and Agrawal, V.P. (1992). Isolation of DNA from *Cheorospondias asillaris* leaves. **Biotech. Biodiv. Lett**, 2, 19-24.
- Akinjogunla, O.J., Adegoke, A.A., Udokang, I.P., Adebayo-Tayo, B.C. (2009). Antimicrobial potential of *Nymphaea lotus* (Nymphaeaceae) against wound pathogens. **Journal of Medicinal Plants Research**, 3(3), 138-141.
- Bakker, F. T., Culham, A., Daugherty, L. and Gibby, M. (1999). A *trnL-F* phylogeny for species of *Pelargonium* (Geraniaceae) with small chromosomes. **Plant Systematics and Evolution**, 216, 309-324.
- Begum, H.A., Ghosal, K.K. and Chattopadhyay, T.K. (2010). Comparative morphology and floral biology of three species of the genus of *Nymphaea* from Bangladesh. **Bangladesh J. Bot**, 39(2), 179-183.

- Bellstedt, D.U., ZYL, L.V., Marais, E.M., Bytebier, B., Villiers, C.A.D., Makwarela, A.M. and Dreyer, L.L. 2008. Phylogenetic relationships, character evolution and of southern African members of *Zygophyllum* (Zygophyllaceae) based on three plastid regions. **Molecular Phylogenetics and Evolution**, 47(3), 932-949.
- Bhandarkar, M.R. and Khan, A. (2004). Antihepatotoxic effect of *Nymphaea stellata* Willd., against carbon tetrachloride-induced hepatic damage in albino rats. **Journal of Ethnopharmacology**, 91, 61–64.
- Chawanje, C.M., Barbeau, W.E. and Grun, I. (2007). **Nutritional potential of the under-utilized indigenous root tuber *Nymphaea petersiana* in Malawi.** สืบค้นเมื่อ 1 พฤศจิกายน 2554, [http://www.istr.org/Symposiums/Tanzania\\_03/Abstracts/ChawanjeAbs.pdf](http://www.istr.org/Symposiums/Tanzania_03/Abstracts/ChawanjeAbs.pdf).
- Conard, H.S. (1905). **A Monograph of the Genus *Nymphaea***. Washington: Carnegie Institution.
- Diao, Y., Chen, L., Yang, G., Zhou, M., Song, Y., Hu, Z., *et al.* (2006). Nuclear DNA C-values in 12 species in Nymphaeales. **CARYOLOGIA**, 59(1), 25-30.
- Dhanabal, S.P., Mohan Maruga Raja M.K., Ramanathan, M. and Suresh, B. (2007). Hypoglycemic activity of *Nymphaea stellata* leaves ethanolic extract in alloxan induced diabetic rats. **Fitoterapia**, 78(4), 288-291.
- Filatov, D.A. and Charlesworth. (1999). DNA polymorphism, haplotype structure and balancing selection in the *Leavenworthia* PgiC locus. **Genetics**, 153, 1423-1434.
- Grob, V., Moline, P., Pfeifer, E., Novelo, A.R. and Rutishauser, R. (2006). Developmental morphology of branching flowers in *Nymphaea Prolifera*. **J Plant Res**, 119, 561–570.
- Hai-Qing, Y., Chun, Z., Chun-Bang, D., Xiao, M. and Yong-Hong, Z. (2010). Maternal donors of polyploids in *Pseudoroegneria* (Poaceae: Triticeae) and related genera inferred from chloroplast *trnL-F* sequences. **Turk J Biol**, 34, 335-342.
- Hossain, A., Kabir, G., Ud-deen, M. M. and Alam, A. M. S. (2007). Cytological Of *Nymphaea* Species Available In Bangladesh. **J. bio-Sci**, 15, 7-13.
- Huang, T.C. (1972). Pollen Flora of Taiwan. National Taiwan University, **Botany. Dept. Press**, 297.
- Kabir, S.R., Zubair, M.A., Nurujjaman, M., Haque, M.A., Hasan, I., Islam, M.F., *et al.* (2011). Purification and characterization of a Ca<sup>2+</sup>-dependent novel lectin from *Nymphaea nouchali* tuber with antiproliferative activities. **Bioscience Reports Immediate Publication**, Portland Press Limited.

- Khoshshokhan, M., Kazempour Osaloo, S., Saadatmand, S. and Attar, F. (2010). Molecular phylogeny of *Rochelia* (*Boraginaceae*) based on nrDNA ITS and cpDNA *trnL-F* sequences. **Iran. J. Bot**, 16 (1), 22-29.
- Krosnick, S.E. and Freudenstein, J.V. (2005). Monophyly and Floral Character Homology of Old World Passiflora (Subgenus *Decaloba*: Supersection Disemma). **Systematic Botany**, 30(1), 139-152.
- Kumar, S., Tamura, K. and Nei, M. (2004). MEGA3: Integrated software for Molecular Evolutionary Genetics Analysis and Sequence Alignment. **Briefing in Bioinformatics**, 5, 150-163.
- Les, D.H., Moody, M.L. and Doran, A.S. (2004). A Genetically Confirmed Intersubgeneric Hybrid in *Nymphaea* L. (*Nymphaeaceae* Salisb.). **HortScience**, 39(2), 219-222.
- Lohne, C., Borsch, T. and Wiersema, J.H. (2007). Phylogenetic analysis of Nymphaeales using fast-evolving and noncoding chloroplast markers. **Botanical Journal of the Linnean Society**, (154) 141–163.
- Marquina, S., Bonilla-Barbosa, J. and Alvarez, L. (2005). Comparative phytochemical analysis of four Mexican *Nymphaea* species. **Phytochemistry**, 66, 921–927.
- Mauricio, R., Stahl, E.A., Korves, T., Tian, D., Kreitman, M. and Bergelson, J. (2003). Natural selection for polymorphism in the disease resistance gene Rps2 of *Arabidopsis thaliana*. **Genetics**, 163, 735-746.
- Maynard, J.R. and Moseley, F. (1961). Morphological Studies of the Nymphaeaceae II. The Flower of *Nymphaea*. **Bot Gaz**, 122(4), 233-259.
- Muntendam, J.B., Povel, G.D.E. and Van Der Velde, G. (1996). Morphometric patterns in the *Nymphaea alba-candida* complex. **Acta Bot, Neerl.** 45(3), 279-302.
- Nakkuntod, M., Su, Y.C.F., Seelanan, T. and Sanders, R.M.K. (2009). Molecular phylogenetic and morphological evidence for the congeneric status of *Goniothalamus* and *Richella* (*Annonaceae*). **Taxon**, 58, 127-132.
- Nicholas, K.B. and Nicholas, H.J.B. (March 21, 2007). **GeneDoc: a tool for editing and annotating multiple sequence alignment**. Retrieved November 2, 2011, from <http://en.biosoft.net/format/GeneDoc.html>.
- Pathummuang, S. and Bandit, V. (2005). Varietal identification of water lily by isozyme analysis. **Journal of Agriculture (Thailand). Warasan Kaset**, 21(1), 15-25.
- Podoplelova, Y. and Ryzhakov, G. (2005). Phylogenetic analysis of the order Nymphaeales based on the nucleotide sequences of the chloroplast ITS2-4 region. **Plant Science**.169, 606–611.
- Prance, G.A. (1980). A Note on the Pollination of *Nymphaea Amazonum* Mart. & Zucc. (*Nymphaeaceae*). **Brittonia**, 32(4), 505-507.

- Rønsted, N., Chase, M.W., Albach, D.C., and Bello M.A. (2002). Phylogenetic relationships within *Plantago* (Plantaginaceae): evidence from nuclear ribosomal ITS and plastid trnL-F sequence data. **Botanical Journal of the Linnean Society**, 139, 323–338.
- Slocum, P.D. (2005). **Waterlilies and Lotuses**. Portland: Cambridge.
- Songpanich, P., Hongtrakul, V. (2010). Intersubgeneric cross in *Nymphaea* spp. L. to develop a blue hardy waterlily. **Scientia Horticulturae**, 124(4), 475-481.
- Swofford, D.L. (1998). PAUP\*: Phylogenetic analysis using parsimony. **Sinauer Associates**, Sunderland, Massachusetts, USA.
- Taberlet, P., Gielly, L., Pautou, G. and Bouvet, J. (1991). Universal primers for amplification of three non-coding regions of chloroplast DNA. **Plant Molecular Biology**, (17), 1105-1109.
- Taylor, M.L. and Osborn, J.M. (2005). Pollen ontogeny in *Brasenia* (Cabombaceae, Nymphaeales). **American Journal of Botany**, 93, 344-356.
- Thippeswamy, B.S., Mishra, B., Veerapur, V.P. and Gupta, G. (2011). Anxiolytic activity of *Nymphaea alba* Linn. in mice as experimental models of anxiety. **Indian Journal of Pharmacology**, 43(1), 50-55.
- Weiguo, Z., Yile, P., Shihai, Z.Z.J., Xuexia, M. and Yongping, H. (2005). Phylogeny of the genus *Morus* (Urticales: Moraceae) inferred from ITS and *trnL-F* sequences. **African Journal of Biotechnology**, 4(6), 563-569.
- Weir, B.S. (April 6, 2010). Bioprospecting environmental metagenomes - BTech Honours report. **NZ Rhizobia Retrieved** November 3, 2010, from <http://www.rhizobia.co.nz/research/bioprospecting.html>
- Wiersema, J.H. (1988). Reproductive Biology of *Nymphaea* (Nymphaeaceae). **Annals of the Missouri Botanical Garden**, 75, 795-804.
- Woods, K., Hilu, K.W., Borsch, T. and Wiersema, J.H. (2005). Pattern of Variation and Systematics of *Nymphaea odorata*: II. Sequence Information from ITS and *trnL-trnF*. **Systematic Botany**, 30(3), 481–493.