

Fluorometric Response of Photosynthetic Microorganism Consortium as Potential Bioindicator for Heavy Metals Detection in Water

Nurul Afiqah Khishamuddin^{1,2}, Wong Ling Shing¹, Chai Mee Kin^{2*},
Voon Betty Wan Niu²

¹*Faculty of Health and Life Sciences, INTI International University, 71800 Nilai, Malaysia.*

²*College of Engineering, Universiti Tenaga Nasional, 43650 Bangi, Malaysia.*

*Corresponding Email: lingshing.wong@newinti.edu.my / lingshing79@yahoo.com.sg
Received: October 14, 2017; Accepted: November 16, 2017

Abstract

This paper reports the fluorometric responses of photosynthetic microorganism consortium in the detection of heavy metals (Cu and Cd). The consortium was collected from natural water body, and then exposed to different concentration of heavy metals. The response was measured based on the fluorescence signal emitted by the consortium before and after the exposure to heavy metals, with excitation and emission wavelengths set at 526 nm and 648 nm respectively. Cell suspension with optical density (OD) of 0.75 at $\lambda = 700$ nm was found to produce best response. The ability of the consortium to respond to both heavy metals within 0.01 – 10.00 mg L⁻¹ was confirmed. Thus, photosynthetic microorganism consortium could be a good natural candidate as heavy metals bioindicator.

Key words: Photosynthetic Microbes; consortium; fluorometric response; heavy metals; bioindicator

1. Introduction

Heavy metals pollution has become a serious environmental issue, affecting not only the flora and fauna at the surrounding area, but threatening human life as well through food chain and drinking water. Cu and Cd are two

heavy metal pollutants which are commonly found in wastewater discharged from mining, electroplating, smelting and pigmentation industries (Li et al., 2004). Due to the overwhelming pollution issues, sensitive scientific tools have been developed for rapid detection of

pollutants. However, most of these tools, e. g. atomic absorption spectrometry and inductively coupled plasma-mass spectrometer require high operational cost and maintenance. Hence, whole cell algae bioindicator can be used as alternative to screen for the presence of heavy metal pollutants.

The search for new cost effective methods for the detection of heavy metals has been focused on the usage of photosynthetic organisms (Bilitewski et al., 2004). Microalgae are generally low cost, easy to cultivate, present abundantly in aquatic ecology, and they are very sensitive to environmental changes (Torres et al., 2000; Chouteau et al., 2004). These microalgae contain chlorophylls and have the ability to produce oxygen (Beams & Kessel, 1977). The energy received from the sun is diffused through the conversion to heat, channeled to photosynthesis process, and some of the energy are released in the form of fluorescence emission. The presence of heavy metals in microalgae will inhibit the photosynthesis process and causes change in the fluorescence emission (Wong et al., 2013). Hence, fluorescence emission is a suitable parameter to be used in the detection of heavy metals. Up to now, several bioindicators have been successfully constructed using microbes, e.g. yeast (Mastura et al., 2013), algae (Singh et al., 2012) and cyanobacteria (Wong et al., 2013).

To date, the research on the detection of heavy metals using cell-based bioindicator are focused on the specific species of lab-based cultures,

such as microalgae *Spirogyra sp.* (Wong & Kiew, 2016) cyanobacteria *Anabaena cylindrica* (Paul et al., 2016) or plant cell *Daucus carota* (Wong et al., 2015). However, the full potential of using photosynthetic microorganisms, especially in the form of consortium from natural environment has not been explored (Hannon et al., 2010). As the responses of pure species of photosynthetic microbes are well studied, the responses of the consortium of photosynthetic microbes remained unknown. Therefore in this paper, the fluorometric response of chlorophyll in the photosynthetic microorganism consortium to the presence of heavy metals is reported. The outcome of this research might be useful in utilizing the photosynthetic microorganism consortium for the construction of biosensor.

2. Materials and Methods

2.1 Culture Medium and Chemicals

Bold Basic stock medium was purchased from Sigma- Aldrich, Malaysia. The stock medium was diluted 50x with deionized water for culturing work. Jaworski Medium was prepared based on the composition provided by Culture Collection of Algae and Protozoa (CCAP), United Kingdom. Heavy metals were obtained from Merck, Malaysia.

2.2 Microalgae culture and Cell Identification

Algae samples were collected from water sources from Biru Kundang Lake, Selangor (3. 251160, 101.526124). The sample was cultured in combination of Bold Basic Medium and Jawoski Medium (ratio 1: 1) to provide the nutrients required by the microalgae. Aeration was carried out daily by manual shaking. Light and dark conditions were at 16 hours and 8 hours respectively. The genus of the algae and cyanobacteria were identified through light microscope (Eclipse E-100 LED, Nikon).

2.3 Standardization of the Number of Cell

The standardization of algae consortium cells were done by spectrophotometer (GeneQuant 100, GE) with $OD_{700nm} = 0.75$. The consortium with $OD_{700nm} = 0.75$ contained optimum density of cells for the fluorescence study, which was experimentally determined. The intensity of fluorescence emission were captured using emission and excitation wavelengths sets at 526 nm and 648 nm respectively using spectrofluorometer (Glomax Multi Jr., Promega).

2.4 Heavy Metals Exposure

Cd solutions were added to 2 mL of consortium culture in four sided clear cuvettes to make the final concentrations of Cd in the algae consortium culture 0.01 mgL^{-1} , 0.05 mgL^{-1} , 0.10 mgL^{-1} , 0.50 mgL^{-1} , 1.00 mgL^{-1} , 5.00 mgL^{-1} , and 10.00 mgL^{-1} respectively. The intensity of the fluorescence emission was measured

before the exposure to Cd, and then measured again after 30 minutes, 60 minutes, 120 minutes, 240 minutes, 360 minutes, and 480 minutes of exposure. The experiment was then repeated with Cu replacing Cd. All exposure tests were conducted in triplicates. The percentage of changes of the intensity of fluorescence emission was calculated using an equation below:

Increase in fluorescence emission (%)

$$= (f_1 - f_0) \times 100\% / (f_0);$$

Where :

f_0 = fluorescence emission before the exposure;

f_1 = fluorescence emission after the exposure

3. Results and Discussion

3.1 Microalgae Identification

Co-culture or consortium had the advantages in certain applications which were impossible to be done by an individual strain or species (Brenner et al. , 2008) . Potential of photosynthetic microorganism consortium as the bio- indicator have been evaluated and it showed that they were able to detected present of heavy metals through the production of fluorescence emission. The consortium contained *Chlorella* sp. , *Haematococcus* sp., cyanobacteria and diatoms.

3.2 Effects of copper and cadmium on photosynthetic microorganism

The density of cell can be determined by the reading of OD at 700 nm, which is the wavelength for absorption of chlorophyll *a* (Desikachary, 1959). The correlation between the optical densities of consortium and fluorescence emission is portrayed in Figure 1. OD has been used to determine the number of fluorescence bacteria (Nawaz *et al.*, 2011), and cyanobacteria (Wong *et al.*, 2013). From the experiment, fluorescence increased from 0.10 A – 0.60 A. However, the intensity started to decreased when the increment of consortium culture exceeded 0.75 A, due to the reabsorption of fluorescence emitted by nearby cells (Védrine *et al.*, 2003). Based on the result, the consortium culture at OD_{700nm} = 0.75 yielded the highest fluorescence intensity, thus had been used for the exposure tests.

The fluorometric response of consortium to seven different concentrations of Cd is shown in

Figure 2. Cd at 0.05 mgL⁻¹ yielded the highest fluorescence intensity compared to other Cd concentrations. However, low fluorescence intensity was observed at concentration of 5 mgL⁻¹ and 10 mgL⁻¹. Lower fluorescence intensity yielded at the higher concentrations of Cd might due to disturbance of metabolic activities of the photosynthetic microbes. In photosynthetic organisms, Cd affected metabolic activities such as photosynthesis in both PS I and PS II (Atal *et al.*, 1991). The effect might due to the difference way of reaction of microbes reacted to the heavy metal (Gadd *et al.*, 2007; Gadd *et al.*, 2009).

The responses of the cells to Cu is shown in Figure 3. The concentration of Cu at 0.05 mgL⁻¹ and 0.50 mgL⁻¹ yielded the highest fluorescence intensity. Cu is an essential element that is required by the metabolic processes of algae (Campanella *et al.*, 2001).




Fig 1. The correlation between OD and fluorescence intensity




Fig 2. Exposure of consortium to different concentration of Cd. The changes of fluorescence intensity were significant ($p < 0.05$) compared to negative control.




Fig 3. Exposure of consortium to different concentration of Cu. The changes of fluorescence intensity were significant ($p < 0.05$) compared to negative control.

However, the metal would exhibit toxicity effect in high concentration (Pandard *et al.*, 1993).

4. Conclusion

The result from this study confirmed that the algae consortium responded to the presence of heavy metals through fluorescence emission. Although the test showed different responses, the consortium produced detectable emission signal to indicate the disturbance by Cu and Cd to the consortium culture, which enabled qualitative detection of the heavy metals. This study showed the consortium could detect the presence of heavy metals even at concentration as low as 0.05mgL^{-1} within 30 minutes of exposure.

5. Acknowledgement

The authors thanked Universiti Tenaga Nasional, Malaysia (grant no: UNIIG J510050639) and INTI International University, Malaysia (grant no: INTI-FHLS-02-05-2016) for the financial support on the project.

References

- Atal, N., Saradhi, P. P., & Mohanty, P. Inhibition of the Chloroplast Photochemical Reactions by Treatment of Wheat Seedlings with Low Concentrations of Cadmium: Analysis of Electron Transport Activities and Changes in Fluorescence Yield. *Plant and Cell Physiology* 1991; 32(7): 943-951.
- Beams, H.W & Kessel, R.G. physical properties of blue-green algae: ultracentrifugation and electron microscope studies. *Protoplasma* 1977; 92: 229-242.
- Brenner, K., You, L., & Arnold, F. H. Engineering microbial consortia: a new frontier in synthetic biology. *Trends in Biotechnology* 2008; 26(9): 483-489.
- Bilitewski, U and Turner, A. *Biosensors in environmental monitoring*: Taylor & Francis 2004.
- Campanella, L., Cubadda, F., Sammartino, M.P., Saoncella, A. An algal biosensor for the monitoring of water toxicity in estuarine environments. *Water Research* 2001; 35(1): 69-76.
- Chouteau, C., Dzyadevych, S., Chovelon, J.M., Durieu, C. Development of novel conductometric biosensors based on immobilized whole cell *Chlorella vulgaris* microalgae. *Biosensors and Bioelectronics* 2004; 19: 1089-1096.
- Desikachary, M. *Cyanophyta*. Indian Council of Agricultural Research 1959.
- Gadd, G. M. Metals, minerals and microbes: geomicrobiology and bioremediation. *Microbiology* 2009; 156(3): 609-643.
- Gadd, G. M., Burford, E. P., Fomina, M., & Melville, K. Mineral transformations and biogeochemical cycles: a geomycological perspective. *Fungi in the Environment* 2007; 77-111.
- Hannon, M., Gimpel, J., Tran, M., Rasal, B., Mayfield, S. Biofuels from algae: challenges and potential. *National Institute of Health* 2011; 1(5): 763-784.
- Li, T.Y. & Ziong, Z.T. a novel response of wild-type duckweed (*Lemna paucicostata* Hegelm.) to heavy metals. *Environ. Toxicol* 2004; 19(2): 95-102.
- Mastura, H., Yamamoto, Y, Muraoka, M., Akaishi, K., Hori, Y., Uemura, K., Tsuji, N., Harada, K., Hirata, K., Bamba, T., Miyasaka, H., Kuroda, K., Ueda, M., Development of surface-engineered yeast cells displaying phytochelatin synthase and their application to cadmium biosensor by the combined use of pyrene-excimer fluorescence. *Biotechnology Progress* 2013; 29(5): 1197-202.
- Nawaz, A., & Ahmed, N. Isolation and characterization of indigenous

- luminescent marine bacteria from Karachi coast. Academic Research International 2011; 1(2): 74-83.
- Pagliano, C., Raviolo, M., Vecchia, F. D., Gabbrielli, R., Gonnelli, C., Rascio, N., Rocca, N. L. Evidence for PSII donor-side damage and photoinhibition induced by cadmium treatment on rice (*Oryza sativa L.*). Journal of Photochemistry and Photobiology B: Biology 2006; 84(1): 70-78.
- Pandard, P., Vasseur, P., Rawson, D.M. Comparison of two types of sensors using eukaryotic algae to monitor pollution of aquatic systems. Water Research 1993; 27: 427-431.
- Paul, S., Wong, L. S., & Ong, G. H. The responses of carotenoids in cyanobacteria *Anabaena cylindrica* as potential bio-indicator for heavy metals detection. Pollution Research 2016; 35(2) 259-262.
- Pinto, E., Sigaud-Kutner, T. C., Leitaõ, M. A., & Okamoto, O. K. Heavy metal-induced oxidative stress in algae. Journal of Phycology 2003; 39(6): 1008-1018.
- Wong, L. S., Lee, Y. H. & Surif, S. Whole cell biosensor using *Anabaena torulosa* with optical transduction for environmental toxicity evaluation. Journal of Sensor 2013; 1-8.
- Wong, L. S., Lee, Y. H. & Surif, S. Performance of a cyanobacteria whole cell-based fluorescence biosensor for heavy metal and pesticide detection. Sensors 2013; (13): 6394-4004.
- Wong, L.S. The interference of bioenergetics in photosynthesis and the detection of heavy metals. Bioenergetics: Open Access 2013; 2(2): 114.
- Wong, L. S., Lee, B. R., Koh, C. E., Ong, Y. Q., & Choong, C. C. A novel in vivo β carotene biosensor for heavy metals detection. Journal of Environmental Biology 2015; 36: 1277-1281.
- Wong, L. S., & Kiew, W. Y. The Detection of Calcium and Sodium using Green Algae *Spirogyra*. Transactions on Science and Technology 2016; 3(1): 8-12.
- Singh, J. and Mittal, S. K. *Chlorella sp.* Based biosensor for selective determination of mercury in presence of silver ions. Sensors and Actuators B: Chemical 2012; 165(1): 48-52.
- Torres, E., Cid, A., Herrero, C., Abalde, J. Effect of cadmium on growth, ATP content, carbon fixation and ultra-structure in the marine diatom *Phaedactylum tricornutum Bohlin*. Water, Air, Soil Pollution 2000; 117: 1-14.
- Védrine, C., Leclerc, J., Durrieu, C., & Tran-Minh, C. Optical whole-cell biosensor using *Chlorella vulgaris* designed for monitoring herbicides. Biosensors and Bioelectronics 2003; 18(4): 457-463.