

Human Health Risk Assessment for Inhabitants of Four Towns of Rajshahi, Bangladesh due to Arsenic, Cadmium and Lead Exposure

Nazmul Islam^{1,*}, Entazul M. Huq¹, Nurul Islam¹, Asadul Islam¹, Rustom Ali¹,
Rehena Khatun¹, Niamul Haque², Rafiqul Islam³, Sultana Nasima Akhter⁴

¹*Department of Chemistry, University of Rajshahi, Rajshahi, Bangladesh*

²*Department of Marine Environmental Engineering, Gyeongsang National University, Tongyeong, South Korea*

³*Department of Pediatrics, Z. H. Sikder Women's Medical College, Dhaka, Bangladesh*

⁴*Institute of Biological Sciences, University of Rajshahi, Rajshahi, Bangladesh*

* Corresponding author: dr.m.nazmul.i@gmail.com
Received: November 30, 2017; Accepted: December 30, 2017

Abstract

We performed a risk assessment for inhabitants of four towns of Rajshahi, Bangladesh beside the silt (toxic metal enriched) carrying river Padma due to arsenic, cadmium and lead exposures from consumption of rice, vegetables, fish, drinking water and dust. As, Cd and Pb concentrations in the samples (n= 300) ranged 0.002-3.441, 0.001-0.017 and 0.001-0.398 $\mu\text{g g}^{-1}$ (ww) respectively. Since $P < 0.05$, Levene and Kruskal-Wallis tests revealed that there were statistically significant differences amongst both standard deviations and medians of the data at 95% confidence level respectively. Generally concentrations of the concerned metals in the samples followed the order: Dust>Rice>Fish>Vegetables>Drinking water. Bangladeshi consumption rate showed that rice alone imposed most of As and Pb load, whereas drinking water most of Cd load. We utilized USEPA risk assessment model and considered potential exposure pathways (diet and non-diet ingestion, and dust inhalation). Ingestion accounted for 95.0-99.9% exposure, whereas inhalation only 0.1-5.0% exposure. Risk, defined as 'Hazard Index' (HI), was mapped for three population sub-groups: *Highly Exposed Child*, *Average Person* and *Senior*. The results showed that all children (0-6 years) were at health risk, whereas 70% of seniors and 60% of average persons (i.e., adults) were safe in consuming the foods. The contribution of the metals in total exposure was

As(69%)>Cd(21%)>Pb(10%). This study highlights the importance of sight-specific human health risk assessment considering more pollutant parameters.

Keywords: human health; risk assessment; arsenic, lead, cadmium, exposure

1. Introduction

British Geological Survey (BGS) in 2001 showed that the groundwater (n=3,540) of shallow aquifer, used for both drinking and irrigation purposes, in 60 out of 64 districts of Bangladesh was contaminated with arsenic (BGS/ DPHE, 2001) . In Chapai Nawabganj district the highest concentration of arsenic in tube-well water was found as 2,400 $\mu\text{g L}^{-1}$ (BGS/DPHE, 2001) , whereas WHO guideline value for drinking water was 50 $\mu\text{g L}^{-1}$ for Bangladesh and 10 $\mu\text{g L}^{-1}$ for many other countries (WHO, 2017). An estimated 24 million people are directly exposed to this contamination and another 75 million are at risk (Engconsult Limited, 2017; WHO, 2016a) . Nearly 10,000 arsenicosis patients had been identified and a few deaths due to arsenic related diseases had also been reported (Engconsult Limited, 2017; Fazal et al 2001; SOES/DCH, 2000). Clearly Bangladesh is facing the probably the largest mass poisoning in history. According to BGS (2001) and, Saha and Ali (2007), the groundwater of Bangladesh was moderately contaminated with lead and least contaminated with cadmium.

The area under irrigation has been increased significantly over the last fifteen years in Bangladesh to raise food production mainly through

installation of shallow tube-wells (BBS, 2017). Islam and his co-authors (2012) found that vegetables were good accumulators of arsenic and arsenic accumulated in vegetables exponentially due to successive harvesting. Das et al. (2004) investigated arsenic concentrations in rice, vegetables and fishes in Bangladesh and found that some of these exceeded some food safety legislation limit (Table 4). Saha and Zaman (2011) investigated the cereals such as rice, wheat and urid beans of Shibganj, Chapai Nawabganj, Bangladesh and found high lead contents ranging 7.31-12.33 $\mu\text{g g}^{-1}$ (dw) . All samples exceeded FAO/ WHO CODEX recommended maximum permissible limit of Pb in foodstuff which was 0.2 $\mu\text{g g}^{-1}$ (FAO/ WHO, 2009b) . Obviously, increased use of arsenic and other metal contaminated groundwater for irrigation purpose suggests that consumption of the irrigated crops and vegetables could be another major exposure route of arsenic and heavy metals in humans in Bangladesh.

There are two major processes by which a chemical can cross the boundary from outside to inside the human body (USEPA, 2016, 1992) . The first process is intake that involves physically moving the chemical through an opening in the outer boundary (usually the mouth or

nose), typically via inhalation, eating, or drinking (Fig. 1) . The second process is uptake that involves absorption of the chemical through skin (Fig 1) , where mass transfer occurs by diffusion. Irrespective of exposure pathways, arsenic, cadmium and lead are cumulative poisons that do not metabolized in other intermediate compounds easily and do not easily break down in environment (Adal, 2017) . Arsenic, defined as Group 1 carcinogen by International Agency for Research on Cancer (IARC, 2012a), causes acute lethality to chronic effects. Chronic arsenic exposure leads to vascular diseases, hypertension, cancer, hyperpigmentation, genotoxicity, diabetes mellitus, repeated abortions, stillbirth, preeclampsia, etc. (USEPA, 2001; ATSDR, 2013; WHO, 2016) . Cadmium is a possible human carcinogen of class B1 (IARC, 2012b). Long-term cadmium exposure

leads to cancer, kidney dysfunction, bone fracture, genotoxicity, etc. (IARC, 2012b; WHO, 2010) . Lead exposure also causes anaemia, hypertension, renal impairment, immunotoxicity, dysfunction of central nervous system and brain, etc. (ATSDR, 2007; IARC, 2006; WHO, 2016b).

2. Materials and Methods

2.1 Study Area

The study area is situated at the northwestern part of Bangladesh, namely, Rajshahi Sadar Upazilla of Rajshahi district, which is under 12 number Ward of Rajshahi City Corporation (RCC) authority. It is located between 24°19'N to 24°22'N and 88°35'E to 88°36'E that covers an area of about 10.5 sq. kilometer (Fig 2). The studied four urban Towns were Rani Bazar, Shaheb Bazar,




Figure 1. Exposure pathways of heavy metals to human beings in nature.

Futhkipara and Kumarpara. The reason for selecting the towns is that the river Padma (also called Ganges) flows adjacent to these Towns that carries a huge amount of silt from India every year as Himalayan drain, a fraction of which contains toxic heavy metals such as Cr, Pb, etc. (Zereen et al., 2000).

2.2 Sampling of Diets and Non-diet

For sampling five houses were selected in random from each town, comprising a total of 20 houses. The five samples that were collected from each house included rice, vegetables, fish, drinking water and dust. The owners of the houses used to collect rice, vegetables and fishes (of the river Padma) from their near-by market Shaheb Bazar, but they fetched drinking water from their adjacent tube-wells. The collected vegetables were fresh and the fishes were kept refrigerated at -20°C until frozen. In order to get aquifer water the tube-wells were flushed with 2-3 tube-well

volumes of groundwater. The water was collected in polyethylene bottles and acidified with conc. HNO_3 to keep pH below 2 (APHA, 2017). The dust samples were collected mainly by sweeping the shelves and floor where they stayed most of the time. At laboratory, the vegetables and fish samples were washed with distilled deionized water (DDW), cut into pieces and dried in an oven at $80-100^{\circ}\text{C}$ for 24 h. Rice was dried in a similar manner without washing. The dried materials were subjected to grinding with ceramic mortar and pestle followed by sieving with 0.5 mm screen. The dust sample after drying at 105°C for 48 h was ground and sieved as before.

2.3 Extraction and Estimation of the Metals

For extraction of As, Cd and Pb, the water samples were subjected to mild digestion with HNO_3 (APHA, 2017), dust samples were digested by modified method of Small




Figure 2. Study area for risk assessment. [Sampling spots are marked by \odot]

and *McCants* with H₂SO₄-HClO₄ (volume ratio of 2 : 3) (Sakamoto et al, 2001; Hesse, 1994; Shimadzu Cookbook) and, vegetables, rice and fish samples were digested through *Wet Oxidation* by HNO₃-H₂SO₄-HClO₄ (volume ratio of 10: 1: 4) (Hesse, 1994; Jackson, 1958; Shimadzu Cookbook). Prior to wet oxidation the samples were pre-digested with 69% (v/v) HNO₃. The digestates, after dilution with 6M HCl and then DDW, were analyzed for total concentrations of As, Cd and Pb by GF-AAS using a Shimadzu AA-6800 (Shimadzu Corporation, Kyoto, Japan) atomic absorption spectrophotometer. It was equipped with an auto-sampler (ASC-6100, Shimadzu) and a graphite furnace (GFA-EX7, Shimadzu), and operated through 'WizAard' software. Certified reference materials (CRM) of As, Cd and Pb standard solutions for AAS (TraceCERT®) (Fluka, Switzerland) were used for calibration purposes. The minimum detection limits of As, Cd and Pb were 0.3, 0.02 and 0.15 µg L⁻¹ respectively.

2.4 Model Development

For estimating total exposure of As, Cd and Pb for the consumers of the studied samples, we first estimate Average Daily Dose (ADD, mg kg⁻¹ d⁻¹). Individual ADDs of arsenic contributed by each of rice, vegetables, fish, drinking water and soil were estimated utilizing the following equation as suggested by U. S. EPA Guidelines for Human

Exposure Assessment (USEPA, 2016; USEPA, 1992):

$${}^{As}ADD_{Diet/Non\ diet} = \frac{{}^{As}C_{Diet/Non\ diet} \cdot IR \cdot ED}{BW \cdot AT} \quad (1)$$

where ${}^{As}C_{Diet/Non-diet}$ is the concentration (mg kg⁻¹, ww) of arsenic in a particular dietary item or soil, *IR* is ingestion rate (kg day⁻¹, ww) for the diet/non-diet, *ED* is exposure duration (yr), *BW* is the body weight (kg) and *AT* is averaging time (yr) (Table 1). We considered bio-accumulation by multiplying equation (1) by relative bio-availability factor, *RBAF*. ${}^{As}ADD_{Ingestion}$ was then estimated as follows:

$${}^{As}ADD_{Ingestion} = \sum {}^{As}ADD_{Diet} + {}^{As}ADD_{Non-diet} \quad (2)$$

$${}^{As}ADD_{Ingestion} = ({}^{As}ADD_{Rice} + {}^{As}ADD_{Veg} + {}^{As}ADD_{Fish} + {}^{As}ADD_{Water}) + {}^{As}ADD_{Soil} \quad (3)$$

The term ${}^{As}ADD_{Soil}$ is included as we passively eat some soil adhered to foods, e.g., vegetables (Hough, 2004; USEPA, 2011). Similarly, ${}^{Pb}ADD_{Ingestion}$ and ${}^{Cd}ADD_{Ingestion}$ were calculated.

For dust inhalation, first concentration of inhaled arsenic, for example, in indoor air was estimated from existing indoor dust using the following equation (USEPA, 2011):

$$[As]_{air} = D \times M_{PM} \times F_{PM} \times F_{count} \quad (4)$$

where, *D* is the concentration of dust in the air [assumed the standard value of 70.0 µg/m³ (USEPA, 2011)], *M_{PM}* is the arsenic concentration on airborne particulate matter (assumed

equal to $[As]_{dust}$ where dust is derived from the soil), F_{PM} is the proportion of particulate matter that is respirable [assumed $PM_{10} \sim 73\%$ as 27% of airborne particles are generally found to be $> 10 \mu m$ in equivalent diameter] and F_{cont} is the fraction of dust that is derived from the contaminated source. For indoors, $F_{cont} = 0.445[M_{soil}]$ because 44.5% of indoor dust is considered to be derived from outdoor soil. This yielded

$$[As]_{air} = (70.0 \mu g/m^3) \times [As]_{dust} (mg/kg) \times 0.27 \times 0.445$$

Or, $[As]_{air} = (0.070 mg/m^3) \times [As]_{dust} \times 0.27 \times 0.445 \mu g/m^3$

Dust inhalation model was the same as equation (1) from where $^{As}ADD_{Inhalation}$ was obtained (USEPA, 2011). Similar calculations were made for Cd and Pb.

Considering the potential routes of entry of the metal into human body, the term $^{As}ADD_{Total}$ was defined as: $^{As}ADD_{Total} = ^{As}ADD_{Ingestion} + ^{As}ADD_{Inhalation}$. Similarly $^{Cd}ADD_{Total}$ and $^{Pb}ADD_{Total}$ were calculated. We ignored dermal adsorption as its contribution to total ADD is extremely small (USEPA, 2001). The parameters involved in ADD estimation are summarized in Table 1. These were adapted from the data of Exposure Factor Handbook (USEPA, 2011), US EPA standard values (USEPA, 2002) and Bangladesh surveys and standards (BBS, 2010). ADD were estimated for three population subgroups: ‘Highly Exposed Child’, ‘Average Person’ and ‘Senior’. The children of 0-6 years age are often referred to as ‘High Exposed Individuals (HEIs)’.

Table 1. Parameters involved in the risk assessment model.

Factors	Unit	Population sub-group		
		Children (0-6 yr)	Average Person	Senior
Age	yr	6	39.5	60.3
Body weight (BW)	kg	18.6	80.8	82.6
Rice ingestion rate	g (ww.) d ⁻¹	100	344	300
Vegetable ingestion rate	g (ww.) d ⁻¹	60	155	120
Fish ingestion rate	g (ww.) d ⁻¹	40	60	60
Water ingestion rate	L d ⁻¹	1.4	2.5	2
Soil and dust ingestion rate	mg (ww.) d ⁻¹	100	50	50
Dust inhalation rate	m ³ d ⁻¹	10.1	16	15.7
Exposure frequency (EF)	d yr ⁻¹	350	350	350
Year	d	365.25	365.25	365.25
Averaging time (AT)	yr	6	30	50
Exposure duration (ED)	yr	6	30	50
Relative Bio-availability Factor (RBAF)	(unit less)	As: 0.4, Cd: 0.6, Pb: 0.6		

Table 2. Metal reference dose (RfD), experimental dose and provisional tolerable weekly intake (PTWI).

Metal	Experimental dose (mg kg ⁻¹ d ⁻¹)	RfD (mg kg ⁻¹ d ⁻¹)			FAO/WHO PTWI (µg/kg BW)
		Children	Adults	Senior	
As	NOAEL: 8.0×10 ⁻⁴ LOAEL: 0.014	3.0×10 ⁻⁴	3.0×10 ⁻⁴	3.0×10 ⁻⁴	15
Cd	NOAEL (food): 0.01 LOAEL: N/A	5.0×10 ⁻⁴	3.0×10 ⁻²	1.0×10 ⁻³	7
Pb	NOAEL: 25 µg/dL LOAEL: 0.02	5.0×10 ⁻⁴	5.0×10 ⁻⁴	5.0×10 ⁻⁴	25

2.5 Risk Characterization

Risk may be characterized using a hazard quotient (HQ). This is the ratio of ADD of a chemical to a reference dose (RfD) defined as the maximum tolerable daily intake of a specific metal that does not result in any deleterious health effects. The RfDs, may be derived from a NOAEL (No Observed Adverse Effect Level) or a LOAEL (Low Observed Adverse Effect Level), were adapted from US EPA’s Health Effects Notebook and Exposure Factors Handbook (USEPA, 2011) and California EPA’s reference dose values (CalEPA, 2009) (Table 2). For arsenic,

$${}^{As}HQ_{Total} = \frac{{}^{As}ADD_{Total}}{RfD} \quad (5)$$

Similarly ${}^{Cd}HQ_{Total}$ and ${}^{Pb}HQ_{Total}$ were calculated. Then HI were estimated using the equation

$$HI = \sum M HQ_{Total} = {}^{As}HQ_{Total} + {}^{Cd}HQ_{Total} + {}^{Pb}HQ_{Total} \quad (6)$$

where M = As, Cd or Pb. Finally maps of HI were constructed for the three population sub-groups.

2.6 Quality Assurance

We followed USEPA approved QA/QC plan with a reagent blank, a duplicate and a spike for every 20 samples. All glassware was kept in 10% (v/v) HNO₃ for 48 h and rinsed with DDW several times before use. Certified reference materials (CRM) of As, Cd and Pb standard solutions for AAS (TraceCERT[®]) (Fluka, Switzerland) were used for calibration purposes. After analyzing every 10 samples, readings of standard solutions were recorded to check the instrument. Two certified reference materials [Tomato Leaves (SRM 1573a) and Lake Sediment (NIES CRM No. 31)] were digested and analyzed in five replicates for total As, Cd and Pb concentrations under the identical experimental conditions. The overall agreement between the certified and the observed results were in the range of 91-104% (Table 3).

2.7 Statistical Analyses

In our investigation each concentration value corresponds to an

average of triplicate measurements. The datasets were treated separately for analyzing basic statistical parameters and for making cross-tabulations and cross-plots. The SPSS (version 20.0) and STATGRAPHICS Centurion (version 18.1.01) statistical software package, and Microsoft Excel (version 12.0.4518.1014) were employed for the purpose. ANOVA and Levene's Variance test were employed for analyses of variance of the data (Table 5). Kruskal-Wallis and Mood's median tests were also employed for analyses of median of the data.

3. Results and Discussion

3.1 Distribution of the Metals in Studied Diets and Non-diet

The statistical analyses of the observed concentrations of arsenic, cadmium and lead in the studied five types of samples, namely, rice, vegetables, fish, drinking water and dust (n=300) are presented in Table 4. The arsenic, cadmium and lead concentrations in the studied dietary samples ranged 0.002-0.527, 0.0003-0.0170 and 0.0002-0.0631 $\mu\text{g g}^{-1}$ (ww) respectively. Our observed concentrations of arsenic, cadmium and lead in the dietary samples are in accordance with many authors (Williams et al, 2006; Alam et al, 2003). It was found that the tube-well water used for drinking purpose were all safe from arsenic and lead contamination, but all were unsafe of cadmium contamination with respect to WHO guideline values (0.010,

0.003 and 0.010 mg L^{-1} respectively for As, Cd and Pb (WHO, 2017). This is probably due to the fact that the studied urban area does not fall into 'hot-spots' of arsenic, where the concentrations of As and other metals are very high (Charlet et al, 2007; BGS/DPHE, 2001). Generally the concentrations of the concerned metals in the studied samples follow the order: Dust > Rice > Fish > Vegetables > Drinking water. The ANOVA table (Table 5) decomposes the variance of the data (n= 100 for each of As, Cd and Pb) into two components: a between-group component and a within a within-group component. Since P values of the F-tests were less than 0.05, there was a statistically significant difference between the means of the variables. Levene's test showed a statistically significant difference amongst the standard deviations at 95% confidence level. Kruskal-Wallis test revealed that there was a statistically significant difference amongst the medians at 95% confidence level, as P values were less than 0.05. The same observations were found in Mood's median test, as P values for the chi-square test were less than 0.05.

3.2 Arsenic, Cadmium and Lead Load

The Joint Food and Agriculture Organization (FAO) / World Health Organization (WHO) Expert Committee on Food Additives (JECFA) (FAO/WHO, 2011, 2009) has set Provisional Tolerable Weekly Intake (PTWI)

Table 3. Analysis of certified reference materials for total arsenic, cadmium and lead.

Sample	Metal	Certified values ($\mu\text{g g}^{-1}$)	Measured values ^a ($\mu\text{g g}^{-1}$)	% of recovery	n ^b
Tomato Leaves (SRM 1573a)	As	0.112 ± 0.004	0.117 ± 0.009	104	5
	Cd	1.52 ± 0.04	1.44 ± 0.12	95	5
	Pb	/	/	/	/
Lake Sediment (NIES CRM No. 31)	As	13.9 ± 1.5	13.5 ± 1.3	97	5
	Cd	0.342 ± 0.043	0.341 ± 0.045	100	5
	Pb	22.0 ± 3.0	20.0 ± 2.6	91	5

^a the samples were analyzed by GF-AAS; ^b No. of measurements.

values for various metals/contaminants, below the dose would not cause any deleterious health effect. The PTWI values for arsenic, cadmium and lead are 15, 7 and 25 $\mu\text{g}/\text{kg}$ body weight respectively. The weekly total load of As, Cd and Pb due to ingestion of the dietary foods at Bangladeshi rate for the three population sub-groups is represented in Fig 3. Obviously rice alone imposes most of the arsenic and lead load, whereas drinking water imposes most of the cadmium load. This suggests that rice (staple food in Bangladesh and many countries of Asia) intake should be reduced that might be compensated by foodstuffs containing more gluten, e.g. wheat. Because it was reported (Bulka et al., 2007) that the diets containing less gluten might be exposed to higher levels of arsenic and mercury.

3.3 Exposure Pathways

Among the ingestion and inhalation processes, dust driven air

was included in inhalation and the others were in ingestion processes. It was found that almost all (99.9%) of cadmium and lead, and 96.6% of arsenic exposure came from ingestion processes (Table 6). On an average, 98.84% exposure is contributed from ingestion processes, while the rest (1.16%) from inhalation process.

3.4 Metalwise Exposure

The hazard index (HI), which is the sum of hazard quotients, is a screening risk assessment technique commonly used to judge whether there is concern for additive effects between chemicals. If the hazard index is less than unity (1.00), it indicates that even if all the metals acted on the same organ or interacted in some other way to cause health effects, the risk of these effects would be low. Since different pollutants may cause similar adverse health effects, USEPA suggests combining hazard quotients associated with different substances. In the investigation we summed up

Table 4. Statistical analyses of arsenic, cadmium and lead data sets of the studied samples.

Sample	Metal	Concentration of metal (mg kg ⁻¹ , ww)						
		Range (Min -Max)	Mean ± SE	95% CI for mean	Median	Standard Deviation	Variance	Skewness
Rice	As	0.045 - 0.276	0.077±0.011	0.054-0.099	0.065	0.049	0.002	4.009
	Cd	0.0003-0.0170	0.007±0.001	0.004-0.010	0.008	0.006	0.000	0.128
	Pb	0.0183-0.0631	0.040±0.003	0.033-0.047	0.045	0.015	0.000	-0.203
Vegetables	As	0.007 - 0.119	0.034±0.006	0.022-0.046	0.033	0.025	0.001	2.103
	Cd	0.0004-0.0033	0.002±0.000	0.001-0.002	0.001	0.001	0.000	0.577
	Pb	0.0020-0.0299	0.007±0.001	0.004-0.010	0.005	0.007	0.000	2.443
Fish	As	0.044 - 0.527	0.144±0.022	0.097-0.190	0.125	0.100	0.010	3.194
	Cd	0.0012-0.0066	0.003±0.000	0.003-0.004	0.003	0.002	0.000	1.013
	Pb	0.0017-0.0407	0.015±0.002	0.011-0.019	0.014	0.009	0.000	1.892
D. water	As	0.002 - 0.004	0.003±0.000	0.002-0.003	0.003	0.001	0.000	0.241
	Cd	0.0037-0.0070	0.005±0.000	0.005-0.006	0.005	0.001	0.000	0.282
	Pb	0.0002-0.0041	0.002±0.000	0.001-0.002	0.001	0.001	0.000	1.251
Dust	As	0.702 - 3.441	1.511±0.172	1.151-1.87	1.150	0.768	0.591	1.236
	Cd	0.0003-0.0132	0.003±0.001	0.001-0.004	0.001	0.003	0.000	1.768
	Pb	0.0065-0.3983	0.044±0.019	0.004-0.083	0.027	0.084	0.007	4.353

^a the guideline values (GV) were according to WHO, FAO/WHO, Chinese and EC regulations; ^b inorganic As; ^c 0.05 in Bangladesh.

Table 5. ANOVA Table for the distribution of the metals in studied samples.

Source	Sum of Squares	Df	Mean Square	F-Ratio	P-Value
Arsenic: Between groups	33.7121	4	8.42803	69.82	0.0000
Arsenic: Within groups	11.4674	95	0.12071		
Cadmium: Between groups	0.000382826	4	0.0000957064	10.28	0.0000
Cadmium: Within groups	0.000884508	95	0.0000093106		
Lead: Between groups	0.029872	4	0.00746793	5.05	0.0010
Lead: Within groups	0.140587	95	0.00147986		

total HQ for all the metals concerned (i.e., $HI = \sum^M HQ_{Total} = {}^{As}HQ_{Total} + {}^{Cd}HQ_{Total} + {}^{Pb}HQ_{Total}$).

The estimated average total HQ values (${}^M HQ_{Total}$) of arsenic, cadmium and lead for the three population sub-groups are represented in Table 7. It is evident that in total exposure the contribution of the metals for all population subgroups follow the order: As (69%) > Cd (21) > Pb (10). Obviously arsenic alone averagely

contributes roughly one-third of total HI. The HQ contributions for specific diets towards HI are in accordance with Fig 3.

3.5 Risk Assessment

Risk assessment strategies are often aimed at population sub-groups. It is common practice to identify vulnerable people in society, such young children or elderly, and assess

potential risk to the health of these population sub-groups (GRCIRAT, 1999). Hough et al (2004) considered young children to be ‘Highly Exposed Individuals (HEIs) ’ . Thus risk assessment can usually focus on highly exposed subpopulations on the

basis that if the risk to the HEI is acceptable then most of the population is protected.

The estimated HI values were utilized to generate HI indexed maps for three population sub-groups (Fig 4) showing clearly where the




Figure 3. Groupwise weekly total load of arsenic, cadmium and lead for consuming the diets. [WHO guideline values (GV) are based on mg/kg body weight (FAO/WHO 2011)]

Table 6. Exposure pathwaywise contribution for three population sub-groups.

Population Sub-groups	Exposure Contribution (%)		
	Metal	Ingestion	Inhalation
Highly Exposed Child	As	99.28	0.72
	Cd	99.98	0.02
	Pb	99.94	0.06
Average person	As	95.55	4.45
	Cd	99.85	0.15
	Pb	99.98	0.02
Senior	As	95.03	4.97
	Cd	99.98	0.02
	Pb	99.96	0.04
Overall Average	As	96.62	3.38
	Cd	99.93	0.07
	Pb	99.96	0.04

Table 7. Metal wise average HQ contributions to HI values.

Population Sub-groups	Metal specific contribution of HQ to HI value (%)			
	AsHQ _{Total}	CdHQ _{Total}	PbHQ _{Total}	HI
Highly Exposed Child	59.88	21.60	18.52	100.00
Average Person	76.97	22.75	0.28	100.00
Senior	69.37	18.82	11.81	100.00
Overall Average	68.74	21.06	10.20	100.00




Figure 4. Map of Hazard Index (HI) values for inhabitants of four towns of RCC. [A: Children, B: Average Person, C: Senior]

greatest potential risks were located. The map (Fig 4 C) reveals that 70% of the *Senior* population sub-group was safe from consumption of the dietary foods as well as from inhalation, as $HI < 1.00$. The average HI for the entire *Senior* population sub-group was 0.96. This suggests that more than two-thirds of this population sub-group would not experience any form of deleterious health effects due to living within and consuming rice, vegetables, fish, drinking water and dust from this urban location.

Fig 4 A reflects that all the *Children* were in health risk due to consumption of the dietary foods and from inhalation, as $HI > 1.00$ for all cases. This is attributed to more consumption of foods with respect to their body weight and more intimacy

with the contaminated dust/soil. The majority (65%) of the children had HI values between 2.00 and 3.00. The average HI for the *Highly Exposed Child* population sub-group was 2.36. The safety position of *Average Person* (i.e., Adult) population sub-group lied intermediate between child and senior. Clearly 60% of the adults were safe (Fig 4B) due to living within and consuming rice, vegetables, fish, drinking water and dust from this urban location. The average HI for the entire *Average Person* population sub-group was 1.00. Overall the human health safety order in the studied region was as follows: Senior > Average Person > Highly Exposed Child.

In UK, similar human health safety order was observed by Hough

et al (2004) for consuming vegetables alone. BGS (BGS/ DPHE, 2001) reported that the groundwater of Bangladesh was also contaminated with Co, Mn, Mo, Rb, Sr, U, etc. Moreover, there are some ‘arsenic hot-spots’ (a few kilometer across) in Chapai Nawabganj district (adjacent to Rajshahi district) and hydraulic modeling of BGS (BGS/DPHE, 2001) showed that the groundwater velocity of shallow aquifer ranged 5-20 mm day⁻¹. Thus this study highlights the importance of sight-specific human health risk assessment considering more pollutant parameters.

Acknowledgment

We gratefully acknowledge the authority of University of Rajshahi for providing University Grants Commission Fund (No. 506-5/ 52/ UGC/ Sci(6)) for the investigation. Our sincere gratitude goes to Professor Dr. Md. Hassan Ahmed, Former Director of Central Science Lab, R. U. , for his kind assistance in analyzing the environmental samples.

References

- Alam MGM, Snow ET, Tanaka A. Arsenic and heavy metal contamination of vegetables grown in Samta village, Bangladesh. *Science of the Total Environment* 2003; 1(3) : 83-96. [https://doi.org/10.1016/S0048-9697\(02\)00651-4](https://doi.org/10.1016/S0048-9697(02)00651-4)
- Adal A. Heavy metal toxicity. Chief editor: Tarabar A. 2017. <http://emedicine.medscape.com/article/814960-overview>. Accessed July 23 2017.
- APHA, AWWA, WEF. Standard methods for the examination of water and wastewater. American Public Health Association, American Water Works Association and Water Environment Federation. 23rd edition, ISBN: 9780875532875. United Book Press, Inc. , Baltimore, Maryland, 2017.
- ATSDR. Arsenic toxicity. Agency for Toxic Substances and Disease Registry, U. S. Department of Health and Human Services, Atlanta, GA, 2013. Federal Registrar: 52FR12866.
- ATSDR. Toxicological profile for lead. Agency for Toxic Substances and Disease Registry, U.S. Department of Health and Human Services, Atlanta, GA, 2007. Federal Registrar: 52FR12866.
- BBS. Yearbook of Agricultural Statistics-2016. 28th series. Bangladesh Bureau of Statistics. Govt. of the PR of Bangladesh, 2017. Available: http://bbs.portal.gov.bd/sites/default/files/files/bbs.portal.gov.bd/page/1b1eb817_9325_4354_a756_3d18412203e2/Yearbook-2016-Final-19-06-2017.pdf
- BBS. Household expenditure survey. Bangladesh Bureau of Statistics, Ministry of Planning, Government of the People’s Republic of Bangladesh, Dhaka, 2010. Available: <http://203.112.218.65:8008/WebTestApplication/userfiles/Image/LatestReports/HIES-10.pdf>
- BGS/ DPHE. Arsenic contamination of groundwater in Bangladesh. BGS Technical Report WC/ 00/ 19, British Geological Survey (UK) and Department of Public Health Engineering (Bangladesh) , (2001) (2): 77-207
- Bulka CM, Davis MA, Karagas MR, Ahsan H, Argos M. The Unintended Consequences of a Gluten-free Diet. *Epidemiology* 2007; 28(3) : 24–25. doi: 10.1097/EDE.0000000000000640
- CalEPA. Development of a Reference Dose (RfD) for Methamphetamine. Office of Environmental Health Hazard Assessment (OEHHA) , California Environmental Protection Agency, California, 2009. Available: <https://oehha.ca.gov/media/downloads/crn/methrfdfinal022609.pdf>

- Charlet L, Chakraborty S, Appelo CAJ, Ross RG, Nath B, Ansari AA, Lanson M, Chatterjee D, Mallik SB. Chemodynamics of an arsenic “hotspot” in a West Bengal aquifer: A field and reactive transport modeling study. *Applied Geochemistry* 2007; 22: 1273–1292. <https://doi.org/10.1016/j.apgeochem.2006.12.022>
- Das HK, Mitra AK, Sengupta PK, Hossain A, Islam F, Rabbani GH. Arsenic concentrations in rice, vegetables, a fish in Bangladesh: a preliminary study. *Environ. Int.* 2004; 30: 383-387. <https://doi.org/10.1016/j.envint.2003.09.005>
- Engconsult Limited. Arsenic calamity of Bangladesh. In: Statistics of arsenic calamity. Engconsult Limited, Toronto, Canada, 2017. Available: <http://www.engconsult.com/arsenic/arsstat.html>.
- FAO/ WHO. Evaluation of certain food additives and contaminants. WHO Technical Report Series, No. 959-960. Joint FAO/ WHO Expert Committee on Food Additives. Geneva, 2011.
- FAO/WHO. Principles and methods for the risk assessment of chemicals in food. In: Environmental Health Criteria 240. Food and Agriculture Organization and the World Health Organization, Geneva, 2009a.
- FAO/ WHO. CODEX General Standard for Contaminants and Toxins in Food and Feed. CODEX STAN 193-1995. Food and Agriculture Organization and the World Health Organization, Geneva, 2009b. Available: http://www.fao.org/fileadmin/user_upload/livestockgov/documents/1_CXS_193e.pdf.
- Fazal MA, Kawachi T, Ichion E. Extent and severity of groundwater arsenic contamination in Bangladesh. *Water Int.* 2001; 26(3) : 370-379. Available: <https://doi.org/10.1080/02508060108686929>.
- GRCIRAT. Developing new approaches to assessing risk to human health from chemicals. Government/Research Councils Initiative on Risk Assessment and Toxicology, Institute for Environment and Health, University of Leicester, UK, 1999. ISBN: 1899110224.
- Hesse PR. A textbook of soil chemical analysis. In: Total (elemental) analysis and some trace elements, Chapter 16, CBS Publishers & Distributor, Delhi, India, 1994; 371-435. ISBN: 9788123918334.
- Hough RL, Breward N, Young SD, Crout NMJ, Tye AM, Moir AM, Thornton I. Assessing potential risk of heavy metal exposure from consumption of home produced vegetables by urban populations. *Environmental Health Perspectives* 2004, 112(2): 215-221. DOI:10.1289/ehp.5589.
- IARC. IARC Monographs: Arsenic, metals, fibres and dusts. Vol. 100 C. International Agency for Research on Cancer, Lyon, France, 2012a. Available: <http://monographs.iarc.fr/ENG/Monographs/vol100C/mono100C.pdf>
- IARC. IARC Monographs on the evaluation of carcinogenic risks to humans: Cadmium and cadmium compounds. Vol. 100 C. International Agency for Research on Cancer, Lyon, France, 2012b. Available: <http://monographs.iarc.fr/ENG/Monographs/vol100C/mono100C-8.pdf>
- IARC. IARC Monographs on the evaluation of carcinogenic risks to humans: Inorganic and organic lead compounds. Vol. 87. International Agency for Research on Cancer, Lyon, France, 2006. Available: <http://monographs.iarc.fr/ENG/Monographs/vol87/mono87.pdf>
- Islam MS, Ahmed MK, Mamun MHA, Eaton DW. Arsenic in the food chain and assessment of population health risks in Bangladesh. *Environment Systems and Decisions* 2017; 37(3) : 344-352. <https://doi.org/10.1007/s10669-017-9635-8>.
- Islam MN, Das BK, Huque ME. Arsenic accumulation in common vegetables from irrigation. *J. Sci. Res.* 2012; 4(3): 675-688. <http://dx.doi.org/10.3329/jsr.v4i3.10494>
- Jackson ML. Soil chemical analysis. In: Plant Tissue Analysis-Mineral Constituents, Chapter 12, Prentice - Hall, Inc. , New Jersey, 1958; 326-338. DOI: 10.1002/jpln.19590850311

- Meharg AA, Rahman MM. Arsenic contamination in Bangladesh paddy field soils: Implications for rice contribution to arsenic consumption. *Environ. Sci. Technol.* 2003; 37(2) : 229-234. DOI: 10.1021/es0259842
- Saha GC, Ali MA. Dynamics of arsenic in agricultural soils irrigated with arsenic contaminated groundwater in Bangladesh. *Sci. Total Environ* 2007; 379: 180-189. <https://doi.org/10.1016/j.scitotenv.2006.08.050>
- Saha N, Zaman MR. Concentration of selected toxic metals in groundwater and some cereals grown in Shibganj area of Chapai Nawabganj. *Current Science* 2011; 101(3): 427-431.
- Sakamoto H, Susa Y, Ishiyama H, Tomiyasu T, Anazawa K. Determination of trace amounts of total arsenic in environmental samples by hydride generation flow injection-AAS using a mixed acid as a pretreatment agent. *Analytical Sciences* 2001; 17: 1067-1071. DOI: 10.2116/analsci.17.1067.
- Shimadzu (----) Atomic Absorption Spectrophotometry CookBook. In: Section 6 (Food staffs analysis) . Shimadzu Corporation, Kyoto, Japan.
- SOES/ DCH. Groundwater arsenic contamination in Bangladesh. School of Environmental Studies, Jadavpur University, India and Dhaka Community Hospital, Dhaka, Bangladesh, 2000. Available: <http://www.engconsult.com/arsenic/article/BANGLAREPORT.pdf>
- USEPA. Guidelines for human exposure assessment (draft) . Risk Assessment Forum, U. S. Environmental Protection Agency, Washington, DC, 2016. Available:https://www.epa.gov/sites/production/files/201602/documents/guidelines_for_human_exposure_assessment_peer_review_draftv2.pdf
- USEPA. Exposure Factors Handbook. U. S. Environmental Protection Agency, Washington, DC, 2011. Federal Registrar: EPA/600/R-09/052F.
- USEPA. Background discussion for soil-plant-human exposure pathway. In: Soil screening guidance. The Office of Emergency and Remedial Response (OERR), U. S. Environmental Protection Agency, Washington, DC, 2002. Federal Registrar EPA/540/R-96/018.
- USEPA. Exposure and Health Effects. U. S. Environmental Protection Agency, Washington, DC, 2001. Available: <https://pdfs.semanticscholar.org/4cec/76fe4f054a4c7984446362b38c8577dda9f3.pdf>
- USEPA. Guidelines for exposure assessment. Risk Assessment Forum, U. S. Environmental Protection Agency, Washington, DC, 1992. Federal Registrar 57(104):22888-22938.
- Williams PN, Islam MR, Adomako EE, Raab A, Hossain SA, Zhu YG, Feldmann J, Meharg AA. Increase in rice grain arsenic for regions of Bangladesh irrigating paddies with elevated arsenic in ground waters. *Environ Sci. Technol.* 2006; 39(15): 5531-40. DOI: 10.1021/es060222i.
- WHO. Guidelines for drinking-water quality. 4th edition. World Health Organization, Geneva. . 2017; 315-320. ISBN: 9789241549950.
- WHO. Arsenic. Fact sheet. In: Health effects. World Health Organization, Geneva, 2016a. Available: <http://www.who.int/mediacentre/factsheets/fs372/en>.
- WHO. Lead poisoning and health. Fact sheet. World Health Organization, Geneva, 2016b. Available: <http://www.who.int/mediacentre/factsheets/fs379/en>.
- WHO. Exposure to cadmium: A major public health concern. World Health Organization, Geneva, 2010. Available: <http://www.who.int/ipcs/features/cadmium.pdf?ua=1>
- Zereen F, Islam F, Habib MA, Bugum D, Zaman MS. Inorganic pollutants in the Padma river. *Environmental Geology* 2000; 39(9) : 1059-1062. Avilable: <https://doi.org/10.1007/s002549900098>.