

REFERENCES

- 1 Jati IRAP, Vadivel V, Biesalski HK. Antioxidant activity of anthocyanins in common legume grains, in bioactive food as dietary interventions for liver and gastrointestinal disease. 2013; 485-97.
- 2 Porres JM, Wu M, Cheng WH. Legumes, genome maintenance, and optimal health, in bioactive food as dietary interventions for the aging population. 2013; 321-34.
- 3 Singh S, Singh HD, Sikka KC. Distribution of nutrients in the anatomical parts of common Indian pulses. Nutrient in Indian pulses. 1968;45: 13-8.
- 4 Graham PH, Vance CP. Legumes: importance and constraints to greater use. Plant Physiol, 2003;131(3): 872-77.
- 5 Zhao Y, Du SK, Wang H, Cai M. In vitro antioxidant activity of extracts from common legumes. Food Chem. 2014; 152: 462-66.
- 6 Zhou K, Slavin M, Lutterodt H, Whent M, Eskin MNA, Yu L. Cereals and legumes. 2013; 3-48.
- 7 Choung MG, Choi BR, An YN, Chu YH, Cho YS. Anthocyanin profile of Korean cultivated kidney bean (*Phaseolus vulgaris* L.). J Agric Food Chem. 2003; 51: 7040-43.
- 8 Kris-Etherton PM, Hecker KD, Bonanome A, Coval SM, Binkoski AE, Hilpert KF, Griel AE, Etherton TD. Bioactive Compounds in Foods: Their Role in the Prevention of Cardiovascular Disease and Cancer. Am J Med. 2002; 113(9B): 71-88.
- 9 Anwar F, Latif S, Przybylski R, Sultana B, Ashraf M. Chemical composition and antioxidant activity of seeds of different cultivars of mungbean. J Food Sci, 2007; 72(7): 503-10.
- 10 ประพันธ์ บุญกลั่นขจร, ประสูติ สิทธิธรรม, สุวิทย์ เลาหศิริวงศ์, สนิท ลวดทอง, จรรย์ จันทลักษณ์. ถั่วลิสงและถั่วอื่นๆบางชนิด. รายงานการสัมมนา สมาคมวิทยาศาสตร์การเกษตรแห่งประเทศไทย. 2521.

- 11 อภิพรรณ พุกภักดี. วิทยาศาสตร์การผลิตพืชตระกูลถั่ว. มหาวิทยาลัยเกษตรศาสตร์. 2533.
- 12 Puwastien P, Raroengpuag M, Sungpuag M, Judprasong K. Thai food composition tables, 1st edition. 1999, Institute of nutrition, Mahidol University (INMU), Thailand.
- 13 Lerer-Metzger M, Rizkalla SW, Luo J, Champ M, Kabir M, Bruzzo F, Bornet F, Slama G. Effects of long-term low-glycaemic index starchy food on plasma glucose and lipid concentrations and adipose tissue cellularity in normal and diabetic rats. *Br J Nutr.* 1996; 75(5): 723-32.
- 14 อัจฉรา อุทโยภาส, รัตนา เสวตาลย์, ชวาลอุทอม ไชยนิติ, เอกสารวิชาการเรื่อง ถั่วเหลือง . กรมส่งเสริมการเกษตร. 2546.
- 15 Rochfort S, Panozzo J. Phytochemical for health, The Role of pulses. *J. Agric. Food Chem.* 2007; 55: 7981-94.
- 16 Fritz J, Shamma S, Elhilali M, Klein D. Rapid task-related plasticity of spectrotemporal receptive fields in primary auditory cortex. *Nat Neurosci.* 2003; 6(11): 1216-23.
- 17 Tripathi RP, Tewari N, Dwivedi N, Tiwari VK. Fighting tuberculosis: an old disease with new challenges. 2005; *Med Res Rev.* 2005; 25(1): 93-131.
- 18 Kumar S, Verma AK, Das M, Jain SK, Dwivedi PD. Clinical complications of kidney bean (*Phaseolus vulgaris* L.) consumption. *Nutrition.* 2013; 29(6): 821-7.
- 19 Tsuda T, Watanabe M, Ohshima K, Norinobu S, Choi SW, Kawakishi S, Osawa T. Antioxidative activity of the anthocyanin pigments cyanidin 3-O- β -D-glucoside and cyanidin. *J Agric Food Chem.* 1994; 42: 2407-10.
- 20 Gibbs BF, Alli I. Characterization of a purified α -amylase inhibitor from white kidney beans (*Phaseolus vulgaris*). *Food Research International*, 1998; 31(3): 217-25.
- 21 Lazzé MC, Pizzala R, Savio M, Stivala LA, Prosperi E, Bianchi L. Anthocyanins protect against DNA damage induced by tert-butylhydroperoxide in rat smooth muscle and hepatoma cells. *Mutat Res.* 2003; 535(1): 103-15.
- 22 ภูวนาท นนทรี. โครงการหนังสือเกษตรชุมชน. มหาวิทยาลัยเกษตรศาสตร์ บางเขน. 2531.

- 23 Chukwumah Y, Walker LT, Verghese M. Peanut skin color: a biomarker for total polyphenolic content and antioxidative capacities of peanut cultivars. *Int J Mol Sci*, 2009; 10(11): 4941-52.
- 24 Azevedo L¹, Gomes JC, Stringheta PC, Gontijo AM, Padovani CR, Ribeiro LR, Salvadori DM. Black bean (*Phaseolus vulgaris* L.) as a protective agent against DNA damage in mice. *Food Chem Toxicol*. 2003; 41(12): 1671-6.
- 25 Tovar J, Granfeldt Y, Bjorck IM. Effect of processing on blood glucose and insulin responses to starch in legumes. *J Agric Food Chem*. 1992; 40: 1846-51.
- 26 Zhang X, Shang P, Qin F, Zhou Q, Gao B, Huang H, Yang H, Shi H, Yu LL. Chemical composition and antioxidative and anti-inflammatory properties of ten commercial mung bean samples. *Food Sci Technol*. 2013; 54(1): 171-8.
- 27 Bed JM, Roberts DCK. The nutritional & biological significance of saponins. *Nutrition research*, 1995; 15(8): 1223-49.
- 28 Udani J, Hardy M, and Madsen DC. Blocking Carbohydrate Absorption and Weight Loss A Clinical Trial Using Phase 2™ Brand Proprietary Fractionated White Bean Extract. *Altern Med Rev* 2004; 9(1): 63-9.
- 29 Bansode RR, Randolph P, Ahmedna M, Hurley S, Hanner T, Baxter SA, Johnston TA, Su M, Holmes BM, Yu J, Williams LL. Bioavailability of polyphenols from peanut skin extract associated with plasma lipid lowering function. *Food Chem*. 2014; 148: 24-9.
- 30 Amarowicz R, Pegg RB. Legumes as a source of natural antioxidants. *Eur. J. Lipid Sci. Technol*. 2008; 110: 865–78.
- 31 Kabagambe EK, Baylin A, Ruiz-Narvarez E, Siles X, Campos H. Decreased consumption of dried mature beans is positively associated with urbanization and nonfatal acute myocardial infarction. *J Nutr*. 2005; 135(7): 770-5.
- 32 Gautam B, Vadivel V, Stuetz W, Biesalski HK. Bioactive compounds extracted from Indian wild legume seeds: antioxidant and type II diabetes-related enzyme inhibition properties. *Int J Food Sci Nutr*, 2012; 63(2): 242-5.

- 33 Marathe SA, Rajalakshmi V, Jamdar SN, Sharma A. Comparative study on antioxidant activity of different varieties of commonly consumed legumes in India. *Food Chem Toxicol*, 2011; 49(9): 2005-12.
34. Dykes L, Rooney LW. Phenolic Compounds in Cereal Grains and Their Health Benefits, in *Cereal Foods World*. 2007: 105-11.
- 35 Kris-Etherton PM, Hecker KD, Bonanome A, Coval SM, Binkoski AE, Hilpert KF, Griel AE, Etherton TD. Bioactive Compounds in Foods: Their Role in the Prevention of Cardiovascular Disease and Cancer. *Am J Med*. 2002; 113(9B): 71-88.
- 36 Jeon SY, Kwon SH, Seong YH, Bae K, Hur JM, Lee YY, Suh DY, Song KS. Beta-secretase(BACE1)-inhibiting stilbenoids from *Smilax Rhizoma*. *Phytomedicine*, 2007; 14(6): 403-8.
- 37 Nickavar B, Amin G. Bioassay-Guided Separation of an α -Amylase Inhibitor Anthocyanin from *Vaccinium arctostaphylos* Berries. *Z. Naturforsch C*. 2010; 65(9-10): 567-70.
- 38 Xu BJ, Chang SK. A comparative study on phenolic profiles and antioxidant activities of legumes as affected by extraction solvents. *J Food Sci*. 2007; 72(2): 159-66.
- 39 Madhujith T, Naczki M, Shahidi F. Antioxidant activity of common bean (*Phaseolus Vulgaris* L.). *J FOOD LIPIDS*. 2004; 11: 220–33.
- 40 Ikarashi N, Takeda R, Ito K, Ochiai W, Sugiyama K. The inhibition of lipase and glucosidase activities by acacia polyphenol. *Evid Based Complement Alternat Med*. 2011; 272-75.
- 41 Yao LH, Jiang YM, Shi J, Tomas-Barberan FA, Datta N, Singanusong R, Chen SS. Flavonoids in Food and Their Health Benefits. *Plant Foods Hum Nutr*. 2004; 59: 113-22.
- 42 Ren W, Qiao Z, Wang H, Zhu L, Zhang L. Flavonoids: promising anticancer agents. *Med Res Rev*, 2003; 23(4): 519-34.
- 43 Boschin G, Arnoldi A, Legumes are valuable sources of tocopherols. *Food Chem*. 2011; 127(3): 1199-203.
- 44 Fairfield KM, Fletcher RH. Vitamins for Chronic Disease Prevention in Adults. *Jama*. 2002; 287(23): 3116-126.

- 45 Freedman JE, Farhat JH, Loscalzo J, Keaney JF. α -Tocopherol Inhibits Aggregation of Human Platelets by a Protein Kinase C-Dependent Mechanism. *Circulation*. 1996; 94: 2434-440.
- 46 Grelaap ER, Giinterb KD. Fatty acid composition and tocopherol content of some legume seeds. *Anim Feed Sci Technol*. 1995; 52: 325-31.
- 47 Moriyama M, Oba K. Comparative study on the vitamin C contents of the food legume seeds. *J Nutr Sci Vitaminol (Tokyo)*. 2008; 54(1): 1-6.
- 48 Nilsson J, Stegmark R, Åkesson B. Total antioxidant capacity in different pea (*Pisum sativum*) varieties after blanching and freezing. *Food Chem*. 2004; 86(4): 501-7.
- 49 Campos P, Saguy A, Ernsberger P, Oliver E, Gaesser G. The epidemiology of overweight and obesity: public health crisis or moral panic? *Int J Epidemiol*. 2006; 35(1): 55-60.
- 50 Popkin BM, Adair LS, Ng SW. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev*. 2012; 70(1): 3-21.
- 51 Padwal RS, Majumdar SR. Drug treatments for obesity: orlistat, sibutramine, and rimonabant. *The Lancet*. 2007; 369(9555): 71-7.
- 52 Heck AM, Yanovski JA, Calis KA. Orlistat, a New Lipase Inhibitor for the Management of Obesity. *Pharmacotherapy*. 2000; 20(3): 270-9.
- 53 Gironés-Vilaplana A, Baenas N, Villaño D, Speisky H, García-Viguera C, Moreno DA. Evaluation of Latin-American fruits rich in phytochemicals with biological effects. *J Funct Foods*. 2014; 7: 599-608.
- 54 Adisakwattana S, Intrawangso J, Hemrid A, Chanathonh B, Mkyneen K. Extracts of Edible Plants Inhibit Pancreatic Lipase, Cholesterol Esterase and Cholesterol Micellization, and Bind Bile Acids. *Food Technol. Biotechnol*. 2012; 50(1): 11-16.
- 55 Joanitti G, Freitas S, Azevedo R. Legume-Derived Bioactive Compounds for the Prevention and Treatment of Breast Cancer, Breast Cancer in Current and Alternative Therapeu Modalities. Esra Gunduz Editor. 2011, InTech. 540 pages.

- 56 Sreerama YN, Takahashi Y, Yamaki K. Phenolic antioxidants in some Vigna species of legumes and their distinct inhibitory effects on alpha-glucosidase and pancreatic lipase activities. *J Food Sci.* 2012; 77(9): 927-33.
- 57 Siah SD, Konczak I, Agboola S, Wood JA, Blanchard CL. *In vitro* investigations of the potential health benefits of Australian-grown faba beans (*Vicia faba* L.): chemopreventative capacity and inhibitory effects on the angiotensin-converting enzyme, alpha-glucosidase and lipase. *Br J Nutr.* 2012; 108: 123-34.
- 58 Sergent T, Vanderstraeten J, Winand J, Beguin P, Schneider YJ. Phenolic compounds and plant extracts as potential natural anti-obesity substances. *Food Chem.*, 2012; 135(1): 68-73.
- 59 Worsztynowicz P, Napierała M, Białasa W, Grajeka W, Olkowicz M. Pancreatic α -amylase and lipase inhibitory activity of polyphenolic compounds present in the extract of black chokeberry (*Aronia melanocarpa* L.). *Process Biochemistry.* 2014; 49(9): 1457-63.
- 60 You Q, Chen F, Wang X, Luo PG, Jiang Y. Inhibitory effects of muscadine anthocyanins on alpha-glucosidase and pancreatic lipase activities. *J Agric Food Chem.* 2011; 59(17): 9506-11.
- 61 Wild S, Roglic G, Green A, Sicree R, King H. Global Prevalence of Diabetes Estimates for the year 2000 and projections for 2030. *Diabetes Care.* 2004; 27(5): 1047-53.
- 62 King H, Aubert RE, Herman WH, Global Burden of Diabetes, 1995–2025 Prevalence, numerical estimates, and projections. *Diabetes Care.* 1998; 21(9): 1414-31.
- 63 Weyer C, Bogardus C, Mott DM, Pratley RE. The natural history of insulin secretory dysfunction and insulin resistance in the pathogenesis of type 2 diabetes mellitus. *J Clin Invest.* 1999; 104(6): 787-94.
- 64 Inzucchi SE. Oral Antihyperglycemic Therapy for Type 2 Diabetes. *Jama.* 2002; 287(3): 360-72.
- 65 Singh K, Kayastha AM. Alpha-Amylase from wheat (*Triticum aestivum*) seeds: Its purification, biochemical attributes and active site studies. *Food Chem.*

- 2014; 162: 1-9.
- 66 Sakulnarmrat K. and Konczak I. Composition of native Australian herbs polyphenolic-rich fractions and in vitro inhibitory activities against key enzymes relevant to metabolic syndrome. *Food Chem.* 2012; 134(2): 1011-9.
- 67 Puls W, Keup U, Krause HP, Thomas G, Hoffmeister F. Glucosidase Inhibition. *Naturwissenschaften.* 1977; 54: 535-6.
- 68 Kumar S, Kumar V, Chandrashekhar MS. *In-vitro* Anti-oxidant and alpha-amylase inhibitory activity of isolated fractions from methanolic extract of *Asystasia Dalzelliana* leaves. *Int J PharmTech Res.* 2011; 3(2): 889-94.
- 69 Chang HB, Kim SH, Kwon YI, Choung DH, Choi WK, Kang TW, Lee S, Kim JG, Chun HS, Ahn SK, Hong CI, Han KH. Novel α -Glucosidase inhibitors, CKD-711 and CKD-711a produced by *Streptomyces sp.* CK-4416. *J Antibiot (Tokyo).* 2002; 55(5): 467-71.
- 70 Chiasson JL, Josse RG, Gomis R, Hanefeld M, Karasik A, Laakso M. Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomised trial. *Lancet.* 2002; 359(9323): 2072-77.
- 71 Chandramohan R, Pari L, Rathinam A, Sheikh BA. Tyrosol, a phenolic compound, ameliorates hyperglycemia by regulating key enzymes of carbohydrate metabolism in streptozotocin induced diabetic rats. *Chem Biol Interact.* 2015; 229: 44-54.
- 72 McDougall JG, Shpiro F, Dobson P, Smith P, Blake A, Stewart D. Different polyphenolic components of soft fruits inhibit alpha-amylase and alpha-glucosidase. *J Agric Food Chem.* 2005; 53: 2760-66.
- 73 Vadivel V, Cheong JN, Biesalski HK. Antioxidant and type II diabetes related enzyme inhibition properties of methanolic extract of an underutilized food legume, *Canavalia ensiformis* (L.) DC: Effect of traditional processing methods. *Food Sci Technol.* 2012; 47(2): 255-60.
- 74 Vadivel V, Biesalski HK, Contribution of phenolic compounds to the antioxidant potential and type II diabetes related enzyme inhibition properties of *Pongamia pinnata* L. Pierre seeds. *Process Biochemistry.* 2011; 46(10): 1973-1980.

- 75 You Q, Chen F, Wang X, Jiang Y, Lin S. Anti-diabetic activities of phenolic compounds in muscadine against alpha-glucosidase and pancreatic lipase. *Food Sci Technol*. 2012; 46(1): 164-8.
- 76 Boath AS, Stewart D, McDougall GJ. Berry components inhibit alpha-glucosidase in vitro: synergies between acarbose and polyphenols from black currant and rowanberry. *Food Chem*. 2012; 135(3): 929-36.
- 77 Kumar S, Kumar V, Rana M, Kumar D. Enzymes inhibitors from plants: an Alternate Approach To Treat Diabetes. *Pharmacognosy Communications*. 2012; 2(2): 18-33.
- 78 Guo H, Xia M. Anthocyanins and Diabetes Regulation, in *Polyphenols in Human Health and Disease*. 2014; 83-93.
- 79 Liu YJ, Zhan J, Liu XL, Wang Y, Ji J, He QQ. Dietary flavonoids intake and risk of type 2 diabetes: a meta-analysis of prospective cohort studies. *Clin Nutr*. 2014; 33(1): 59-63.
- 80 Sancho RAS, Pastore GM. Evaluation of the effects of anthocyanins in type 2 diabetes. *Food Res Int*. 2012; 46(1): 378-86.
- 81 Ademiluyi AO, Oboh G. Soybean phenolic-rich extracts inhibit key-enzymes linked to type 2 diabetes (alpha-amylase and alpha-glucosidase) and hypertension (angiotensin I converting enzyme) in vitro. *Exp Toxicol Pathol*, 2013; 65(3): 305-9.
- 82 Vadivel V, Stuetz W, Scherbaum V, Biesalski HK. Total free phenolic content and health relevant functionality of Indian wild legume grains: Effect of indigenous processing methods. *Journal of Food Composition and Analysis*, 2011; 24(7): 935-43.
- 83 Girish TK, Pratape VM, Rao UJSP. Nutrient distribution, phenolic acid composition, antioxidant and alpha-glucosidase inhibitory potentials of black gram (*Vigna mungo* L.) and its milled by-products. *Food Res Int*. 2012; 46(1): 370-77.
- 84 Yao Y, Cheng X, Wang L, Wang S, Ren G. A determination of potential alpha-glucosidase inhibitors from Azuki Beans (*Vigna angularis*). *Int J Mol Sci*. 2011; 12(10): 6445-51.

- 85 Chockalingam, A., Campbell NR, Fodor JG, Worldwide epidemic of hypertension. *Canadian Journal of Cardiology*. 2006; 22(7): 553-55.
- 86 Kearney PM, Whelton M, Reynolds K, Whelton PK, He J. Worldwide prevalence of hypertension: a systematic review. *Journal of Hypertension*. 2004; 21(1): 11-19.
- 87 Weber MA, Schiffrin EL, White WB, White, Mann S, Lindholm LH, Kenerson JG, *et al.* Clinical practice guidelines for the management of hypertension in the community a Statement by the American Society of Hypertension and the International Society of Hypertension. *J CLIN HYPERTENS*. 2013: 1-13.
- 88 Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *The Lancet*. 2005; 365(9455): 217-23.
- 89 ศุภนิมิต ที่มขุณหเถยร. Inhibitors of Renin-angiotensin system: 17-33.
- 90 Chen ZY, Peng C, Jiao R, Wong YM, Yang N, Huang Y. Anti-hypertensive nutraceuticals and functional foods. *J Agric Food Chem*. 2009; 57(11): 4485-99.
- 91 Israili ZH, Hall WD. Cough and Angioneurotic Edema Associated with Angiotensin-Converting Enzyme Inhibitor Therapy: A Review of the Literature and Pathophysiology. *Ann Intern Med*. 1992; 117(3): 234-42.
- 92 Modena, M.G., Hypertension, Menopause and Natural Antioxidants in Foods and the Diet, in *Natural antioxidant and hypertension*. 2014; 225-31.
- 93 Eriz G, Sanhueza V, Roeckel M, Fernandez K. Inhibition of the angiotensin-converting enzyme by grape seed and skin proanthocyanidins extracted from *Vitis vinífera L. cv. País*. *Food Science and Technology*. 2011; 44(4): 860-65.
- 94 Simaratanamongkol A, Umehara K, Noguchi H, Panichayupakaranant P. Identification of a new angiotensin-converting enzyme (ACE) inhibitor from Thai edible plants. *Food Chem*. 2014; 165: 92-7.
- 95 Lacaille-Dubois, Franck U, Wagner H. Search for potential angiotensin converting enzyme (ACE)-inhibitors from plants. *Phytomedicine*. 2001; 8(1): 47-52.

- 96 Balasuriya BWN, Rupasinghe HPV. Plant flavonoids as angiotensin converting enzyme inhibitors in regulation of hypertension. *Funct. food health dis.* 2011; 5: 172-88.
- 97 Kannappan R, Gupta SC, Kim JH, Reuter S, Aggarwal BB. Neuroprotection by spice-derived nutraceuticals: you are what you eat! *Mol Neurobiol.* 2011; 44(2): 142-59.
- 98 Bendlin BB, Carlsson CM, Gleason CE, Johnson SC, Sodhi A, Gallagher CL, *et al.* Midlife predictors of Alzheimer's disease. *Maturitas.* 2010; 65(2): 131-7.
- 99 Rao AA, Sridhar GR, Das UN. Elevated butyrylcholinesterase and acetylcholinesterase may predict the development of type 2 diabetes mellitus and Alzheimer's disease. *Med Hypotheses.* 2007; 69(6): 1272-6.
- 100 Lee JW, Lee YK, Ban JO, Ha TY, Yun YP, Han SB, *et al.* Green tea (-)-epigallocatechin-3-gallate inhibits beta-amyloid-induced cognitive dysfunction through modification of secretase activity via inhibition of ERK and NF-kappaB pathways in mice. *J Nutr.* 2009; 139(10): 1987-93.
- 101 Orhan I, Kartal M, Tosun F, Sener B. Screening of Various Phenolic Acids and Flavonoid Derivatives for their Anticholinesterase Potential. *Z Naturforsch.* 2007; 62c: 829-32.
- 102 Jung HA, Min BS, Yokozawa T, Lee JH, Kim YS, Choi JS, Anti-Alzheimer and antioxidant activity of *Coptidis Rhizoma* alkaloids. *Biol Pharm Bull.* 2009; 32(8): 1433-8.
- 103 Brewer GJ, Torricelli JR, Lindsey AL, Kunz EZ, Neuman A, Fisher DR, *et al.* Age-related toxicity of amyloid-beta associated with increased pERK and pCREB in primary hippocampal neurons: reversal by blueberry extract. *J Nutr Biochem.* 2010; 21(10): 991-8.
- 104 Yordi EG, Perez EM, Matos MJ, Villares EU. Antioxidant and Pro-Oxidant Effects of Polyphenolic Compounds and Structure-Activity Relationship Evidence, in *Nutrition, Well-Being and Health*, D.J. Bouayed, Editor. 2012; InTech. p. 224.
- 105 Hogervorst E, Sadjimim T, Yesufu A, Kreager P, Rahardjo TB. High tofu intake

- is associated with worse memory in elderly Indonesian men and women. *Dement Geriatr Cogn Disord*. 2008; 26: 50-7.
- 106 Henry-Vitrac C, Berbille H, Merillon JM, Vitrac X. Soy isoflavones as potential inhibitors of Alzheimer β -amyloid fibril aggregation in vitro. *Food Rev Int*. 2010; 43(8): 2176-8.
- 107 Choi DY, Lee YJ, Hong JT, Lee HJ. Antioxidant properties of natural polyphenols and their therapeutic potentials for Alzheimer's disease. *Brain Res Bull*. 2012; 87(2-3): 144-53.
- 108 Nepote V, Grosso NR, and Guzman CA. Optimization of extraction of phenolic antioxidants from peanut skins. *J Sci Food Agric*. 2005; 85(1): 33-38.
- 109 Merken HM, Beecher GR. Liquid chromatographic method for the separation and quantification of prominent flavonoid aglycones. *Journal of Chromatography*. 2000; 897: 177-84.
- 110 Fukumoto LR, Mazza G. Assessing antioxidant and prooxidant activities of phenolic compounds. *J Agric Food Chem*. 2000; 48: 3597-604.
111. Benzie IF, Strain JJ. The Ferric Reducing Ability of Plasma (FRAP) as a Measure of "Antioxidant Power": The FRAP Assay. *Anal Biochem*. 1996; 239: 70-6.
- 112 Ou B, Hampsch-Woodill M, Prior RL. Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe. *J Agric Food Chem*. 2001; 49: 4610-26.
- 113 Ainsworth EA, Gillespie KM. Estimation of total phenolic content and other oxidation substrates in plant tissues using Folin-Ciocalteu reagent. *Nat Protoc*. 2007; 2: 875-7.
- 114 Choi SJ, Hwang JM, Kim SI. A Colorimetric Microplate Assay Method for High Throughput Analysis of Lipase Activity. *J Biochem Mol Biol*. 2003; 36(4): 417-20.
- 115 Funke I. and Melzig MF. Traditionally used plants in diabetes therapy – phytotherapeutics as inhibitors of α -amylase activity. *Braz J. Pharmacogn*. 2006; 16(1): 1-5.
- 116 Schwager SL, Carmona KA, Sturrock ED. A high-throughput fluorimetric assay

- for angiotensin I-converting enzyme. *Nature Protocols*. 2006; 1(4): 1961-4.
- 117 Alu'datt MH, Rababah T, Ereifej K, Alli I. Distribution, antioxidant and characterisation of phenolic compounds in soybeans, flaxseed and olives. *Food Chem*. 2013; 139(1-4): 93-9.
- 118 Huang MT, Wood AW, Newmark HL, Sayer JM, Yagi H, Jerina DM, *et al*. Inhibition of the mutagenicity of bay-region diol-epoxides of polycyclic aromatic hydrocarbons by phenolic plant flavonoids. *Carcinogenesis*, 198; 4(12): 1631-37.
- 119 Kanu P, Zhu k, Kanu JB, Zhou H, Qian H, Zhu K. RETRACTED: Biologically active components and nutraceuticals in sesame and related products: a review and prospect. *Trends in Food Science & Technology*. 2007; 18(12): 599-608.
- 120 Chatonnet A, Lockridge O, Comparison of butyrylcholinesterase and acetylcholinesterase. *Biochem J*. 1989; 260: 625-34.
- 121 Hart JJ, Tako E, Kochian LV, Glahn RP, Identification of black bean (*Phaseolus vulgaris* L.) polyphenols that inhibit and promote iron uptake by Caco-2 Cells. *J Agric Food Chem*. 2015; 63(25): 5950-6.
- 122 Elizabeth RGN, Annete H, Francisco GLR, Javier IPF, Graciela ZG, Alberto GIJ. Antioxidant and antimutagenic activity of phenolic compounds in three different colour groups of common bean cultivars (*Phaseolus vulgaris*). *Food Chem*. 2007; 103(2): 521-7.
- 123 Lin LZ, Harnly JM, Pastor-Corrales MS, Luthria DL. The polyphenolic profiles of common bean (*Phaseolus vulgaris* L.). *Food Chem*. 2008; 107(1): 399-410.
- 124 Yao Y, Chen F, Wang M, Wang J, Ren G. Antidiabetic Activity of Mung Bean Extracts in Diabetic KK-Ay Mice. *J Agric Food Chem*. 2008; 56: 8869-73.
- 125 Craft BD, Kosińska A, Amarowicz R, Pegg RB. Antioxidant properties of extracts obtained from raw, dry-roasted, and oil-roasted US peanuts of commercial importance. *Plant Foods Hum Nutr*. 2010; 65(3): 311-8.

- 126 Góes-Favoni SP, Carrão-Panizzi MC, Beleia A. Changes of isoflavone in soybean cotyledons soaked in different volumes of water. *Food Chem.* 2010; 119(4): 1605-12.
- 127 Kunyanga CN, Imungia JK, Okotha MW, Biesalski HK, Vadivel V. Total phenolic content, antioxidant and antidiabetic properties of methanolic extract of raw and traditionally processed Kenyan indigenous food ingredients. *Food Sci Technol.* 2012; 45(2): 269-76.
- 128 Bawadi HA, Bansode RR, Trappey A, Truax RE, Losso JN. Inhibition of Caco-2 colon, MCF-7 and Hs578T breast, and DU 145 prostatic cancer cell proliferation by water-soluble black bean condensed tannins. *Cancer Lett.* 2005; 218(2): 153-62.
- 129 Xu B, Chang SKC. Effect of soaking, boiling, and steaming on total phenolic content and antioxidant activities of cool season food legumes. *Food Chem.* 2008; 110(1): 1-13.
- 130 Adeniyani OO, Ibukun EO, Ogunbolude Y, Eseigbe MI. Effect of Boiling on the Nutritional Composition and Antioxidant Properties of Beniseed (*Sesamum indicum* L.). *Food Science and Quality Management.* 2013; 11: 39-49.
- 131 Kalyarat K, Tongyonk L, Kangsadalampai K. Antimutagenic and co-mutagenic activities of some legume seeds and their seed coats. *J. Med. Plants Res.* 2012; 6(22): 3845-51.
- 132 Ferracane R, Pellegrini N, Visconti A, Graziani G, Chiavaro E, Miglio C, *et al.* Effects of different cooking methods on antioxidant profile, antioxidant capacity, and physical characteristics of artichoke. *J Agric Food Chem.* 2008; 56: 8601–8.
- 133 Chumyam A, Whangchaia K, Jungklanga J, Faiyuec B, Saengnil K. Effects of heat treatments on antioxidant capacity and total phenolic content of four cultivars of purple skin eggplants. *ScienceAsia.* 2013; 39(3): 246-51.
- 134 Turkmen N, Sari F, Velioglu Y. The effect of cooking methods on total phenolics and antioxidant activity of selected green vegetables. *Food Chem.* 2005; 93(4): 713-8.
- 135 Yasmin A, Zeb A, Khalil AW, Paracha GMUD, Khattak B. Effect of Processing

- on Anti-nutritional Factors of Red Kidney Bean (*Phaseolus vulgaris*) Grains. Food Bioprocess Technol. 2008; 1(4): 415-9.
- 136 Duenas M, Hernandez T, Estrella I. Assessment of in vitro antioxidant capacity of the seed coat and the cotyledon of legumes in relation to their phenolic contents. Food Chem. 2006; 98(1): 95-103.
- 137 Reynoso-Camacho R, Ramos-Gomez M, Loarca-Pina G, Bioactive components in common beans (*Phaseolus vulgaris* L.), in Advances in Agricultural and Food Biotechnology, R.G.G.-G.a.I. Torres-Pacheco, Editor. 2006: Research and Graduate Studies in Food Science, School of Chemistry Universidad Autónoma de Queretaro, Queretaro, Qro., 76010 México. p. 217- 236.
- 138 Ranilla LGL, Genovese MIS, Lajolo FM. Polyphenols and Antioxidant Capacity of Seed Coat and Cotyledon from Brazilian and Peruvian Bean Cultivars (*Phaseolus vulgaris* L.). J Agric Food Chem. 2007; 55: 90-8.
- 139 Beninger CW, Hosfield GL. Flavonol Glycosides from Montcalm Dark Red Kidney Bean: Implications for the Genetics of Seed Coat Color in *Phaseolus vulgaris* L. J Agric Food Chem. 1999; 47: 4079-82.
- 140 Choung MG, Choi BR, An YN, Chu YH, Cho YS. Anthocyanin profile of korean cultivated kidney bean (*Phaseolus vulgaris* L.). J Agric Food Chem. 2003; 51: 7040-3.
- 141 Troszyńska A, Estrella I, LoHpez-Amores ML, Hernandez T. Antioxidant activity of pea (*Pisum sativum* L.) seed coat acetone extract. Food Sci Technol. 2002; 35(2): 158-64.
- 142 Romani A, Vignolini P, Galardi C, Mulinacci N, Benedettelli S, Heimler D. Germplasm characterization of Zolfino Landraces (*Phaseolus vulgaris* L.) by Flavonoid Content. J Agric Food Chem. 2004; 52: 3838-42.
- 143 Luthria DL, Pastor-Corrales MA. Phenolic acids content of fifteen dry edible bean (*Phaseolus vulgaris* L.) varieties. Food Comp Anal. 2006; 19(2-3): 205-11.
- 144 Cai S, Wang O, Wang M, He J, Wang Y, Zhang D, *et al.* In vitro inhibitory effect on pancreatic lipase activity of subfractions from ethanol extracts of fermented Oats (*Avena sativa* L.) and synergistic effect of three phenolic

- acids. *J Agric Food Chem.* 2012; 60(29): 7245-51.
- 145 Karamać M, Amarowicz R. Inhibition of pancreatic lipase by phenolic acids – examination in vitro. *Z. Naturforsch.* 1996; 51C: 903-5.
- 146 Liu S, Li D, Huang B, Chen Y, Lu X, Wang Y. Inhibition of pancreatic lipase, alpha-glucosidase, alpha-amylase, and hypolipidemic effects of the total flavonoids from *Nelumbo nucifera* leaves. *J Ethnopharmacol.* 2013; 149(1): 263-9.
- 147 McDougall GJ, Kulkarni NN, Stewart D, Berry polyphenols inhibit pancreatic lipase activity in vitro. *Food Chem.* 2009; 115(1): 193-9.
- 148 Randhir R, Kwon YI, Shetty K. Effect of thermal processing on phenolics, antioxidant activity and health-relevant functionality of select grain sprouts and seedlings. *Innovative Food Science & Emerging Technologies.* 2008; 9(3): 355-64.
- 149 Duenas M, Estrella I, Hernandez T. Occurrence of phenolic compound in the seed coat and the cotyledon of pea (*Pisum sativum* L.). *Europ Food Resea Techn.* 2004; 219: 116-23.
- 150 Sternby B, Hartmann D, Borgström B, Nilsson A. Degree of in vivo inhibition of human gastric and pancreatic lipases by Orlistat (Tetrahydrolipstatin, THL) in the stomach and small intestine. *Clin Nutr.* 2002; 21(5): 395-402.
- 151 Karamać M, Amarowicz R. Inhibition of pancreatic lipase by phenolic acids— examination in vitro. *Z Naturforsch C.* 1996; 51: 903-5.
- 152 Li YQ, Yang P, Gao F, Zhang ZW, Wu B. Probing the interaction between 3 flavonoids and pancreatic lipase by methods of fluorescence spectroscopy and enzymatic kinetics. *Europ Food Resea Techn.* 2011; 233(2): 63-9.
- 153 Narita Y, Inouye K. Inhibition of Porcine Pancreas α -Amylase by Chlorogenic Acids from Green Coffee Beans and Cinnamic Acid Derivatives. 2015: p. 757-63.
- 154 Kaur R, Gupta AK, Taggar GK. Characterisation and inhibition studies of *Helicoverpa armigera* (Hubner) gut alpha-amylase. *Pest Manag Sci.* 2014; 1-10.

- 155 Wongsap P, Chaiwarit J, Zamaludien A, In vitro screening of phenolic compounds, potential inhibition against α -amylase and α -glucosidase of culinary herbs in Thailand. *Food Chem.* 2012; 131(3): 964-71.
- 156 Wang H, Du YJ, Song HC, α -Glucosidase and α -amylase inhibitory activities of guava leaves. *Food Chem.* 2010; 123(1): 6-13.
- 157 Peng X, Zheng Z, Cheng KW, Shan F, Ren GX, Chen F, *et al.* Inhibitory effect of mung bean extract and its constituents vitexin and isovitexin on the formation of advanced glycation endproducts. *Food Chem.* 2008; 106(2): 475-81.
- 158 Farsi E, Shafaei A, Hor SY, Ahamed MBK, Yam MF, Attitalla IH. Correlation between enzymes inhibitory effects and antioxidant activities of standardized fractions of methanolic extract obtained from *Ficus deltoidea* leaves. *Afr. J. Biotechnol.* 2011; 10(67): 15184-94.
- 159 Yang JP, He H, Lu YH. Four flavonoid compounds from *Phyllostachys edulis* leaf extract retard the digestion of starch and its working mechanisms. *J Agric Food Chem.* 2014; 62(31): 7760-70.
- 160 Choo CY, Sulong NY, Man F, Wong TW, Vitexin and isovitexin from the leaves of *ficus deltoidea* with in-vivo alpha-glucosidase inhibition. *J Ethnopharmacol.* 2012; 142(3): 776-81.
- 161 Nickavara B, Amin G. Enzyme Assay Guided Isolation of an α -Amylase Inhibitor Flavonoid from *Vaccinium arctostaphylos* Leaves. *Iran J Pharm Res.* 2011;10(4): 849-53.
- 162 Li Z, Alli I, Kermasha S. In-vitro α -amylase inhibitor activity-phytate relationships in proteins from *Phaseolus* beans. *Food Res Int.* 1993; 26(3): 195-201.
- 163 Sama K, Sivaraaj R, Rajiv P. In vitro antidiabetic activity of anthocyanin extract of *Asystasia gangetica* (Chinese violet) flower. *Asian Journal of Plant Science and Research.* 2013; 3(2): 88-92.
- 164 Xu B, Chang SK. Phytochemical profiles and health-promoting effects of cool-season food legumes as influenced by thermal processing. *J Agric Food Chem.* 2009; 57(22): 10718-31.

- 165 Al Kazaz M, Desseaux V, Marchis-Mouren G, Prodanov E, Santimone M. The mechanism of porcine pancreatic alpha-amylase Inhibition of maltopentaose hydrolysis by acarbose, maltose and maltotriose. *Eur J Biochem.* 1998; 252: 100-7.
- 166 Talcott ST, Moore JP, Lound-Singleton AJ, Percival SP. Ripening associated phytochemical changes in mangoes (*Mangifera indica*) following thermal quarantine and low-temperature storage. *J Food Sci.* 2005; 70: 337–41.
- 167 Bland MJ, RA L. Isolation and Characterization of a Peanut Maturity-Associated Protein. *J Agric Food Chem.* 2000; 48: 3275-79.
- 168 Seo A, Morr CV. Activated carbon and ion exchange treatments for removing phenolics and phytate from peanut protein products. *J Food Sci.* 1985; 50(1): 262-63.
- 169 Mattila P, Hellström J. Phenolic acids in potatoes, vegetables, and some of their products. *Food Res Int.* 2007; 20(3-4): 152-60.
- 170 Phan-Thien KY, Wright GC, Lee NA. Peanut antioxidants: Part 2. Quantitation of free and matrix-bound phytochemicals in five selected genotypes with diverse antioxidant capacity by high performance liquid chromatography (HPLC). *Food Sci Technol.* 2014; 57(1): 312-19.
- 171 Laranjinha J, Vieira O, Madeira V, Almeida L. Two Related Phenolic Antioxidants with Opposite Effects on Vitamin E Content in Low Density Lipoproteins Oxidized by Ferrylmyoglobin: Consumption vs Regeneration. *Arch. Biochem. Biophys.* 1995; 323(2): 373–81.
- 172 Chen CYO, Blumberg JB. Phytochemical composition of nuts. *Asia Pac J Clin Nutr.* 2008; 17(S1): 329-32
- 173 Kerem Z, Bilkis I, Flaishman MA, Sivan L. Antioxidant activity and inhibition of alpha-glucosidase by trans-resveratrol, piceid, and a novel trans-stilbene from the roots of *Israeli Rumex bucephalophorus* L. *J Agric Food Chem.* 2006; 54(4): 1243-7.
- 174 Matsui T, Ueda T, Oki T, Sugita K, Terahara N, Matsumoto K. Alpha-glucosidase inhibitory action of natural acylated anthocyanins. 1. survey of natural pigments with potent inhibitory activity. *J Agric Food Chem.* 2001; 49: 1948-51.

- 175 Rojo LE, Ribnicky D, Logendra S, Poulev A, Rojas-Silva P, *et al.* In vitro and in vivo anti-diabetic effects of anthocyanins from Maqui Berry (*Aristotelia chilensis*). *Food Chem.* 2012; 131(2): 387-96.
- 176 Aparicio-Fernandez X, Yousef GG, Loarca-Pina G, de Mejia E, Lila MA. Characterization of polyphenolics in the seed coat of Black Jamapa bean (*Phaseolus vulgaris* L.). *J Agric Food Chem.* 2005; 11: 4615-22.
- 177 Ho HM, Chen RY, Leung LK, Chan FL, Huang Y, Chen ZY. Difference in flavonoid and isoflavone profile between soybean and soy leaf. *Biomed Pharmacother.* 2002; 56: 289-95.
- 178 Wang Y, Ma L, Li Z, Du Z, Liu Z, Qin J, *et al.* Synergetic inhibition of metal ions and genistein on alpha-glucosidase. *FEBS Lett.* 2004; 576(1-2): 46-50.
- 179 Lee DS, Lee SH. Genistein, a soy isoflavone, is a potent alpha-glucosidase inhibitor. *FEBS Letters.* 2001; 501: 84-6.
- 180 Duncan AC, Jager AK, van Staden J. Screening of Zulu medicinal plants for angiotensin converting enzyme (ACE) inhibitors. *J Ethnopharmacol.* 1999; 68: 63-70.
- 181 Al Shukor N, Van Camp J, Gonzales GB, Staljanssens D, Struijs K, Zotti MJ, *et al.* Angiotensin-converting enzyme inhibitory effects by plant phenolic compounds: a study of structure activity relationships. *J Agric Food Chem.* 2013; 61(48): 11832-9.
- 182 Ojeda D, Jiménez-Ferrer E, Zamilpa A, Herrera-Arellano A, Tortoriello J, Alvarez L. Inhibition of angiotensin convertin enzyme (ACE) activity by the anthocyanins delphinidin- and cyanidin-3-O-sambubiosides from *Hibiscus sabdariffa*. *J Ethnopharmacol.* 2010; 127(1): 7-10.
- 183 Parichatikanond W, Pinthong D, Mangmoo S. Blockade of the Renin-Angiotensin System with Delphinidin, Cyanin, and Quercetin. *Planta Med.* 2012; 78: 1626-32.
- 184 Wei L, Clauser E, Alhenc-Gelas F, Corvol P. The Two homologous domains of human angiotensin I-converting enzyme interact differently with competitive inhibitors. *J Biol Chem.* 1992; 267(19): 13398-405.

- 185 Li RS, Wang XB, Hu XJ, Kong LY. Design, synthesis and evaluation of flavonoid derivatives as potential multifunctional acetylcholinesterase inhibitors against Alzheimer's disease. *Bioorg Med Chem Lett*. 2013; 23(9): 2636-41.
- 186 Chen J, Quan MH, Cheng YQ, Sun J, Li LT. Acetylcholinesterase inhibitory activity of Chinese sufu (fermented tofu) ethanol-extract. *Food Chem*. 2012; 134(3): 1263-6.
- 187 Sharififar F, Moshafi MH, Shafazand E, Koohpayeh A. Acetyl cholinesterase inhibitory, antioxidant and cytotoxic activity of three dietary medicinal plants. *Food Chem*. 2012; 130(1): 20-3.
188. Lee JH, Choung MG. Determination of optimal acid hydrolysis time of soybean isoflavones using drying oven and microwave assisted methods. *Food Chem*. 2011; 129(2): 577-82.
- 189 Zubik L, Meydani M. Bioavailability of soybean isoflavones from aglycone and glucoside forms in American women. *Am J Clin Nutr*. 2003; 77:1459–65.
- 190 Lee, CH, Yang L, Xu JZ, Yeung SYV, Huang Y, *et al*. Relative antioxidant activity of soybean isoflavones and their glycosides. *Food Chem*. 2005; 90(4): 735-41.
- 191 Santos CSP, Cunha S, Casal S. Bioactive components in potatoes as influenced by thermal processing, in processing and impact on active components in food. 2015; p 111-119.
- 192 Sertovic E, Mujic I, Jokic S, Alibabic V, Saric Z. Effect of soybean cultivars on the content of isoflavones in soymilk. *Rom Biotech Lett*. 2012; 17(2): 7151-9.
- 193 Khan MT, Orhan I, Senol FS, Kartal M, Sener B, Dvorská M, *et al*. Cholinesterase inhibitory activities of some flavonoid derivatives and chosen xanthone and their molecular docking studies. *Chem Biol Interact*. 2009; 181(3): 383-9.
- 194 Khadri A, Neffatib M, Smiti S, Fale P, Lino ARL, Serralheiro MLM, *et al*. Antioxidant, antiacetylcholinesterase and antimicrobial activities of *Cymbopogon schoenanthus* L. Spreng (lemon grass) from Tunisia. *Food Sci Technol*. 2010; 43(2): 331-6.

- 195 Choi JS, Islam MN, Ali MY, Kim EJ, Kim YM, Jung HA. Effects of C-glycosylation on anti-diabetic, anti-Alzheimer's disease and anti-inflammatory potential of apigenin. *Food Chem Toxicol.* 2014; 64: 27-33.
- 196 Citron M. Beta-secretase as a target for the treatment of Alzheimer's disease. *J Neurosci Res.* 2002; 70(3): 373-9.
- 197 Sathya M, Premkumar P, Karthick C, Moorthi P, Jayachandran KS, Anusuyadevi M. BACE1 in Alzheimer's disease. *Clin Chim Acta.* 2012; 414: 171-8.
- 198 Shimmyo Y, Kihara T, Akaike A, Niidome T, Sugimoto H. Flavonols and flavones as BACE-1 inhibitors: structure-activity relationship in cell-free, cell-based and in silico studies reveal novel pharmacophore features. *Biochim Biophys Acta.* 2008; 1780(5): 819-25.
- 199 Kumar NS, Nisha N. Phytomedicines as potential inhibitors of β amyloid aggregation: significance to Alzheimer's disease. *Chin J Nat Med.* 2014; 12(11): 801-818.
- 200 Anderson RL, Wolf WJ, Compositional changes in trypsin inhibitors, phytic acid, saponins and isoflavones related to soybean processing. *J Nutr.* 1995; 125(3): 581S-8S.
- 201 Karpagam V, Sathishkumar N, Sathiyamoorthy S, Rasappan P, Shila S, Kim YJ, *et al.* Identification of BACE1 inhibitors from Panax ginseng saponins-An Insilco approach. *Comput Biol Med.* 2013; 43(8): 1037-44.
- 202 Dhole VJ, Reddy KS. Genetic variation for phytic acid content in mungbean (*Vigna radiata L. Wilczek*). *The Crop Journal.* 2015; 3(2): 157-162.
- 203 Abe TK, Taniguchi M. Identification of myo-inositol hexakisphosphate (IP6) as a beta-secretase 1 (BACE1) inhibitory molecule in rice grain extract and digest. *FEBS Open Bio.* 2014; 4: 162-7.
- 204 Anand P, Singh B. Flavonoids as lead compounds modulating the enzyme targets in Alzheimer's disease. *Med Chem Res.* 2012; 22(7): 3061-75.

