

บทที่ 4
ผลการศึกษา

Table 1 Patient characteristics

Age- years (mean ±SD)	50.5 ± 2.4	56.6 ± 1.9	0.054
Gender (n,%)			1.000
Male	13 (65)	13 (65)	
Female	7 (35)	7 (35)	
BMI(kg/m2)	23.4±2.7	21.9±2.6	0.081
Operation time-mins (mean ±SD)	79.5±20	74±13.5	0.314
ASA (n,%)			0.504
I	7 (35)	5 (25)	
II	13 (65)	14 (70)	
III		1(5)	

Table 2 morphine use Nausea and Vomiting pain score and patient satification

Total amount of Morphine use (mg in 24 hrs) (mean ±SD)	3.9±3.7	5.6±3.7	0.121
Nausea and vomitng (n,%)	3(65%)	4(20%)	0.677
Numeric pain score(mean ±SD)			
0 hr	1.1±1.8	4.8 ±1.1	0.001
4 hr	2.2 ±1.9	4.3 ±1.7	<0.001
12 hr	3.8 ±2.0	3.4 ±2.1	0.513
24 hr	2.4 ±0.9	3.4 ±2.1	0.293
Satification (mean ±SD)	4.4±0.7	3.8±0.7	0.014

GA: General anesthesia

RA : Regional anesthesia

