

LIST OF TABLES

Table	Page
2.1 Cytokine abbreviations, sources, and functions.....	27
<u>Study I</u>	
4.1 The comparison of physiological characteristics and blood chemical data variables between the control group (CON) and the allergic rhinitis patients group (AR).....	69
4.2 Cytokine in serum and nasal secretion.....	70
<u>Study II</u>	
4.1 The comparison of resting heart rate (bpm) between pre- and post training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	73
4.2 The comparison of systolic blood pressure (mmHg) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	74
4.3 The comparison of diastolic blood pressure (mmHg) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	75
4.4 The comparison of body weight (kg) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	76
4.5 The comparison of body mass index (kg/m^2) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	77

Table	Page
4.6 The comparison of body fat (%) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	78
4.7 The comparison of maximum oxygen consumption; VO_2 max (ml/kg/min) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	79
4.8 The comparison of forced vital capacity; FVC (liters) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	80
4.9 The comparison of forced expiratory volume at 1 second; FEV1 (liters) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	81
4.10 The comparison of slow vital capacity; SVC (liters) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	82
4.11 The comparison of maximum voluntary ventilation; MVV (liters/min) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	83
4.12 The comparison of percent difference of the physiological characteristics variables among in control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	84
4.13 The comparison of white blood cell (cells/mm ³) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	86

Table	Page
4.14 The comparison of red blood cell (mcells/mm ³) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	87
4.15 The comparison of hemoglobin (g/dl) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	88
4.16 The comparison of hematocrit (%) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	89
4.17 The comparison of cholesterol (mg/dl) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	90
4.18 The comparison of triglyceride (mg/dl) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	91
4.19 The comparison of high density lipoprotein cholesterol; HDL-C (mg/dl) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	92
4.20 The comparison of low density lipoprotein cholesterol; LDL-C (mg/dl) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	93
4.21 The comparison of total Immunoglobulin E; Total IgE (IU/ml) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	94

Table	Page
4.22 The comparison of specific immunoglobulin E; specific IgE (<i>D.pteronyssinus</i>) (kUA/L) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	95
4.23 The comparison of plasma Vit C (mg/dl) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	96
4.24 The comparison of malondialdehyde; MDA ($\mu\text{mol/L}$) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	97
4.25 The comparison of percent difference of the blood chemical variables among in control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	98
4.26 The comparison of interleukin-2 (pg/ml) between pre- and post-training in three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	100
4.27 The comparison of percent difference of interleukin-2 after 5 minutes nasal challenge by house dust mite (<i>D.pteronyssinus</i>) between pre- and post-training in three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	101
4.28 The comparison of interleukin-4 (pg/ml) between pre- and post-training in three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	104

Table	Page
4.29 The comparison of percent difference of interleukin-4 after 5 minutes nasal challenge by house dust mite (<i>D.pteronyssinus</i>) between pre- and post-training in three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	105
4.30 The comparison of interleukin-13 (pg/ml) between pre- and post-training in three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	108
4.31 The comparison of percent difference of interleukin-13 after 5 minutes nasal challenge by house dust mite (<i>D.pteronyssinus</i>) between pre- and post-training in three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	109
4.32 The comparison of percent difference of cytokine levels in nasal secretion after 5 minutes nasal challenge by house dust mite (<i>D.pteronyssinus</i>) among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	112
4.33 The comparison of peak nasal inspiratory flow; PNIF (liter/sec) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	114
4.34 The comparison of PNIF after nasal challenge by house dust mite (<i>D.pteronyssinus</i>) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	115
4.35 The comparison of nasal blood flow (AU) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	118

Table	Page
4.36 The comparison of NBF after nasal challenge by house dust mite (<i>D.pteronyssinus</i>) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	120
4.37 The comparison of nasal congestion (level) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C)...	123
4.38 The comparison of itching (level) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	124
4.39 The comparison of sneezing (level) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	125
4.40 The comparison of rhinorrhea (level) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	126
4.41 The comparison of total rhinitis symptoms (level) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	127
4.42 The comparison of percent difference of the rhinitis symptoms variables among in control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	128
4.43 The comparison of nasal congestion after nasal challenge by house dust mite (<i>D.pteronyssinus</i>) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	130

Table	Page
4.44 The comparison of itching after nasal challenge by house dust mite (<i>D.pteronyssinus</i>) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	133
4.45 The comparison of sneezing after nasal challenge by house dust mite (<i>D.pteronyssinus</i>) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	136
4.46 The comparison of rhinorrhea after nasal challenge by house dust mite (<i>D.pteronyssinus</i>) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	139
4.47 The comparison of total rhinitis symptoms after nasal challenge by house dust mite (<i>D.pteronyssinus</i>) between pre- and post-training and among three groups of subjects: control group (CON), exercise group (EX) and exercise combined vitamin C supplementation group (EX + Vit. C).....	142