

REFERENCES

- Akbarsha, M. A., & Manivannan, B. (1993). Biochemical changes in the testis and male accessory organs of albino rats on treatment with *Andrographis paniculata* Nees. **Comparative Animal Physiology**, **28**(5), 421-426.
- _____, & Murugaian, P. (2000). Aspects of the male reproductive toxicity/male antifertility property of andrographolide in albino rats: effect on the testis and the cauda epididymidal spermatozoa. **Phytotherapy Research**, **14**(6), 432-435.
- _____, Manivannan, B., Hamid, K.S. & Vijayan, B. (1990). Antifertility effect of *Andrographis paniculata* (Nees) in male albino rats. **Indian Journal of Experimental Biology**, **28**(5), 421-426
- Akokuwah, G.A., Zhari, I., Mariam, A., & Yam, M.F. (2009). Absorption of andrographolides from *Andrographis paniculata* and its effect on CCl₄-induced oxidative stress in rats. **Food and Chemical Toxicology**, **47**(9), 2321-2326.
- _____, Zhari, I., Norhayati, I., & Mariam, A. (2006). HPLC and HPTLC densitometric determination of andrographolides and antioxidant potential of *Andrographis paniculata*. **Journal of Food Composition and Analysis**, **19**(2-3), 118-126.
- Allen, R.G., & Tresini, M. (2000). Oxidative stress and gene regulation. **Free Radical Biology and Medicine**, **28**(3), 463-499.
- Angus, W.G.R., Larsen, M.C., & Jefcoate, C.R., 1999. Expression of CYP1A1 and CYP1B1 depends on cell-specific factors in human breast cancer cell lines: role of estrogen receptor status. **Carcinogenesis**, **20**(6), 947-955.
- Barilla, J.M.S. (1999). *Andrographis paniculata: can herbs fight common ailments, cancer, and chronic viral infections?* New Canaan, Conn: Keats Publishing.

- Barker, C.W., Fagan, J.B., & Pasco, D.S. (1994). Down-regulation of P4501A1 and P4501A2 mRNA expression in isolated hepatocytes by oxidative stress. **Journal of Biological Chemistry**, **269**(6), 3985-3990.
- Barouki, R., & Morel, Y. (2001). Repression of cytochrome P450 1A1 gene expression by oxidative stress: mechanisms and biological implications. **Biochemical Pharmacology**, **61**(5), 511-516.
- Bartsch, H., Nair, U., Risch, A., Rojas, M., Wikman, H. & Alexandrov, K. (2000). Genetic polymorphism of CYP genes, alone or in combination, as a risk modifier of tobacco-related cancers, **Cancer Epidemiology, Biomarkers and Prevention**, **9**(1),3-28.
- Batkhuu, J., Hattori, K., Takano, F., Fushiya, S., Oshiman, K., & Fujimiya, Y. (2002). Suppression of NO production in activated macrophages *in vitro* and *ex vivo* by neoandrographolide isolated from *Andrographis paniculata*. **Biological and Pharmaceutical Bulletin**, **25**(9), 1169.
- Beischlag, T.V., & Perdew, G.H. (2005). ER α -AHR-ARNT protein-protein interactions mediate estradiol dependent transrepression of dioxin-inducible gene transcription. **Journal of Biological Chemistry**, **280**(22), 21607-21611.
- Blankenship, A., & Matsumara, F. (1997). 2,3,7,8,-Tetrachlorodibenzo- p-dioxin (TCDD)-induced activation of a protein tyrosine kinase, pp60Src in murine hepatic cytosol using a cell-free system. **Molecular Pharmacology**, **52**(4), 667-675.
- Brosens, J.J., & Parker, M.G. (2003). Oestrogen receptor hijacked. **Nature**, **423**(6939), 487-488.
- Burgos, R.A., Caballero, E.E., Sfinchez, N.S., Schroeder, R.A., Wikman G.K., & Hancke J.L. (1997). Testicular toxicity assesment of *Andrographis paniculata* dried extract in rats. **Journal of Ethnopharmacology**, **58**(3), 219-224.
- Caceres, D.D., Hancke, J.L., Burgos, R.A., & Wikman, G.K. (1997). Prevention of common colds with *Andrographis paniculata* dried extract: a pilot double blind trial. **Phytomedicine**, **4**(2), 101-104.

- Calabrese, C., Berman, S.H., Babish, J.G., Ma, X., Shinto, L., Dorr, M., et al. (2000). A phase I trial of andrographolide in HIV positive patients and normal volunteers. **Phytotherapy Research**, **14**(5), 333-338.
- Chandrasekaran, C.V., Thiyagarajan, P., Sundarajan, K., Goudar, K.S. Deepak, M., Murali, B., et al. (2009). Evaluation of the genotoxic potential and acute oral toxicity of standardized extract of *Andrographis paniculata* (KalmCold™). **Food and Chemical Toxicology**, **47**(8), 1892-1902.
- Chang, H.J., Park, J.S., Lee, E.K., Kim, M.H., Baek, M.K., Kim, H.R., et al. (2009). Ascorbic acid suppresses the 2,3,7,8-tetrachlorodibenzo- p-dioxin (TCDD)-induced CYP1A1 expression in human HepG2 cells. **Toxicology In vitro**, **23**(4), 622-626.
- Chang, K., Lii, C., Tsai, C., Yang, A., & Chen, H. (2008). Modulation of the expression of the p class of glutathione S-transferase by *Andrographis paniculata* extracts and andrographolide. **Food and Chemical Toxicology**, **46**(3), 1079-1088.
- Chao, W.W., & Lin, B.F. (2010). Review isolation and identification of bioactive compounds in *Andrographis paniculata* (Chuanxinlian). **Chinese Medicine**, **5**(1), 17.
- Chem, W., & Liang, X. (1982). Deoxyandrographolide 19 β -D-glucoside from the leaves of *Andrographis paniculata*. **Planta Medica**, **45**(2), 245-246.
- Chen, J., Hsiao, G., Lee, A., Wu, C., & Yen, M. (2004). Andrographolide suppresses endothelial cell apoptosis via activation of phosphatidyl inositol-3-kinase/Akt pathway. **Biochemical Pharmacology**, **67**(7), 1337-1345.
- Cheung, H.Y., Cheung, C.S., & Kong, C.K., (2001). Determination of bioactive diterpenoids from *Andrographis paniculata* by micellar electrokinetic chromatography. **Journal of Chromatography A**, **930**(1-2), 171-176.
- Chi, A.C., Appleton, K., Henriod, J.B., Krayner, J.W., Marlow, N.M., Bandyopadhyay, D., et al. (2009). Differential induction of CYP1A1 and CYP1B1 by benzo[a]pyrene in oral squamous cell carcinoma cell lines and by tobacco smoking in oral mucosa. **Oral Oncology**, **45**(11) 980-985.

- Chiou, W.F., Chen, C.F., & Lin, J.J. (2000). Mechanisms of suppression of inducible nitric oxide synthase (iNOS) expression in RAW264.7 cells by andrographolide. **British Journal of Pharmacology**, **129**(8), 1553-1560.
- _____, Lin, J.J., & Chen, C.F. (1998). Andrographolide suppresses the expression of inducible nitric oxide synthase in macrophage and restores the vasoconstriction in rat aorta treated with LPS. **British Journal of Pharmacology**, **125**(2), 327-334.
- Chomczynski, P., & Sacchi, N. (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. **Analytical Biochemistry**, **162**(1), 156-159.
- Choudhury, B.R., Haque, S.J., & Poddar M.K. (1987). *In vivo* and *in vitro* effects of kalmegh (*Andrographis paniculata*) extract and andrographolide on hepatic microsomal drug metabolizing enzymes. **Planta Medica**, **53**(2), 135-140.
- Ciolino, H.P., Wang, T.T., & Yeh, G.C. (1998). Diosmin and diosmetin are agonists of the aryl hydrocarbon receptor that differentially affect cytochrome P450 1A1 activity. **Cancer Research**, **58**(1), 197-206.
- Dalton, T., Sherzer, H. & Puga, A. (1999). Regulation of gene expression by reactive oxygen. **Annual Review of Pharmacology and Toxicology**, **39**(1), 67-101.
- Danielson, P.B. (2002). "The cytochrome P450 superfamily: biochemistry, evolution and drug metabolism in humans". **Current Drug Metabolism**, **3**(6), 561-97.
- Datta, A., & Sukul, N.C. (1982). Filaricidal properties of a wild herb, *Andrographis paniculata*. **Journal of Helminthology**, **56**(2), 81-84.
- Delescluse, C., Lemaire, G., Sousa, G., & Rahmani, R. (2000). Is CYP1A1 induction always related to AHR signaling pathway? **Toxicology**, **153**(1-3), 73-82.
- Deng, W.L. (1978). Preliminary studies on the pharmacology of *Andrographis* product dihydroandrographolide sodium succinate. **Newsletter Chinese Herbal Medicine**, **8**(1), 26-28.
- Eaton, D.L., Ramsdell, H.S., & Neal, G.E. (1994). Bio-transformation of aflatoxins. In D.L. Eaton & J.D. Groopman (Eds.). **The toxicology of aflatoxins, human health, veterinary and agricultural significance**. San Diego: Academic Press.

- Eisenbrand, G., & Tang, W. (1993). Food-borne heterocyclic amines chemistry, formation, occurrence and biological activities. **Toxicology**, **84**(1) 1-82.
- Eugster, H.P., Probst, M., Wurgler, F.E., & Sengstay, C. (1993). Caffeine, estradiol, and progesterone interact with human CYP1A1 and CYP1A2. Evidence from cDNA-directed expression in *Saccharomyces cerevisiae*. **Drug Metabolism and Disposition**, **21**(1), 43-49.
- Fabricant, D.S., & Farnsworth N.R. (2001). The value of plants used in traditional medicine for drug discovery. **Environmental Health Perspectives**, **109**(1), 69-75.
- Farber, E. (1984). The biochemistry of preneoplastic liver: a common metabolic pattern in hepatocyte nodules. **Canadian Journal of Biochemistry and Cell Biology**, **62**(6), 486-494.
- Fernandez-Salguero, P.M., Hilbert, D.M., Rudikoff, S., Ward, J.M., & Gonzalez, F.J. (1996). Aryl-hydrocarbon receptor-deficient mice are resistant to 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin-induced Toxicity. **Toxicology and Applied Pharmacology**, **140**(1), 173-179.
- Gallagher, E.P., Kunze, K.L., Stapleton, P.L., & Eaton D.L. (1996). The kinetics of aflatoxin B1 oxidation by human cDNA-expressed and human liver microsomal cytochromes P450 1A2 and 3A4. **Toxicology and Applied Pharmacology**, **141**(3), 595-606.
- George, J., Goodwin, B., Liddle, C., Tapner, M., & Farrell, G.C. (1997). Time-dependent expression of cytochrome P450 genes in primary cultures of well-differentiated human hepatocytes. **Journal of Laboratory and Clinical Medicine**, **129**(6), 638-648.
- Gleiter, C.H., & Gundert-Remy, U. (1996). Gender differences in pharmacokinetics. **European Journal of Drug Metabolism and Pharmacokinetics**, **21**(2), 123-128.
- Gonzalez, F.J., & Gelboin, H.V. (1994). Role of human cytochrome P450 in the metabolic activation of chemical carcinogens and toxins. **Drug Metabolism Reviews**, **26**(1-2), 165-183.

- Goth-Goldstein, R., Stampfer, M.R., Erdmann, C.A., & Russell, M. (2000). Interindividual variation in CYP1A1 expression in breast tissue and the role of genetic polymorphism. **Carcinogenesis**, **21**(11), 2119-2122.
- Guengerich, F.P. (1988). Carcinogenesis and Cancer Chemotherapy Roles of Cytochrome P-450 Enzymes in Chemical. **Cancer Research**, **48**, 2946-2954.
- _____. (2008). "Cytochrome p450 and chemical toxicology". **Chemical Research in Toxicology**, **21**(1), 70-83.
- _____, Wu, Z.L., & Bartleson, C. J. (2005). Function of human cytochrome P450s: Characterization of the remaining orphans. **Biochemical and Biophysical Research Communication**, **338**(1), 465-469.
- Hanapi, N.A., Azizi, J., Ismail, S. & Mansor, S.M. (2010). Evaluation of selected Malaysian medicinal plant on phase I drug metabolizing enzyme, CYP2C9, CYP2D6, and CYP3A4 activities *in vitro*. **International Journal of Pharmacology**, **6**(4), 494-499.
- Handa, S.S., & Sharma, A. (1990). Hepatoprotective activity of andrographolide from *Andrographis paniculata* against carbon tetrachloride. **Indian Journal of Medical Research**, **92**(2), 276-283.
- Harris, R.Z., Benet, L.Z., & Schwartz, J.B. (1995). Gender effects in pharmacokinetics and pharmacodynamics. **Drug**, **50**(2), 222-239.
- Hayashi, S., Watanabe, J., Nakachi, K., & Kawajiri, K. (1991). Genetic linkage of lung cancer-associated MspI polymorphisms with amino acid replacement in the heme binding region of the human Cytochrome P450 1A1 gene. **The Journal of Biochemistry**, **110**(3), 407-411.
- Hayes, J.D., Flanagan, J.U., & Jowsey, I.R. (2005). Glutathione transferases. **Annual Review of Pharmacology and Toxicology**, **45**(1), 51-88.
- Higashi, M.K., Veenstra, D.L., Kondo, L.M., Wittkowsky, A.K., Srinouanprachanh, S.L., Farin, F.M., et al. (2002). Association between CYP2C9 genetic variants and anticoagulation-related outcomes during warfarin therapy. **The Journal of the American Medical Association**, **287**(13), 1690-1698.

- Iruretagoneya, M.I., Tobar, J.A., Gonzalez, P.A., Sepulveda, S.E., Figueroa, C.A., Burgos, R.A., et al. (2005). Andrographolide interferes with T-cell activation and reduces experimental autoimmune encephalomyelitis in the mouse. **Journal of Pharmacology Experimental Therapeutics**, **312**(1), 366-372.
- Iwanari, M., Nakajima, M., Kizu, R., Hayakawa K., & Yokoi, T. (2002). Induction of CYP1A1, CYP1A2, and CYP1B1 mRNAs by nitropolycyclic aromatic hydrocarbons in various human tissue-derived cells: chemical-, cytochrome P450 isoform-, and cell-specific differences. **Archives of Toxicology**, **76**(5-6), 287-298.
- Jacquet, M., Lambert, V., Baudoux, E., Muller, M., Kremers, P., & Gielen, J. (1996). Correlation between P450 CYP1A1 inducibility, Msp1 genotype and lung cancer incidence. **European Journal of Cancer**, **32**(10), 1701-1706.
- Jain, D.C., Gupta, M.M., Saxena, S., & Kumar, S. (2000). LC analysis of hepatoprotective diterpenoids from *Andrographis paniculata*. **Journal of Pharmaceutical and Biomedical Analysis**, **22**(4), 705-709.
- Jaruchotikamol, A., Jarukamjorn, K., Sirisangtrakul, W., Sakuma, T., Kawasaki, Y., & Nemoto, N. (2007). Strong synergistic induction of CYP1A1 expression by andrographolide plus typical CYP1A inducers in mouse hepatocytes. **Toxicology and Applied Pharmacology**, **224**(2), 156-162.
- Jarukamjorn, K. (2008). *Andrographis paniculata*: a review of aspects of regulatory mechanisms of hepatic CYP1A enzymes. **Boletin Latinoamericano y del Caribe de Plantas Medicinales y Aromaticas**, **7**(2), 108-115.
- _____, Don-in, K., Makejaruskul, C., Laha, T., Daodee, S., Pearaksa, P., et al. (2006). Impact of *Andrographis paniculata* crude extract on mouse hepatic cytochrome P450 enzymes. **Journal of Ethnopharmacology**, **105**(3), 464-467.
- _____, Sakuma, T., & Nemoto, N. (2002). Sexual dimorphic expression of mouse hepatic CYP2B: alterations during development or after hypophysectomy. **Biochemical Pharmacology**, **63**(11), 2037-2041.
- Kapil, A., Koul, I.B., Banerjee, S.K., & Gupta, D. (1993). Antihepatotoxic effects of major constituents of *Andrographis paniculata*. **Biochemical Pharmacology**, **46**(1), 182-185.

- Kim, J.H., Stansbury, K.H., Walker, N.J., Trush, M.A., Strickland, P.T., & Sutter, T.R. (1998). Metabolism of benzo[a]pyrene and benzo[a]pyrene-7,8-diol by human cytochrome P450 1B1. **Carcinogenesis**, **19**(10), 1847-1853.
- Kim, J.Y., & Lee, S.M. (2006). Vitamins C and E protect hepatic cytochrome P450 dysfunction induced by polymicrobial sepsis. **European Journal of Pharmacology**, **534**(1-3), 202-209.
- Kimura, S., Gonzalez, F.J., & Nebert, D.W. (1986). Tissue-specific expression of the mouse dioxin-inducible P(1)450 and P(3)450 genes: Differential transcriptional activation and mRNA stability in liver and extrahepatic tissues. **Molecular and Cellular Biology**, **6**(5), 1471-1477.
- Kohle, C., Gschaidmeier, H., Lauth, D., Topell, S., Zitzer, K., & Bock, K.W. (1999). 2,3,7,8,-Tetrachlorodibenzo-p-dioxin (TCDD)-mediated membrane translocation of c-Src protein kinase in liver WB-F344 cells. **Archives Toxicology**, **73**(3), 152-158.
- Kumar, R.A., Sridevi, K., Kumar, N.V., Nanduri, S., & Rajagopal, S. (2004). Anticancer and immunostimulatory compounds from *Andrographis paniculata*. **Journal of Ethnopharmacology**, **92**(2-3), 3291-3295.
- Lai, K.P., Wong, M.H., & Wong, C.K.C. (2004). Modulation of AhR-mediated CYP1A1 mRNA and EROD activities by 17 β -estradiol and dexamethasone in TCDD-induced H411E cells. **Toxicological Sciences**, **78**(1), 41-49.
- Larsen-su, S.A., & Williams, D.E. (2001). Transplacental exposure to indole-3-carbinol induces sex-specific expression of cyp1a1 and cyp1b1 in the liver of fischer 344 neonatal rats. **Toxicological Sciences**, **64**(2), 162-168.
- LeCluyse, E.L., Madan, A., Hamilton, G., Carroll, K., DeHaan, R., & Parkinson, A. (2000). Expression and regulation of cytochrome P450 enzymes in primary cultures of human hepatocytes. **Journal of Biochemical Molecular Toxicology**, **14**(4), 177-188.
- Li, J., Cheung, H., Zhang, Z., Chan, G.K.L., & Fong, W. (2007). Andrographolide induces cell cycle arrest at G2/M phase and cell death in HepG2 cells via alteration of reactive oxygen species. **European Journal of Pharmacology**, **568**(1-3), 31-44.

- Liddle, C., Goodwin, B.J., George, J., Tapner, M., & Farrell, G.C. (1998). Separate and interactive regulation of cytochrome P450 3A4 by triiodothyronine, dexamethasone, and growth hormone in cultured hepatocytes. **The Journal of Clinical Endocrinology and Metabolism**, **83**(7), 2411-2416.
- Lili, J., Kaikai, S., Jun, L., Ying, C., Tianyu, L., & Zhengtao, W. (2009). Intracellular glutathione regulates andrographolide-induced cytotoxicity on hepatoma Hep3B cells. **Redox Report**, **14**(9), 176-184.
- Ma, Q., & Whitlock Jr., J.P. (1997). A novel cytoplasmic protein that interacts with the Ah receptor, contains tetratricopeptide repeat motifs, and augments the transcriptional response to 2,3,7,8-tetrachlorodibenzo-pdioxin. **The Journal of Biological Chemistry**, **272**(14), 8878-8884.
- Madav, S., Tripathi, C., Tandan, S.K., & Mishra, S. (1995). Analgesic, antipyretic, and antiulcerogenic effects of andrographolide. **Indian Journal of Pharmaceutical Sciences**, **57**(3), 121-125.
- Mandal, S.C., Dhara, A.K., & Maiti, B.C. (2001). Studies on psychopharmacological activity of *Andrographis paniculata* extract. **Phytotherapy Research**, **15**(3), 253-256.
- Matthews, J., & Gustafsson, J. (2006). Estrogen receptor and aryl hydrocarbon receptor signaling pathways. **Nuclear Receptor Signaling**, **4**(1), e016.
- Merawin, L.T., Arifah, A.K., Sani, R.A., Somchit, M.N., Zuraini, A., Ganabadi, S. et al. (2010). Screening of microfilaricidal effects of plant extracts against *Dirofilaria immitis*. **Research in Veterinary Science**, **88**(1), 142-147.
- Mkrtchyan, A., Panosyan, V., Panossian, A., Wikman, G., Wagner, H. (2005). A phase I clinical study of *Andrographis paniculata* fixed combination Kan Jang™ versus ginseng and valerian on the semen quality of healthy male subjects. **Phytomedicine**, **12**(6-7), 403-409.
- Mollerup, S., Ryberg, D., Hewer, A., Phillips, D.H., & Haugen, A. (1999). Sex differences in lung CYP1A1 expression and DNA adduct levels among lung cancer patients. **Cancer Research**, **59**, 3317-3320.
- Morel, Y., & Barouki, R. (1998). Down-regulation of cytochrome P450 1A1 gene promoter by oxidative stress. **The Journal of Biological Chemistry**, **273**(41), 26969-26976.

- Morel, Y., & Barouki, R. (1999). Repression of gene expression by oxidative stress. **Biochemical Journal**, **342**(3), 481-496.
- Murray, G.I., Taylor, V.E., McKay, J.A., Weaver, R.J., Ewen, S.W., Melvin, W.T., et al. (1995). The immunohistochemical localization of drug-metabolizing enzymes in prostate cancer. **The Journal of Pathology**, **177**(2), 147-152.
- Nakachi, K., Imai, K., Hayashi, S., & Kawajiri, K. (1993). Polymorphisms of the CYP1A1 and glutathione S-transferase genes associated with susceptibility to lung cancer in relation to cigarette dose in a Japanese population. **Cancer Research**, **53**, 2994-2999.
- Nagaraj, N.S., Beckers, S., Mensah, J.K., Waigel, S., Vigneswaran, N., & Zacharias, W. (2006). Cigarette smoke condensate induces cytochromes P450 and aldo-keto reductases in oral cancer cells. **Toxicology Letter**, **165**(2), 182-194.
- Nebert, D.W. (1989). The Ah locus: Genetic differences in toxicity, cancer, mutation, and birth defects. **Critical Review in Toxicology**, **20**(3), 153-174.
- _____, Dalton, T.P., Okey, A.B., & Gonzalez, F.J. (2004). Role of aryl hydrocarbon receptor-mediated induction of the CYP1 enzymes in environmental toxicity and cancer. **The Journal of Biological Chemistry**, **279**(23), 23847-23850.
- Nelson, D.R. (2009). The Cytochrome P450 Homepage. **Human Genomics**, **4**(1), 59-65.
- Nemoto, N., & Sakurai, J. (1992). Differences in regulation of gene expression between Cyp1a-1 and Cyp1a-2 in adult mouse hepatocytes in primary culture. **Carcinogenesis**, **13**(12), 2249-2254.
- _____, & Sakurai, J. (1995). Glucocorticoid and sex hormones as activating or modulating factors for expression of Cyp2b-9 and Cyp2b-10 in the mouse liver and hepatocytes. **Archives of Biochemistry and Biophysics**, **319**(1), 286-292.
- _____, Sakurai, J., Tazawa, A., & Ishikawa, T. (1989). Proline-dependent expression of aryl hydrocarbon hydroxylase in C57BL/6 mouse hepatocytes in primary culture. **Cancer Research**, **49**, 5863-5869.

- Nerurkar, P.V., Okinaka, L., Aoki, C., Seifried, A., Lum-Jones, A., Wilkens, L.R. et al. (2000). CYP1A1, GSTM1, and GSTP1 genetic polymorphisms and urinary 1-hydroxypyrene excretion in non-occupationally exposed individuals, **Cancer Epidemiology, Biomarkers and Prevention**, **9**, 1119-1122.
- NiceData Software. (2004). **Traditional Chinese Medicines Database Version 2.1**. Cambridge, Massachusetts: CambridgeSoft Corporation.
- Ohtake, F., Takeyama, K.I., Matsumoto, T., Kitagawa, H., Yamamoto, Y., Nohara, K., et al. (2003). Modulation of oestrogen receptor signaling by association with the activated dioxin receptor. **Nature**, **423**(6393), 545-550.
- Okey, A.B. (1990). Enzyme induction in the cytochrom P450 system. **Pharmacology and Therapeutics**, **45**(2), 241-298.
- Ooi, J.P., Kuroyanagi, M., Sulaiman, S.F., Muhammad, T.S.T., & Tan, M.L. (2011). Andrographolide and 14-deoxy-11, 12-didehydroandrographolide inhibit cytochrome P450s in HepG2 hepatoma cells. **Life Sciences**, **88**(9-10), 447-454.
- Pan, Y., Abd-Rashid, B.A., Ismail, Z., Ismail, R., Mak, J.W., Pook, P.C.K., et al. (2011a). *In vitro* modulatory effects of *Andrographis paniculata*, *Centella asiatica* and *Orthosiphon stamineus* on cytochrome P450 2C19 (CYP2C19). **Journal of Ethnopharmacology**, **133**(2), 881-887.
- _____, Abd-Rashid, B.A. Ismail, Z., Ismail, R., Mak, J.W., Pook, P.C.K., et al. (2011b). *In vitro* determination of the effect of *Andrographis paniculata* extracts and andrographolide on human hepatic cytochrome P450 activities. **Journal of Natural Medicines**, **65**(3-4), 440-447.
- Panossian, A., Hovhannisyan, A., Mamikonyan, G., Abrahamian, H., Hambardzumyan, E., Gabrielian, E., et al. (2000). Pharmacokinetic and bioavailability of andrographolide from *Andrographis paniculata* fixed combination Kan Jang in rats and human. **Phytomedicine**, **7**(5), 351-364.
- Paolini, M., Pozzetti, L., Pedulli, G.F., Marchesi, E., & Cantelli-Forti, G. (1999). The nature of prooxidant activity of vitamin C. **Life Sciences**, **64**(23), PL273-PL278.

- Patarapanich, C., Laungcholatan, S., Mahaverawat, N., Chaichantipayuth C., & Pummangura, S. (2007). HPLC determination of active diterpene lactones from *Andrographis paniculata* Nees planted in various seasons and regions in Thailand. **Thai Journal of Pharmaceutical Sciences**, **31**(1), 91-99.
- Pekthong, D., Blanchard, N., Abadie, C., Bonet, A., Heyd, B., Manton, G., et al. (2009). Effect of *Andrographis paniculata* extract and Andrographolide on hepatic cytochrome P450 mRNA expression and monooxygenase activities after *in vivo* administration to rats and *in vitro* in rat and human hepatocyte culture. **Chemico- Biological Interactions**, **179**(2-3), 247-255.
- _____, Martin, H., Abadie, C., Bonet, A., Heyd, B., Manton, G., et al. (2008). Differential inhibition of rat and human hepatic cytochrome P450 by *Andrographis paniculata* extract and andrographolide. **Journal of Ethnopharmacology**, **115**(3), 432-440.
- Perry, L.M. (1980). **Medicinal plants of East Southeast Asia: Attributed Properties and Uses**. Cambridge, MA: MIT Press.
- Pompella, A., Visvikis, A., Paolicchi, A., Tata, V.D., & Casini A.F. (2003). The changing faces of glutathione, a cellular protagonist. **Biochemical Pharmacology**, **66**(8), 1499-1503.
- Price, R.J., Scott, M.P., Walters, D.G., Stierum, R.H., Groten, J.P., Meredith, C., et al. (2004). Effect of thiabendazole on some rat hepatic xenobiotic metabolising enzymes. **Food and Chemical Toxicology**, **42**(6), 899-908.
- Puri, A., Axena, R., Saxena, R.P., Saxena, K.C., Srivastava, V., & Tandon, J.S. (1993). Immunostimulant agents from *Andrographis paniculata*. **Journal of Natural Products**, **56**(7), 995-999.
- Quinones, L., Lucas, D., Godoy, J., Cáceres, D., Berthou, F., Varela, N., et al. (2001). CYP1A1, CYP2E1 and GSTM1 genetic polymorphisms. The effect of single and combined genotypes on lung cancer susceptibility in Chilean people. **Cancer Letters**, **174**(1), 35-44.
- Rahman, N.N.A., Furuta, T., Kojima, S., Takane, K., & Mohd, M.A. (1999). Antimalarial activity of extracts of Malaysian medicinal plants. **Journal of Ethnopharmacology**, **64**(3), 249-254.

- Rajagopal, S., Kumar, R.A., Deevi, D.S., Satyanarayana, C., & Rajagopalan, R. (2003). Andrographolide, a potential cancer therapeutic agent isolated from *Andrographis paniculata*. **Journal of Experimental Therapeutics Oncology**, **3**(3), 147-158.
- Rau, S.E., Bend, J.R., Arnold, J.M.O., Tran, L.T., Spence, J.D., & Bailey, D.G. (1997). Grapefruit juice-terfenadine single-dose interaction: magnitude, mechanism, and relevance. **Clinical Pharmacology and Therapeutics**, **61**(2), 401-409.
- Ricci, M.S., Toscano, D.G., Mattingly, C.J., & Toscano, W.A. (1999). Estrogen receptor reduces CYP1A1 induction in cultured human endometrial cells. **Journal of Biological Chemistry**, **274**(6), 3430-3438.
- Roberts-Thompson, S.J., McManus, M.E., Tukey, R.H., Gonzalez, F.F., & Holder, G.M. (1993). The catalytic activity of four expressed human cytochrome P450s towards benzo[a]pyrene and the isomers of its proximate carcinogen. **Biochemical and Biophysical Research Communications**, **192**(3), 1373-1379.
- Royall, J.A., & Ischiropoulos, H. (1993). Evaluation of 2,7 dichlorofluorescein and dihydrorhodamine 123 as fluorescent probes for intracellular hydrogen peroxide in culture endothelial cells. **Archives of Biochemistry and Biophysics**, **302**(2), 348-355.
- Safe, S., & Wormke, M. (2003). Inhibitory aryl hydrocarbon receptor-estrogen receptor α cross-talk and mechanisms of action. **Chemical Research in Toxicology**, **16**(7), 807-816.
- Sakuma, T., Endo, Y., Mashino, M., Kuroiwa, M., Ohara, A., Jarukamjorn, K., et al. (2002). Regulation of the expression of two female-predominant CYP3A mRNAs (CYP3A41 and CYP3A44) in mouse liver by sex and growth hormones. **Archives of Biochemistry and Biophysics**, **404**(2), 234-242.
- _____, Ohtake, M., Katsurayama, Y., Jarukamjorn, K., & Nemoto, N. (1999). Induction of CYP1A2 by phenobarbital in the livers of aryl hydrocarbon-responsive and -nonresponsive mice. **Drug Metabolism and Disposition**, **27**(3), 379-384.

- Sato, K. (1989). Glutathione transferases as markers of preneoplasia and neoplasia. **Advances in Cancer Research**, **52**(1), 205-255.
- Sen, C.K., & Packer, L. (1996). Antioxidants and redox regulation of gene transcription. **Federation of American Society for Experimental Biology**, **10**(7), 709-720.
- Seow, A., Zhao, B., Lee, E.J.D., Poh, W., Teh, M., Eng, P., et al. (2001). Cytochrome P4501A2 (CYP1A2) activity and lung cancer risk: a preliminary study among Chinese women in Singapore. **Carcinogenesis**, **22**(4), 673-677.
- Shama, A., Lal, K., & Handa, S.S. (1992). Standardization of Indian crude drug Kalmegh by high pressure liquid chromatographic determination of andrographolide. **Phytochemical Analysis**, **3**(3), 129-131.
- Sharma, P.V. (1983). **Charaka Samhita Chankhambhia Orientalia, Vol. II**, Varanasi, India: Indian Medical Science Series, Sri Satguru Publications, a division of Indian Books Centre.
- Sheeja, K., Guruvayoorappan, C., & Kuttan, G. (2007). Antiangiogenic activity of *Andrographis paniculata* extract and andrographolide. **International Immunopharmacology**, **7**(2), 211-221.
- Shen, Y.C., Chen, C.F., & Chiou, W.F. (2000). Suppression of rat neutrophil reactive oxygen species production and adhesion by the diterpenoid lactone andrographolide. **Planta Medica**, **66**(4), 314.
- _____, Chen, C.F., & Chiou, W.F. (2002). Andrographolide prevents oxygen radical production by human neutrophils: possible mechanism(s) involved in its anti-inflammatory effect. **British Journal of Pharmacology**, **135**(2), 399-406.
- Shi, M.D., Lin, H.H., Chiang, T.A., Tsai L.Y., Tsai, S.M., Lee Y.C., et al. (2009). Andrographolide could inhibit human colorectal carcinoma Lovo cells migration and invasion via down-regulation of MMP-7 expression. **Chemico-Biological Interactions**, **180**(3), 344-352.
- _____, Lin, H.H., Lee, Y.C., Chao, J.K., Lin, R.A., & Chen, J.H. (2008). Inhibition of cell-cycle progression in human colorectal carcinoma Lovo cells by andrographolide. **Chemico-Biological Interactions**, **174**(3), 201-210.

- Shimada, T., & Fujii-Kuriyama, Y. (2004). Metabolic activation of polycyclic aromatic hydrocarbons to carcinogens by cytochromes P450 1A1 and 1B1. **Cancer Sciences**, **95**(1), 1-6.
- _____, Hayes, C.L., Yamazaki, H., Amin, S., Hecht, S.S., Guengerich, F.P., et al. (1996). Activation of chemically diverse procarcinogens by human cytochrome P450 1B1. **Cancer Research**, **56**, 2979-2984.
- _____, Gillam, E.M.J., Sutter, T.R., Strickland, P.T., Guengerich, F.P., & Yamazaki, H. (1997). Oxidation of xenobiotics by recombinant human cytochrome P450 1B1. **Drug Metabolism and Disposition**, **25**(5), 617-622.
- _____, Yamazaki, H., Mimura, M., Inui, Y., & Guengerich, F.P. (1994). Interindividual variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: studies with liver microsomes of 30 Japanese and 30 Caucasians. **Journal of Pharmacology and Experimental Therapeutics**, **270**(1), 414-23.
- Sigel, A., Sigel, H., & Sigel, R. (2007). **The Ubiquitous Roles of Cytochrome P450 Proteins: Metal Ions in Life Sciences**. New York: Wiley.
- Singh, R.P., Banerjee, S., & Rao, A.R. (2001). Modulatory influence of *Andrographis paniculata* on mouse hepatic and extrahepatic carcinogen metabolizing enzymes and antioxidation status. **Phytotherapy Research**, **15**(5), 382-390.
- Singha, P.K., Roy, S., & Dey, S. (2007). Protective activity of andrographolide and arabinogalactan proteins from *Andrographis paniculata* Nees. against ethanol-induced toxicity in mice. **Journal of Ethnopharmacology**, **111**(1), 13-21.
- Siripong, P., Kongkathip, B., Preechanokool, K., Picha, P., Tunsuwan, K., & Taylor, W.C. (1992). Cytotoxic diterpenoid constituents from *Andrographis paniculata* Nees. Leaves. **Journal of the Sciences Society of Thailand**, **18**(2), 187-194.
- Smith, P.L., Maloney, K.N., Pothen, R.G., Clardy J., & Clapham, D.E., (2006). Bisandrographolide from *Andrographis paniculata* activates TRPV4 channels. **The Journal of Biological Chemistry**, **281**(40), 29897-29904.

- Stucker, I., Jacquet, M., de Waziers, I., Cenee, S., Beaune, P., Kremers, P., et al. (2000). Relation between inducibility of CYP1A1, GSTM1 and lung cancer in a French population. **Pharmacogenetics**, **10**(7), 617-627.
- Sueyoshi, T., & Negishi, M. (2001). Phenobarbital response elements of cytochrome P450 genes and nuclear receptors. **Annual Review of Pharmacology and Toxicology**, **41**(1), 123-143.
- Sun, B., & Fukuhara, M. (1997). Effects of co-administration of butylated hydroxytoluene, butylated hydroxyanisole and flavonoids on the activation of mutagens and drug-metabolizing enzymes in mice. **Toxicology**, **122**(1-2), 61-72.
- The Geneva Foundation for Medical Education and Research. (2004). **Traditional Chinese Medicines Database**. Retrieved June 20, 2009, from http://www.gfmer.ch/TMCAM/TNCAM_database_system.htm.
- Trevigen, Inc. (2007). **HT Glutathione Assay Kit**. Retrieved December 3, 2009, from http://www.trevigen.com/docs/protocol_7511-100-K.pdf.
- Ueta, E., Suzuki, E., Nanba, E., Tadokoro, Y., Otsuka, Y., & Kurata, T. (2001). Regulation of cigarette smoke-induced cytochrome P4501A1 gene expression in osteogenic disorder Shionogi rat liver and in lung by large ascorbic acid dose. **Bioscience, Biotechnology and Biochemistry**, **65**(11), 2548-2551.
- Uno, S., Dalton, T.P., Derkenne, S., Curran, C.P., Miller, M.L., Shertzer, H.G., et al. (2004). Oral exposure to benzo[a]pyrene in the mouse: detoxication by inducible cytochrome P450 is more important than metabolic activation. **Molecular Pharmacology**, **65**(5), 1225-1237.
- Vecchini, F., Lenoir-Viale, M.C., Cathelineau, C., Magdalou, J., Bernard, B.A., & Shroot, B. (1994). Presence of a retinoid responsive element in the promoter region of the human cytochrome P4501A1 gene. **Biochemical and Biophysical Research and Communications**, **201**(3), 1205-1212.
- Verma, N., & Vinayak, M. (2008). Antioxidant action of *Andrographis paniculata* on lymphoma. **Molecular Biology Reports**, **35**(4), 535-540.
- Visen, P.K., Shukla, B., Patnaik, G.K., & Dhawan, B.N. (1993). Andrographolide protects rat hepatocytes against paracetamol-induced damage. **Journal of Ethnopharmacology**, **40**(2), 131-136.

- Williams, J.A., Hyland, R., Jones, B.C., Smith, D.A., Hurst, S., Goosen, T.C., et al. (2004). Drug drug interactions for UDP-glucuronosyltransferase substrates: A pharmacokinetic explanation for typically observed low exposure AUC_i/AUC ratios. **Drug Metabolism and Disposition**, **32**(11), 1201-1208.
- Williams, P.A., Cosme, J., Sridhar, V., Johnson, E.F., & McRee, D.E. (2000). The crystallographic structure of a mammalian microsomal cytochrome P450 monooxygenase: Structural adaptations for membrane binding and functional diversity. **Molecular Cell**, **5**(1), 121-132.
- Woo, A.Y.H., Waye, M.M.Y., Tsui, S.K.W., Yeung, S.T.W., & Cheng C.H.K. (2008). Andrographolide up-regulates cellular-reduced glutathione level and protects cardiomyocytes against hypoxia/reoxygenation injury. **The Journal of Pharmacology and Experimental Therapeutics**, **325**(1), 226-235.
- Xia, Y.F., Ye, B.Q., Li, Y.D., Wang, J.G., He, X.J., Lin, X., et al. (2004). Andrographolide attenuates inflammation by inhibition of NF-κB activation through covalent modification of reduced cysteine 62 of p50. **The Journal of Immunology**, **173**(6), 4207-4217.
- Xu, Y., Marshall, R.L., & Mukkur, T.K.S., (2006). An investigation on the antimicrobial activity of *Andrographis paniculata* extract and andrographolide *in vitro*. **Asian Journal of Plant Sciences**, **5**(3), 527-530.
- Yamazaki, H., Hatanaka, N., Kizu, R., Hayakawa, K., Shimada, N., Guengerich, F.P., et al. (2000). Bioactivation of diesel exhaust particle extracts and their major nitrated polycyclic aromatic hydrocarbon components, 1-nitropyrene and dinitropyrenes, by human cytochromes P450 1A1, 1A2, and 1B1. **Mutation Research**, **472**(1-2), 129-138.
- Yoon, K., Kim, J.H., Gil, H., Hwang, H., Hwangbo, B., & Lee J.S. (2008). CYP1B1, CYP1A1, MPO, and GSTP1 polymorphisms and lung cancer risk in never-smoking Korean women. **Lung Cancer**, **60**(1), 40-46.
- Yoshinari, K., Yoda, N., Toriyabe, T., & Yamazoe, Y. (2010). Constitutive androstane receptor transcriptionally activates human CYP1A1 and CYP1A2 genes through a common regulatory element in the 5'-flanking region. **Biochemical Pharmacology**, **79**(2), 261-269.

- Zaher, H., Yang, T.J., Gelboin, H.V., Fernandez-Salguero, .P, & Gonzalez, F.J. (1998). Effect of phenobarbital on hepatic CYP1A1 and CYP1A2 in the Ahr-null mouse. **Biochemical Pharmacology**, **55**(2), 235-238.
- Zaridah, M.Z., Idid, S.Z., Wan Omar, A., & Khozirah, S. (2001). *In vitro* antifilarial effects of three plant species against adult worms of subperiodic *Brugia malayi*. **Journal of Ethnopharmacology**, **78**(1), 79-84.
- Zhao, H.Y., & Fang, W.Y. (1991). Antithrombotic effect of *Andrographis paniculata* Nees in preventing myocardial infarction. **Chinese Medical Journal**, **104**(9), 770-775.
- Zhang, C.Y., & Tan, B.K. (1996). Hypotensive activity of aqueous extract of *Andrographis paniculata* in rats. **Clinical and Experimental Pharmacology and Physiology**, **23**(8), 675-678.
- _____, & Tan, B.K. (1997). Mechanisms of cardiovascular activity of *Andrographis paniculata* in the anaesthetized rat. **Journal of Ethnopharmacology**, **56**(2), 97-101.
- _____, & Tan, B.K. (1998). Vasorelaxation of rat thoracic aorta caused by 14-deoxyandrographolide. **Clinical and Experimental Pharmacology and Physiology**, **25**(6), 424-429.
- _____, & Tan, B.K. (1999). Effects of 14-deoxyandrographolide and 14-deoxy-11,12-didehydroandrographolide on nitric oxide production in cultured human endothelial cells. **Phytotherapy Research**, **13**(2), 157-159.
- _____, Kuroyangi, M., & Tan, B.K. (1998) Cardiovascular activity of 14-deoxy-11,12-didehydroandrographolide in the anaesthetised rat and isolated right atria. **Pharmacological Research**, **38**(6), 413-417.
- Zhang, Z., Chan, G.K., Li, J., Fong, W., & Cheung, H. (2008). Molecular interaction between andrographolide and glutathione follows second order kinetics. **Chemical and Pharmaceutical Bulletin**, **56**(9), 1229-1233.