

## REFERENCES

- Adwitiya, P., Ashwini, P., Avinash, A. K., Badri, R., Kajal, D., Vomsi, P., & Srividya, S. (2009). Mutagenesis of *Bacillus thuringiensis* IAM 12077 for increasing poly ( $\beta$ -) hydroxybutyrate (PHB) production. **Turkey journal of biology**, **33**, 225-230
- Almodares, A., & Hadi, M.R. (2009). Production of bioethanol from sweet sorghum : A review. **Journal of Agricultural Research**, **4**(9), 772 - 780.
- Anderson, A.J., & Dawes, E.A. (1990). Occurrence, metabolism, metabolic role, and industrial uses of bacterial polyhydroxyalkanoates. **Microbiology Review**, **54**, 450-72.
- Antonopoulou, G., Gavala, H.N., Skiadas, I.V., Angelopoulos, K., & Lyberatos, G. (2008). Biofuels generation from sweet sorghum: Fermentative hydrogen production and anaerobic digestion of the remaining biomass. **Bioresource Technology**, **99**, 110-119.
- Braunegg, G., Lefebvre, G., & Genser, K.L. (1998). Polyhydroxyalkanoates, biopolyesters from renewable resources: physiological and engineering aspects. **Journal of Biotechnology**, **65**, 127-161.
- Byrom, D. (1987). Polymer synthesis by microorganisms: Technology and Economics. **Trends Biotechnology**, **5**, 246-250.
- Chien, C.C., Chen, C.C., Choi, M.H., Kung, S.S., & Wei, Y.H. (2007). Production of poly- $\beta$ -hydroxybutyrate (PHB) by *Vibrio* spp. isolated from marine environment. **Journal of Biotechnology**, **132**, 259-263.
- Choi, J., & Lee, S.Y. (1999). Factors affecting the economics of polyhydroxy alkaonoate production by bacterial fermentation. **Applied Microbiology and Biotechnology**, **51**, 13-21.
- Choi, M.H., Song, J.J., & Yoon, S.C. (1995). Biosynthesis of copolyesters by *Hydrogenophaga pseudoflava* from various lactones. **Canadian Journal Microbiology**, **41**, 60-67.

- Doi, Y., Kunioka, M., Nakamura, Y., & Soga, K. (1986). Nuclear magnetic resonance studies on poly( $\beta$ -hydroxybutyrate) and a copolyester of  $\beta$ -hydroxybutyrate and  $\beta$ -hydroxyvalerate isolated from *Alcaligenes eutrophus* H16. **Macromolecules**, **19**, 2860-2864.
- Du, G., Chen, J., Yu, J., & Lun, S. (2001). Continuous production of poly-3-hydroxybutyrate by *Ralstonia eutropha* in a two-stage culture system. **Journal of Biotechnology**, **88**(1), 59-65.
- Edwards, U., Rogall, T., Bocker, H., Emde, M., & Bottger, E. (1989). Isolation and direct complete nucleotide determination of entire genes, characterization of a gene coding for 16S ribosomal DNA. **Nucleic Acids Research**, **17**, 7843-7853.
- Faccin, D.J.L., Martins, I., Cardozo, N.S.M., Rech, R., Ayub, M.A.Z., Alves, T.L.M., et al. (2009). Optimisation of C:N ratio and minimal initial carbon source for poly(3-hydroxybutyrate) production by *Bacillus megaterium*. **Journal of Chemical Technology and Biotechnology**, **84**, 1756-1761.
- Fernandez, D., Rodriguez, E., Bassas, M., & Vinas, M. (2005). Agro - industrial oily wastes as substrates for PHA production by the new strain *Pseudomonas aeruginosa* NCIB 40045: Effect of culture conditions. **Biochemistry Engineering Journal**, **26**, 159-167.
- Gouda, M.K., Swellam, A.E., & Omar, S.H. (2001). Production of PHB by a *Bacillus megaterium* strain using sugarcane molasses and corn steep liquor as sole carbon and nitrogen sources. **Microbiology Research**, **156**, 201-207.
- Grothe, E., Moo-Young, M., & Chisti, Y. (1999). Fermentation optimisation for the production of poly[ $\beta$ -hydroxybutyric acid) microbial thermoplastic. **Enzyme and Microbial Technology**, **25**, 132-141.
- Guan, L., Cho, K.H., & Lee, J.H. (2011). Analysis of the cultivable bacterial community in jeotgal, a Korean salted and fermented seafood, and identification of its dominant bacteria. **Food Microbiology**, **28**, 101-113.
- Halami, P.M. (2007). Production of polyhydroxyalkanoate from starch by the native isolate *Bacillus cereus* CFR06. **World Journal of Microbiology and Biotechnology**, **24**, 805-812.

- Hetényi, K., Gál, K., Németh, Á., & Sevela, B. (2010). Use of sweet sorghum juice for lactic acid fermentation: preliminary steps in a process optimisation. **Journal of Chemical Technology and Biotechnology**, **85**, 872-877.
- Ishizaki, A., Tanaka, K., & Taga, N. (2001). Microbial production of poly-D-3-hydroxybutyrate from CO<sub>2</sub>. **Applied Microbiology and Biotechnology**, **57**, 6-12.
- Jan, S., Roblot, C., Courtois, J., & Courtois, B. (1996). <sup>1</sup>H NMR spectroscopic determination of poly-3-hydroxybutyrate extracted from microbial biomass. **Enzyme Microbiology Technology**, **18**, 195-201.
- Jiang, Y., Song, X., Gong, L., Li, P., Dai, C., & Shao, W. (2008). High poly (β-hydroxybutyrate) production by *Pseudomonas fluorescens* A2a5 from inexpensive substrates. **Enzyme and Microbial Technology**, **42**, 167-172.
- Kaewkannetra, P., Tanonkeo, P., Tanamool, V., & Imai, T., (2008). Biorefinery of squeeze sweet sorghum juice into value added product of biopolymer. **Journal of Biotechnology**, 136, Supplement, S412.
- Katsura, K., Kawasaki, Potacharoen, H. W., Saono, S., Seki, T., Yamada, Y. et al. (2001). *Asaia siamensis* sp. nov., an acetic acid bacterium in the alpha-proteobacteria. **International Journal of Systematic and Evolutionary Microbiology**, **51**, 559-63.
- Khanna, S., & Srivastava, A.K., (2005). Statistical media optimisation studies for growth and PHB production by *Ralstonia eutropha*. **Process Biochemistry**, **40**, 2173-2182.
- Kim, B.S., Lee, S.C., Lee, S.Y., Chang, H.N., Chang, Y.K., & Woo, S.I. (1994). Production of poly(3-hydroxybutyric acid) by fed-batch culture of *Alcaligenes eutrophus* with glucose concentration control. **Biotechnology and Bioengineering**, **43**(9), 892-898.
- Kim, P., Kim, J.H., & Oh, D.-K. (2003). Improvement in cell yield of *Methylobacterium* sp. by reducing the inhibition of medium components for poly-β-hydroxybutyrate production. **World Journal of Microbiology and Biotechnology**, **19**, 357-361.

- Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. **Journal of Molecular Evolution**, **16**, 111-120.
- Koller, M., Bona, R., Chiellini, E., Fernandes, E.G., Horvat, P., Kutschera, C., et al. (2008). Polyhydroxyalkanoate production from whey by *Pseudomonas hydrogenovora*. **Bioresource Technology**, **99**, 4854-4863.
- Kominek, L.A., & Halvorson, H.O. (1965). Metabolism of poly-beta-hydroxybutyrate and acetoin in *Bacillus cereus*. **Journal of Bacteriology**, **90**, 1251-9.
- Kulpreecha, S., Boonruangthavorn, A., Meksiriporn, B., & Thongchul, N. (2009). Inexpensive fed-batch cultivation for high poly(3-hydroxybutyrate) production by a new isolate of *Bacillus megaterium*. **Journal of Bioscience and Bioengineering**, **107**, 240-245.
- Laopaiboon, L., Nuanpeng, S., Srinophakun, P., Klanrit, P., & Laopaiboon, P. (2009). Ethanol production from sweet sorghum juice using very high gravity technology: Effects of carbon and nitrogen supplementations. **Bioresource Technology**, **100**, 4176-4182.
- Law, J.H., & Slepecky, R.A. (1961). Assay of poly-beta-hydroxybutyric acid. **Journal of Bacteriology**, **82**, 33-36.
- Lee, K.M., Gilmore, D.F. (2005). Formulation and process modeling of biopolymer (polyhydroxyalkanoates: PHAs) production from industrial wastes by novel crossed experimental design. **Process Biochemistry**, **40**, 226-249.
- Lee, S., Ka, J.O., & Song, H.G. (2012). Growth promotion of *Xanthium italicum* by application of rhizobacterial isolates of *Bacillus aryabhattai* in microcosm soil. **Journal of Microbiology**, **50**, 45-49.
- Lemoigne, M. (1926). Products of dehydration and polymerization of the acid oxybutyric. **Bulletin Society Chemistry Biology**, **8**.
- Liang, Y., Sarkany, N., Cui, Y., Yesuf, J., Trushenski, J., & Blackburn, J.W. (2010). Use of sweet sorghum juice for lipid production by *Schizochytrium limacinum* SR21. **Bioresource Technology**, **101**, 3623-3627.
- Luengo, J.M., Garcia, B., Sandoval, A., Naharro, G., & Olivera, E.R. (2003). Bioplastics from microorganisms. **Current Opinion Microbiology**, **6**, 251-260.

- Mahmoudi, M., Sharifzadeh-Baei, M., Najafpour, G.D., Tabandeh, F., & Eisazadeh, H. (2010). Kinetic model for polyhydroxybutyrate (PHB) production by *Hydrogenophaga pseudoflava* and verification of growth conditions. **African Journal of Biotechnology**, **9**, 3151-3157.
- Mamma, D., Christakopoulos, P., Koullas, D., Kekos, D., Macris, B.J., & Koukios, E. (1995). An alternative approach to the bioconversion of sweet sorghum carbohydrates to ethanol. **Biomass and Bioenergy**, **8**, 99-103.
- Massoud, M.I., & Abd El-Razek, A. (2010). Suitability of *Sorghum bicolor* L. stalks and grains for bioproduction of ethanol. **Annals of Agricultural Sciences**, **56**, 83-87.
- Miyake, M., Miyamoto, C., Schnackenberg, J., Kurane, R., & Asada, Y. (2000). Phosphotransacetylase as a key factor in biological production of polyhydroxybutyrate. **Applied Biochemistry and Biotechnology**, **86**, 1039–1044.
- Modi, S., Koelling, K., & Vodovotz, Y. (2011). Assessment of PHB with varying hydroxyvalerate content for potential packaging applications. **European Polymer Journal**, **47**, 179-186.
- Mokhtari-Hosseini, Z.B., Vasheghani-Farahani, E., Heidarzadeh-Vazifekhoran, A., Shojaosadati, S.A., Karimzadeh, R., & Darani, K.K. (2009). Statistical media optimisation for growth and PHB production from methanol by a methylotrophic bacterium. **Bioresource Technology**, **100**, 2436-2443.
- Nakata, H.M. (1963). Effect of pH on Intermediates produced during growth and sporulation of *Bacillus cereus*. **Journal of Bacteriology**, **86**, 577-81.
- Oliveira, F.C., Dias, M.L., Castilho, L.R., & Freire, D.M.G. (2007). Characterization of poly(3-hydroxybutyrate) produced by *Cupriavidus necator* in solid-state fermentation. **Bioresource Technology**, **98**, 633-638.
- Ostle, A.G., & Holt, J.G. (1982). Nile blue A as a fluorescent stain for poly- $\beta$ -hydroxybutyrate. **Applied and Environmental Microbiology**, **44**, 238-241.
- Pachekoski, W.M., Agnelli, J.A.M., & Belem, L.P. (2009). Thermal, mechanical and morphological properties of poly (hydroxybutyrate) and polypropylene blends after processing. **Material Research**, **12**, 159-164.

- Palleroni, N.J. & Palleroni, A.V. (1978). *Alcaligenes latus*, a new species of hydrogen-utilizing bacteria. **International Journal of System Bacteriology**, **28**, 416–424.
- Pandian, S.R., Deepak, V., Kalishwaralal, K., Rameshkumar, N., Jeyaraj, M., & Gurunathan, S. (2010). Optimisation and fed-batch production of PHB utilizing dairy waste and sea water as nutrient sources by *Bacillus megaterium* SRKP-3. **Bioresource Technology**, **101**, 705-711.
- Philip, S., Keshavarz, T., & Roy, I. (2007). Polyhydroxyalkanoates: biodegradable polymers with a range of applications. **Journal of Chemical Technology and Biotechnology**, **82**, 233-247.
- Ramadas, N.V., Singh, S.K., Soccol, C.R., & Pandey, A. (2009). Polyhydroxybutyrate production using agro-industrial residue as substrate by *Bacillus sphaericus* NCIM 5149. **Brazilian Archives of Biology and Technology**, **52**, 17-23.
- Reddy, C.S.K., Ghai, R., & Kalia, V.C. (2003). Polyhydroxyalkanoates: an overview. **Bioresource Technology**, **87**, 137-146.
- Redzwan, G., Gan, S.N., & Tan, I.K.P. (1997). Isolation of polyhydroxyalkanoate-producing bacteria from an integrated-farming pond and palm-oil mill effluent ponds. **World Journal Microbiology Biotechnology**, **13**, 707-709.
- Saitou, N., & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. **Molecular Biology and Evolution**, **4**, 406-25.
- Santimano, M.C., Prabhu, N.N., & Garg, S. (2009). PHA production using low-cost agro-industrial wastes by *Bacillus* sp. strain COL1/A6. **Research Journal of Microbiology**, **4**, 89-96.
- Schlegel, H. G., Lafferty, R., & Krauss, I. (1970). The isolation of mutants not accumulating poly- $\beta$ -hydroxybutyric acid. **Archives of Microbiology**, **71**, 283-294.
- Shivaji, S., Chaturvedi, P., Begum, Z., Pindi, P.K., Manorama, R., Padmanaban, D. A., et al. (2009). *Janibacter hoylei* sp. nov., *Bacillus isronensis* sp. nov. and *Bacillus aryabhatai* sp. nov., isolated from cryotubes used for collecting air from the upper atmosphere. **International Journal of Systematic and Evolutionary Microbiology**, **59**, 2977-86.

- Sindhu, R., Ammu, B., Binod, P., Deepthi, S.K., Ramachandran, K.B., Soccol, C.R., et al. (2011). Production and characterization of poly-3-hydroxybutyrate from crude glycerol by *Bacillus sphaericus* NII 0838 and improving its thermal properties by blending with other polymers. **Brazilian Archives of Biology and Technology**, **54**, 783-794.
- Singh, M., Patel, S.K.S., & Kalia, V.C. (2009). *Bacillus subtilis* as potential producer for polyhydroxyalkanoates. **Microbial Cell Factories**, **8**(38), 1-11.
- Sipos, B., Reczey, J., Somorai, Z., Kadar, Z., Dienes, D., & Reczey, K. (2009). Sweet sorghum as feedstock for ethanol production: Enzymatic hydrolysis of steam-pretreated bagasse. **Applied Biochemistry and Biotechnology**, **153**, 151-162.
- Slater, S.C., Voige, W.H., & Dennis, D.E. (1988). Cloning and expression in *Escherichia coli* of the *Alcaligenes eutrophus* H16 poly-3-hydroxybutyrate biosynthetic pathway. **Bacteriology**, **170**, 4431-4436.
- Spiekermann, P., Rehm, B.H.A., Kalscheuer, R., Baumeister, D., & Steinbüchel, A. (1999). A sensitive, viable-colony staining method using Nile red for direct screening of bacteria that accumulate polyhydroxyalkanoic acids and other lipid storage compounds. **Archives of Microbiology**, **171**, 73-80.
- Steinbüchel, A., & Valentin, H.E. (1995). Diversity of bacterial polyhydroxyalkanoic acids. **FEMS Microbiology Letter**, **128**, 219-228.
- Suwannasing, W., Mooamart, S., & Keawkannetra, P. (2011). Yields of Polyhydroxyalkanoates (PHAs) during batch fermentation of sugar cane juice by *Alcaligenes latus* and *Alcaligenes eutrophus*. **Journal of Life Sciences**, **5**(11), 960-966.
- Tajima, K., Igari, T., Nishimura, D., Nakamura, M., Satoh, Y., & Munekata M. (2003). Isolation and characterization of *Bacillus sp* INT005 accumulating polyhydroxyalkanoate (PHA) from gas field soil. **Journal of Bioscience and Bioengineering**, **95**, 77-81.
- Tanamool, V., Imai, T., Danvirutai, P., & Kaewkannetra, P. (2011). Biosynthesis of polyhydroxyalkanoate (PHA) by *Hydrogenophaga sp.* isolated from soil environments during batch fermentation. **Journal of Life Sciences**, **5**(12), 1003-1012.

- Tanamool, V., & Kaewkannetra, P. (Eds). Direct screening of potential polyhydroxy alkananoates (PHAs) bacterial from soil environment using sweet sorghum as a sole carbon source. **Proceeding of the World Congress on Engineering and Technology (CET2011): October 28-30 2011, Wuhan University, Shanghai, China.** (pp. 397-400). Shanghai: Wuhan University.
- Tavares, L. Z., da Silva, E. S., & da Cruz Pradella, J. G. (2004). Production of poly(3-hydroxybutyrate) in an airlift bioreactor by *Ralstonia eutropha*. **Biochemical Engineering Journal**, **18**(1), 21-31.
- Thakur, P., Borah, B., Baruah, S., & Nigam, J. (2001). Growth-associated production of poly-3 hydroxybutyrate by *Bacillus mycoides*. **Folia Microbiology**, **46**, 488-494.
- Thirumala, M., Reddy, S., & Mahmood, S. (2010). Production and characterization of PHB from two novel strains of *Bacillus* spp. isolated from soil and activated sludge. **Journal of Industrial Microbiology and Biotechnology**, **37**, 271-278.
- Valappil, S.P., Peursum, D., Langley, G.J., Herniman, J.M., Boccaccini A.R., Bucke, C., et al. (2007). Polyhydroxyalkanoate (PHA) biosynthesis from structurally unrelated carbon sources by a newly characterized *Bacillus* spp. **Journal of Biotechnology**, **127**, 475–487.
- Vroman, I., & Tighzert, L. (2009). Review: biodegradation polymer. **Materials**, **2** (2), 307-344.
- Wang, F., & Lee, S.Y. (1997). Poly (3-hydroxybutyrate) production with high productivity and high polymer content by a fed-batch culture of *Alcaligenes latus* under nitrogen limitation. **Apply Environment Microbiology**, **63**, 3703-3706.
- Wang, J.G., & Bakken, L.R. (1998). Screening of soil bacteria for poly- $\beta$ -hydroxybutyric acid production and its role in the survival of starvation. **Microbiology Ecology**, **35**, 94-101.
- Wei, Y. H., Chen, W. C., Huang, C. K., Wu, H. S., Sun, Y. M., Lo, C. W., et al. (2011). Screening and Evaluation of Polyhydroxybutyrate-Producing Strains from Indigenous Isolate *Cupriavidus taiwanensis* Strains. **International Journal of Molecular Science**, **12**(1), 252-265.

- Willem, A., Busse, J., Goor, M., Pot, B., Falsen, E., Janzent, E., et al. (1989). *Hydrogenophaga*, a new genus of hydrogen-oxidizing bacteria that includes *Hydrogenophaga flava* comb. nov. (formerly *Pseudomonas flava*), *Hydrogenophaga palleronii* (formerly *Pseudomonas palleronii*), *Hydrogenophaga pseudoflava* (formerly *Pseudomonas pseudoflava* and *Pseudomonas carboxydoflava*), and *Hydrogenophaga taeniospiralis* (formerly *Pseudomonas taeniospiralis*). **System Bacteriology**, **39**, 319-333.
- Wu, Q., Huang, H. Hu, G.H., Chen, J., Ho, K.P., & Chen, G.Q. (2001). Production of poly-3-hydroxybutyrate by *Bacillus* sp. JMa5 cultivated in molasses media. **Antonie Van Leeuwenhoek**, **80**, 111-118.
- Xu, S., Luo, R., Wu, L., Xu, K., & Chen, G.O. (2006). Blending and characterizations of microbial poly (3-hydroxybutyrate) with dendrimers. **Journal Applied Polymer Science**, **102**, 3782-3790.
- Yeza, A., Halasz, A., Levadoux, W., & Hawari, J. (2007). Production of poly- $\beta$ -hydroxybutyrate (PHB) by *Alcaligenes latus* from maple sap. **Applied Microbiology and Biotechnology**, **77**, 269-274.
- Yilmaz, M., Soran, H., & Beyatli, Y. (2005). Determination of poly- $\beta$ -hydroxybutyrate (PHB) production by some *Bacillus* spp. **World Journal of Microbiology and Biotechnology**, **21**, 565-566.
- Yoon, S.C., & Choi, M.H. (1999). Local sequence dependence of polyhydroxyalkanoic acid degradation in *Hydrogenophaga pseudoflava*. **The Journal of Biology Chemistry**, **274**, 37800-37808.
- Yüksekdağ, Z. N., Aslım, B., Beyatl, Y., & Mercan, N. (2004). Effect of carbon and nitrogen sources and incubation times on poly-beta-hydroxybutyrate (PHB) synthesis by *Bacillus subtilis* 25 and *Bacillus megaterium* 12. **African Journal of Biotechnology**, **3**, 63-66.