

REFERENCES

- 1 Materska M, Perucka I. Antioxidant activity of the main phenolic compounds isolated from hot pepper fruit (*Capsicum annuum* L). *J Agric Food Chem.* 2005 Mar 9;53(5):1750-6.
- 2 Lee Y, Howard LR, VillalÓN B. Flavonoids and antioxidant activity of fresh pepper (*Capsicum annuum*) cultivars. *J Food Sci.* 1995;60(3):473-6.
- 3 Sun T, Xu Z, Wu CT, Janes M, Prinyawiwatkul W, No HK. Antioxidant Activities of Different Colored Sweet Bell Peppers (*Capsicum annuum* L.). *J Food Sci.* 2007;72(2):S98-S102.
- 4 Matsufuji H, Ishikawa K, Nunomura O, Chino M, Takeda M. Anti-oxidant content of different coloured sweet peppers, white, green, yellow, orange and red (*Capsicum annuum* L.). *J Food Sci Tech.* 2007;42(12):1482-8.
- 5 Mayne ST. Beta-carotene, carotenoids, and disease prevention in humans. *FASEB J.* 1996 May;10(7):690-701.
- 6 Stevenson DE, Hurst RD. Polyphenolic phytochemicals – just antioxidants or much more? *Cell Mol Life Sci.* 2007;64(22):2900-16.
- 7 Birari RB, Bhutani KK. Pancreatic lipase inhibitors from natural sources: unexplored potential. *Drug Discovery Today.* 2007;12(19-20):879-89.
- 8 Yoshioka M, St-Pierre S, Drapeau V, Dionne I, Doucet E, Suzuki M, *et al.* Effects of red pepper on appetite and energy intake. *Briti J Nutri.* 1999;82(02):115-23.
- 9 Menichini F, Tundis R, Bonesi M, Loizzo M, Conforti F, Statti G, *et al.* The influence of fruit ripening on the phytochemical content and biological activity of *Capsicum chinense* Jacq. cv Habanero. *Food Chem.* 2009;114(2):553-60.
- 10 Kwon Y-I, Apostolidis E, Shetty K. Evaluation of Pepper (*Capsicum annuum*) for Management of Diabetes and Hypertension. *J Food Biochem.* 2007;31(3):370-85.

- 11 Ranilla LG, Kwon Y-I, Apostolidis E, Shetty K. Phenolic compounds, antioxidant activity and *in vitro* inhibitory potential against key enzymes relevant for hyperglycemia and hypertension of commonly used medicinal plants, herbs and spices in Latin America. *Biore Techn.* 2010;101(12):4676-89.
- 12 Chaiyasit K, Khovidhunkit W, Wittayalertrpanya S. Pharmacokinetic and the effect of capsaicin in *Capsicum frutescens* on decreasing plasma glucose level. *J Med Assoc Thai.* 2009 Jan;92(1):108-13.
- 13 Yang D, Luo Z, Ma S, Wong WT, Ma L, Zhong J, *et al.* Activation of TRPV1 by dietary capsaicin improves endothelium-dependent vasorelaxation and prevents hypertension. *Cell Metabo.* 2010;12(2):130-41.
- 14 Meckelmann SW, Riegel DW, van Zonneveld MJ, Ríos L, Peña K, Ugas R, *et al.* Compositional Characterization of Native Peruvian Chili Peppers (*Capsicum* spp.). *J Agric Food Chem.* 2013;61(10):2530-7.
- 15 Guil-Guerrero JL, Martínez-Guirado C, del Mar Reboloso-Fuentes M, Carrique-Pérez A. Nutrient composition and antioxidant activity of 10 pepper (*Capsicum annuum*) varieties. *Europ Food Resea Techn.* 2006;224(1):1-9.
- 16 Rubio C, Hardisson A, Martin RE, Baez A, Martin MM, Alvazex R. Mineral composition of the red and green pepper (*Capsicum annuum*) from Tenerife Island. *Europ Food Resea Techn.* 2002;214(6):501-4.
- 17 Conforti F, Statti GA, Menichini F. Chemical and biological variability of hot pepper fruits (*Capsicum annuum* var. *acuminatum* L.) in relation to maturity stage. *Food Chem.* 2007;102(4):1096-104.
- 18 Deepa N, Kaur C, George B, Singh B, Kapoor HC. Antioxidant constituents in some sweet pepper (*Capsicum annuum* L.) genotypes during maturity. *LWT - Food Scien Techn.* 2007;40(1):121-9.
- 19 Howard LR, Talcott ST, Brenes CH, Villalon B. Changes in phytochemical and antioxidant activity of selected pepper cultivars (*Capsicum* Species) As influenced by maturity. *J Agric Food Chem.* 2000;48(5):1713-20.
- 20 Bae H, Jayaprakasha GK, Jifon J, Patil BS. Variation of antioxidant activity and the levels of bioactive compounds in lipophilic and hydrophilic extracts

- from hot pepper (*Capsicum* spp.) cultivars. Food Chem. 2012;134(4):1912-8.
- 21 Bureau of Non Communicable Disease. Non communicable disease. 2012. Available from: <http://thaincd.com/information-statistic/non-communicable-disease-data.php>.
 - 22 Social and Quality of Life Data Base System. The number and rate of mortality from major diseases. 2012. Available from: http://social.nesdb.go.th/SocialStat/StatReport_Final.aspx?reportid=367&template=1R2C&yeartype=M&subcatid=15.
 - 23 Grove KA, Sae-tan S, Kennett MJ, Lambert JD. (-)-Epigallocatechin-3-gallate inhibits pancreatic lipase and reduces body weight gain in high fat-fed obese mice. Obesity (Silver Spring). 2012;20(11):2311-3.
 - 24 da Rocha MB, Souza FV, dos Santos Estevam C, Pizza C, Sant'ana AE, Marcal RM. Antispasmodic effect of 4'-methylepigallocatechin on guinea pig ileum. Fitoterapia. 2012;83(7):1286-90.
 - 25 Yoshikawa M, Shimoda H, Nishida N, Takada M, Matsuda H. *Salacia reticulata* and its polyphenolic constituents with lipase inhibitory and lipolytic activities have mild antiobesity effects in rats. J Nutr. 2002;132(7):1819-1824.
 - 26 Zheng CD, Duan YQ, Gao JM, Ruan ZG. Screening for anti-lipase properties of 37 traditional Chinese medicinal herbs. J Chin Med Assoc. 2010 ;73(6):319-24.
 - 27 Shin JE, Joo Han M, Kim DH. 3-Methylethergalangin isolated from *Alpinia officinarum* inhibits pancreatic lipase. Biol Pharm Bull. 2003 ;26(6):854-7.
 - 28 Zhao HL, Sim JS, Shim SH, Ha YW, Kang SS, Kim YS. Antiobese and hypolipidemic effects of platycodin saponins in diet-induced obese rats: evidences for lipase inhibition and calorie intake restriction. Int J Obes (Lond). 2005;29(8):983-90.
 - 29 Yoshizumi K, Hirano K, Ando H, Hirai Y, Ida Y, Tsuji T, et al. Lupane-type saponins from leaves of *Acanthopanax sessiliflorus* and their inhibitory activity on pancreatic lipase. J Agric Food Chem. 2006;54(2):335-41.

- 30 Han LK, Kimura Y, Kawashima M, Takaku T, Taniyama T, Hayashi T, *et al.* Anti-obesity effects in rodents of dietary teasaponin, a lipase inhibitor. *Int J Obes Relat Metab Disord.* 2001;25(10):1459-64.
- 31 Zheng Q, Koike K, Han LK, Okuda H, Nikaido T. New biologically active triterpenoid saponins from *Scabiosa tschiliensis*. *J Nat Prod.* 2004;67(4):604-13.
- 32 Ninomiya K, Matsuda H, Shimoda H, Nishida N, Kasajima N, Yoshino T, *et al.* Carnosic acid, a new class of lipid absorption inhibitor from sage. *Bioorg Med Chem Lett.* 2004;14(8):1943-6.
- 33 Sheng L, Qian Z, Zheng S, Xi L. Mechanism of hypolipidemic effect of crocin in rats: crocin inhibits pancreatic lipase. *Eur J Pharmacol.* 2006;543(1-3):116-22.
- 34 Yoshioka M, Doucet E, Drapeau V, Dionne I, Tremblay A. Combined effects of red pepper and caffeine consumption on 24 h energy balance in subjects given free access to foods. *Brit J Nutri.* 2001;85(02):203-11.
- 35 Yu Q, Wang Y, Yu Y, Li Y, Zhao S, Chen Y, *et al.* Expression of TRPV1 in rabbits and consuming hot pepper affects its body weight. *Mol Biol Rep.* 2012;39(7):7583-9.
- 36 Kawada T, Hagihara K, Iwai K. Effects of capsaicin on lipid metabolism in rats fed a high fat diet. *J Nutr.* 1986;116(7):1272-8.
- 37 Diepvens K, Westerterp KR, Westerterp-Plantenga MS. Obesity and thermogenesis related to the consumption of caffeine, ephedrine, capsaicin, and green tea. 2007;292(1):R77-R85.
- 38 Kawabata F, Inoue N, Yazawa S, Kawada T, Inoue K, Fushiki T. Effects of CH-19 sweet, a non-pungent cultivar of red pepper, in decreasing the body weight and suppressing body fat accumulation by sympathetic nerve activation in humans. *Biosci Biotechnol Biochem.* 2006;70(12):2824-35.
- 39 Jeon G, Choi Y, Lee SM, Kim Y, Jeong HS, Lee J. Anti-obesity activity of methanol extract from hot pepper (*Capsicum annuum* L.) seeds in 3T3-L1 adipocyte. *Food Sci Biotechnol.* 2010;19(4):1123-7.

- 40 Ahn IS, Do MS, Kim SO, Jung HS, Kim YI, Kim HJ, et al. Antiobesity effect of Kochujang (Korean fermented red pepper paste) extract in 3T3-L1 adipocytes. *J Med Food*. 2006;9(1):15-21.
- 41 Cha YS, Kim SR, Yang JA, Back HI, Kim MG, Jung SJ, et al. Kochujang, fermented soybean-based red pepper paste, decreases visceral fat and improves blood lipid profiles in overweight adults. *Nutr Metab (Lond)*. 2013;10(1):24.
- 42 Kobayashi K, Ihara S, Kobata A, Itoh K, Kusunoki N, Yoshizaki F. Inhibitory effect of Myrica bark on lipase activity in mouse plasma and gastrointestinal tract. *J Med Food*. 2008;11(2):289-93.
- 43 Kang JH, Tsuyoshi G, Le Ngoc H, Kim HM, Tu TH, Noh HJ, et al. Dietary capsaicin attenuates metabolic dysregulation in genetically obese diabetic mice. *J Med Food*. 2011;14(3):310-5.
- 44 Yamashita K, Sugawara S, Sakairi I. Effects of an alpha-glucosidase inhibitor, acarbose, on blood glucose and serum lipids in streptozotocin-induced diabetic rats. *Horm Metab Res*. 1984;16(4):179-82.
- 45 Kumar S, Kumar V, MS C. *In-vitro* anti-oxidant and alpha-amylase inhibitory activity of isolated fraction from methanolic extract of *Asystasia dalzelliana* leaves. *Int J of PharmTech Res*. 2011;3(2):889-894.
- 46 Kwon YI, Son HJ, Moon KS, Kim JK, Kim JG, Chun HS, et al. Novel alpha-glucosidase inhibitors, CKD-711 and CKD-711a produced by *Streptomyces* sp. CK-4416. II. Biological properties. *J Antibiot (Tokyo)*. 2002;55(5):462-6.
- 47 He Q, Lv Y, Yao K. Effects of tea polyphenols on the activities of α -amylase, pepsin, trypsin and lipase. *Food Chemistry*. 2007;101(3):1178-82.
- 48 Schmidt JS, Lauridsen MB, Dragsted LO, Nielsen J, Staerk D. Development of a bioassay-coupled HPLC-SPE-ttNMR platform for identification of α -glucosidase inhibitors in apple peel (*Malus \times domestica* Borkh.). *Food Chem*. 2012;135(3):1692-9.
- 49 Wang Y, Yang Z, Wei X. Sugar compositions, α -glucosidase inhibitory and amylase inhibitory activities of polysaccharides from leaves and flowers of

- Camellia sinensis obtained by different extraction methods. *Int J Biol Macrom.* 2010;47(4):534-9.
- 50 Wu T, Zhou X, Deng Y, Jing Q, Li M, Yuan L. *In vitro* studies of *Gynura divaricata* (L.) DC extracts as inhibitors of key enzymes relevant for type 2 diabetes and hypertension. *J Ethnoph.* 2011;136(2):305-8.
- 51 Oboh G, Ademiluyi AO, Faloye YM. Effect of combination on the antioxidant and inhibitory properties of tropical pepper varieties against α -amylase and α -glucosidase activities *in vitro*. *J Med Food.* 2011;14(10):1152-8.
- 52 Ani V, Naidu KA. Antihyperglycemic activity of polyphenolic components of black/bitter cumin *Centratherum anthelminticum* (L.) Kuntze seeds. *Europ Food Res Techn.* 2008;226(4):897-903.
- 53 Kim JS, Kwon CS, Son KH. Inhibition of alpha-glucosidase and amylase by luteolin, a flavonoid. *Biosci Biotechnol Biochem.* 2000;64(11):2458-61.
- 54 Lee YA, Cho EJ, Tanaka T, Yokozawa T. Inhibitory activities of proanthocyanidins from persimmon against oxidative stress and digestive enzymes related to diabetes. *J Nutr Sci Vitaminol (Tokyo).* 2007;53(3):287-92.
- 55 Liu S, Li D, Huang B, Chen Y, Lu X, Wang Y. Inhibition of pancreatic lipase, α -glucosidase, α -amylase, and hypolipidemic effects of the total flavonoids from *Nelumbo nucifera* leaves. *J Ethnopharm.* 2013;149(1):263-9.
- 56 Tadera K, Minami Y, Takamatsu K, Matsuoka T. Inhibition of alpha-glucosidase and alpha-amylase by flavonoids. *J Nutr Sci Vitaminol (Tokyo).* 2006;52(2):149-53.
- 57 Lo Piparo E, Scheib H, Frei N, Williamson G, Grigorov M, Chou CJ. Flavonoids for controlling starch digestion: structural requirements for inhibiting Human α -Amylase. *J Med Chem.* 2008;51(12):3555-61.
- 58 Yan J, Zhang G, Pan J, Wang Y. α -Glucosidase inhibition by luteolin: Kinetics, interaction and molecular docking. *International J Biol Macrom.* 2014;64(0):213-23.
- 59 Loizzo MR, Tundis R, Menichini F, Statti GA, Menichini F. Influence of ripening stage on health benefits properties of *Capsicum*

- annuum* Var. *acuminatum* L.: *in vitro* studies. J Med Food. 2008;11(1):184-9.
- 60 Tundis R, Menichini F, Bonesi M, Conforti F, Statti G, Menichini F, *et al.* Antioxidant and hypoglycaemic activities and their relationship to phytochemicals in *Capsicum annuum* cultivars during fruit development. LWT - Food Scien Techn. 2013;53(1):370-7.
- 61 Burnier M, Biollaz J. Pharmacokinetic optimisation of angiotensin converting enzyme (ACE) inhibitor therapy. Clin Pharmacokin. 1992;22(5):375-84.
- 62 Kumar S, Maurya DR, Chandra M. ACE Inhibition versus angiotensin-II antagonism in heart failure. Asian Cardiova Thorac Annals. 2000;8(2):195-203.
- 63 Hong F, Ming L, Yi S, Zhanxia L, Yongquan W, Chi L. The antihypertensive effect of peptides: A novel alternative to drugs? Peptides. 2008;29(6):1062-71.
- 64 Li G-H, Le G-W, Shi Y-H, Shrestha S. Angiotensin I-converting enzyme inhibitory peptides derived from food proteins and their physiological and pharmacological effects. Nutri Res. 2004;24(7):469-86.
- 65 Ojeda D, Jimenez-Ferrer E, Zamilpa A, Herrera-Arellano A, Tortoriello J, Alvarez L. Inhibition of angiotensin convertin enzyme (ACE) activity by the anthocyanins delphinidin- and cyanidin-3-O-sambubiosides from *Hibiscus sabdariffa*. J Ethnopharmacol. 2010;127(1):7-10.
- 66 Dong J, Xu X, Liang Y, Head R, Bennett L. Inhibition of angiotensin converting enzyme (ACE) activity by polyphenols from tea (*Camellia sinensis*) and links to processing method. Food Funct. 2011;2(6):310-9.
- 67 Lacaille-Dubois MA, Franck U, Wagner H. Search for potential angiotensin converting enzyme (ACE)-inhibitors from plants. Phytomedicine. 2001;8(1):47-52.
- 68 Loizzo MR, Said A, Tundis R, Rashed K, Statti GA, Hufner A, *et al.* Inhibition of angiotensin converting enzyme (ACE) by flavonoids isolated from *Ailanthus excelsa* (Roxb) (Simaroubaceae). Phytotherapy Res. 2007;21(1):32-6.

- 69 Berry SE, Mulla UZ, Chowienczyk PJ, Sanders TAB. Increased potassium intake from fruit and vegetables or supplements does not lower blood pressure or improve vascular function in UK men and women with early hypertension: a randomised controlled trial. *Brit J Nutri.* 2010;104(12):1839-47.
- 70 Judprasong K, Charoenkiatkul, S., Thiyajai, P., Sukprasansap, M. Nutrients and bioactive compounds of Thai indigenous fruits. *Food Chem.* 2013;140:507-12.
- 71 Azar A, Nekoei M, Larijani K, Bahraminasab S. Chemical composition of the essential oils of *Citrus sinensis* cv. *valencia* and a quantitative structure-retention relationship study for the prediction of retention indices by multiple linear regression. *Journal of the Serbian Chemical Society.* 2011;76(12):1627-37.
- 72 Choi SJ, Hwang JM, Kim SI. A colorimetric microplate assay method for high throughput analysis of lipase activity. *J Biochem Mol Biol.* 2003;36(4):417-20.
- 73 Ingrid F, Matthias FM. Traditionally used plants in diabetes therapy: phytotherapeutics as inhibitors of alpha-amylase activity. *Sociedade Brasileira de Farmacognosia.* 2006;16(1).
- 74 You Q, Chen F, Wang X, Luo PG, Jiang Y. Inhibitory effects of muscadine anthocyanins on alpha-glucosidase and pancreatic lipase activities. *J Agric Food Chem.* 2011;59(17):9506-11.
- 75 Schwager SL, Carmona AK, Sturrock ED. A high-throughput fluorimetric assay for angiotensin I-converting enzyme. *Nat Protoc.* 2006;1(4):1961-4.
- 76 Frank CA, Nelson RG, Simonne EH, Behe BK, Simonne AH. Consumer preferences for color, price, and vitamin C content of bell peppers. *HortScience.* 2001;36(4):795-800.
- 77 Simonne AH, Simonne EH, Eitenmiller RR, Mills HA, Green NR. Ascorbic acid and provitamin A contents in unusually colored bell peppers (*Capsicum annuum* L.). *J Food Composit Anal.* 1997;10(4):299-311.
- 78 Stahl W, Sies H. Antioxidant activity of carotenoids. *Molec Spects Med.* 2003;24(6):345-51.

- 79 Winklhofer-Roob BM, Rock E, Ribalta J, Shmerling DH, Roob JM. Effects of vitamin E and carotenoid status on oxidative stress in health and disease. Evidence obtained from human intervention studies. *Molec Spects Med.* 2003;24(6):391-402.
- 80 Russo VM, Howard LR. Carotenoids in pungent and non-pungent peppers at various developmental stages grown in the field and glasshouse. *J Scien Food Agri.* 2002;82(6):615-24.
- 81 Kaur C, Kapoor HC. Antioxidants in fruits and vegetables – the millennium's health. *Int J Food Scien Techn.* 2001;36(7):703-25.
- 82 Marín A, Ferreres F, Tomás-Barberán FA, Gil MI. Characterization and quantitation of antioxidant constituents of sweet pepper (*Capsicum annuum* L.). *J Agric Food Chem.* 2004;52(12):3861-9.
- 83 Kim J-S, Ahn J, Lee S-J, Moon B, Ha T-Y, Kim S. Phytochemicals and antioxidant activity of fruits and leaves of paprika (*Capsicum annuum* L., var. Special) cultivated in Korea. *J Food Scien.* 2011;76(2):C193-C8.
- 84 Maoka T, Mochida K, Kozuka M, Ito Y, Fujiwara Y, Hashimoto K, *et al.* Cancer chemopreventive activity of carotenoids in the fruits of red paprika *Capsicum annuum* L. *Cancer Letters.* 2001;172(2):103-9.
- 85 Howard LK, Wildman REC. Antioxidant vitamin and phytochemical content of fresh and processed pepper fruit (*Capsicum annuum*). *Handbook of Nutraceutu Func Foods.* 2nd ed. 2006:165-191.
- 86 Croft KD. The chemistry and biological effects of flavonoids and phenolic acids. *Annals of the New York Academy of Sciences.* 1998;854(1):435-42.
- 87 Sergent T, Vanderstraeten J, Winand J, Beguin P, Schneider Y-J. Phenolic compounds and plant extracts as potential natural anti-obesity substances. *Food Chem.* 2012;135(1):68-73.
- 88 Actis-Goretta L, Ottaviani JJ, Fraga CG. Inhibition of angiotensin converting enzyme activity by flavanol-rich foods. *J Agric Food Chem.* 2006;54(1):229-34.
- 89 Stagg GV, Millin DJ. The nutritional and therapeutic value of tea—a review. *J Scien of Food Agric.* 1975;26(10):1439-59.

- 90 Ghasemnezhad M, Sherafati M, Payvast GA. Variation in phenolic compounds, ascorbic acid and antioxidant activity of five coloured bell pepper (*Capsicum annum*) fruits at two different harvest times. *J Func Foods*. 2011;3(1):44-9.
- 91 Miean KH, Mohamed S. Flavonoid (myricetin, quercetin, kaempferol, luteolin, and apigenin) content of edible tropical plants. *J Agric Food Chem*. 2001;49(6):3106-12.
- 92 Govindarajan VS, Salzer UJ. Capsicum-production, technology, chemistry, and quality part 1: History, botany, cultivation, and primary processing. *C R C Critical Reviews in Food Science and Nutrition*. 1985;22(2):109-76.
- 93 Luning PA, van der Vuurst de Vries R, Yuksel D, Ebbenhorst-Seller T, Wichers HJ, Roozen JP. Combined instrumental and sensory evaluation of flavor of fresh bell peppers (*Capsicum annuum*) Harvested at Three Maturation Stages. *J Agri Food Chem*. 1994;42(12):2855-61.
- 94 Forero MD, Quijano CE, Pino JA. Volatile compounds of chile pepper (*Capsicum annuum* L. var. *glabriusculum*) at two ripening stages. *J Flav Fragr*. 2009;24(1):25-30.
- 95 Eggink PM, Maliepaard C, Tikunov Y, Haanstra JPW, Bovy AG, Visser RGF. A taste of sweet pepper: Volatile and non-volatile chemical composition of fresh sweet pepper (*Capsicum annuum*) in relation to sensory evaluation of taste. *Food Chem*. 2012;132(1):301-10.
- 96 Luning PA, de Rijk T, Wichers HJ, Roozen JP. Gas Chromatography, Mass Spectrometry, and Sniffing Port Analyses of Volatile Compounds of Fresh Bell Peppers (*Capsicum annuum*) at Different Ripening Stages. *J Agri Food Chem*. 1994;42(4):977-83.
- 97 Yu C-W, Li W-H, Hsu F-L, Yen P-L, Chang S-T, Liao VH-C. Essential Oil alloaromadendrene from mixed-type *Cinnamomum osmophloeum* leaves prolongs the lifespan in *Caenorhabditis elegans*. *J Agric Food Chem*. 2014;62(26):6159-65.
- 98 Bryant RJ, McClung AM. Volatile profiles of aromatic and non-aromatic rice cultivars using SPME/GC-MS. *Food Chem*. 2011;124(2):501-13.

- 99 Lekshmi PC, Arimboor R, Indulekha PS, Nirmala Menon A. Turmeric (*Curcuma longa* L.) volatile oil inhibits key enzymes linked to type 2 diabetes. *Int J Food Scien Nutri*. 2012;63(7):832-4.
- 100 Raina VK, Srivastava SK, Jain N, Ahmad A, Syamasundar KV, Aggarwal KK. Essential oil composition of *Curcuma longa* L. cv. *Roma* from the plains of northern India. *J Flav Fragra*. 2002;17(2):99-102.
- 101 Karamac M, Amarowicz R. Inhibition of pancreatic lipase by phenolic acids--examination *in vitro*. *Z. Naturforsch*. 1996;51:903-905.
- 102 Van Buggenhout S, Alming M, Lemmens L, Colle I, Knockaert G, Moelants K, *et al*. *In vitro* approaches to estimate the effect of food processing on carotenoid bioavailability need thorough understanding of process induced microstructural changes. *Trends in Food Scien Techn*. 2010;21(12):607-18.
- 103 Proteopedia. Lipase. 2012, Available from: http://www.proteopedia.org/wiki/index.php/Sandbox_51.
- 104 Enzymology at interfaces and physiology of lipolysis. Inhibition of gastrointestinal lipolysis by orlistat during digestion of test meals in healthy volunteers. 2001. Available from: <http://eipl.cnrs-mrs.fr/news.php?newsid=2>.
- 105 Sternby B, Hartmann D, Borgström B, Nilsson Å. Degree of *in vivo* inhibition of human gastric and pancreatic lipases by orlistat (Tetrahydrolipstatin, THL) in the stomach and small intestine. *Clinical Nutri*. 2002;21(5):395-402.
- 106 Li Y-Q, Yang P, Gao F, Zhang Z-W, Wu B. Probing the interaction between 3 flavonoids and pancreatic lipase by methods of fluorescence spectroscopy and enzymatic kinetics. *Europ Food Resea Techn*. 2011;233(1):63-9.
- 107 Funke I, Melzig MF. Effect of different phenolic compounds on α -amylase activity: screening by microplate-reader based kinetic assay. *Die Pharmazie - An Int J Pharma Scien*. 2005;60(10):796-7.
- 108 Ohnuki K, Haramizu S, Oki K, Watanabe T, Yazawa S, Fushiki T. Administration of capsiate, a non-Pungent capsaicin analog, promotes

- energy metabolism and suppresses body fat accumulation in mice. *Biosci, Biotechn, and Biochem.* 2001;65(12):2735-40.
- 109 Iida T, Moriyama T, Kobata K, Morita A, Murayama N, Hashizume S, *et al.* TRPV1 activation and induction of nociceptive response by a non-pungent capsaicin-like compound, capsiate. *Neuropharmacology.* 2003;44(7):958-67.
- 110 MacGrego EA, Janecek S, Svensson B. Relationship of sequence and structure to specificity in the α -amylase family of enzymes. 2001;1546:1-20.
- 111 Nielsen JE, Beier L, Otzen D, Borchert TV, Frantzen HB, Andersen KV, *et al.* Electrostatics in the active site of an α -amylase. *Europ J Biochem.* 1999;264(3):816-24.
- 112 Li C, Begum A, Numao S, Park KH, Withers SG, Brayer GD. Acarbose rearrangement mechanism implied by the kinetic and structural analysis of human pancreatic α -amylase in complex with analogues and their elongated counterparts. *Biochemistry.* 2005;44(9):3347-57.
- 113 Aghajari N, Roth M, Haser R. Crystallographic evidence of a transglycosylation reaction: ternary complexes of a psychrophilic α -amylase. *Biochem.* 2002;41(13):4273-80.
- 114 Mahmud T. The C7N aminocyclitol family of natural products. *Natur Produ Repor.* 2003;20(1):137-66.
- 115 Hallmann E, Rembiałkowska E. Characterisation of antioxidant compounds in sweet bell pepper (*Capsicum annuum* L.) under organic and conventional growing systems. *J Scien Food Agri.* 2012;92(12):2409-15.
- 116 Adisakwattana S, Chantarasinlapin P, Thammarat H, Yibchok-Anun S. A series of cinnamic acid derivatives and their inhibitory activity on intestinal α -glucosidase. *J Enzyme Inhibit Med Chem.* 2009;24(5):1194-200.
- 117 Brás NF, Fernandes PA, Ramos MJ, Cerqueira NMFS. Glycosidases - a mechanistic overview. 2012.
- 118 Kim M-J, Lee S-B, Lee H-S, Lee S-Y, Baek J-S, Kim D, *et al.* Comparative study of the inhibition of α -glucosidase, α -amylase, and cyclomaltodextrin glucanotransferase by acarbose, isoacarbose, and acarviosine-glucose. *Archiv Biochem Biophys.* 1999;371(2):277-83.

- 119 Roujeinikova A, Raasch C, Sedelnikova S, Liebl W, Rice DW. Crystal structure of thermotoga maritima 4- α -glucanotransferase and its acarbose complex: implications for substrate specificity and catalysis. *J Molecu Bio.* 2002;321(1):149-62.
- 120 Kameda K, Takaku T, Okuda H, Kimura Y, Okuda T, Hatano T, et al. Inhibitory effects of various flavonoids isolated from leaves of persimmon on angiotensin-converting enzyme activity. *J Natur Products.* 1987 ;50(4):680-3.
- 121 Oh H, Kang D-G, Kwon J-W, Kwon T-O, Lee S-Y, Lee D-B, *et al.* Isolation of angiotensin converting enzyme (ACE) inhibitory flavonoids from *Sedum sarmentosum*. *Biol Pharm Bull.* 2004;27(12):2035-7.
- 122 Guerrero L, Castillo J, Quiñones M, Garcia-Vallvé S, Arola L, Pujadas G, *et al.* Inhibition of angiotensin-converting enzyme activity by flavonoids: structure-activity relationship studies. *PLoS One.* 2012;7(11):e49493.
- 123 Kayahara H, Miao Z, Fujiwara G. Synthesis and biological activities of ferulic acid derivatives. *Anticancer Res.* 1999;19(5A):3763-8.
- 124 Pinto MDS, Ranilla LG, Apostolidis E, Lajolo FM, Genovese MI, Shetty K. Evaluation of antihyperglycemia and antihypertension potential of native peruvian fruits using *in vitro* models. *J Med Food.* 2009;12(2):278-91.
- 125 Zhang C, Wu s, XU D. Catalytic mechanism of angiotensin-converting enzyme and effects of the chloride ion. *J phys chem B.* 2013;117:6635-6645.
- 126 Wei L, Clauser E, Alhenc-Gelas F, Corvol P. The two homologous domains of human angiotensin I-converting enzyme interact differently with competitive inhibitors. *J Biol Chem.* 1992;267(19):13398-405.
- 127 Kim MH, Shin DR, Yoo OJ, Lee H, Lee JO. Crystal structure of *Drosophila* angiotensin I-converting enzyme bound to captopril and lisinopril. *Elsevier.* 2003;538:65-70.
- 128 Nileeka Balasuriya BW, Vasantha Rupasinghe HP. Plant flavonoids as angiotensin converting enzyme inhibitors in regulation of hypertension. *Func Food Health Disea.* 2001;1(5):172-188.

- 129 Chen C-H, Lin J-Y, Lin C-N, Hsu S-Y. Inhibition of angiotensin-I-converting enzyme by tetrahydroxyxanthenes isolated from tripterospermum lanceolatum. *J Natur Produ.* 1992;55(5):691-5.