

References

1. Biesalski HK, Nohr D. Importance of vitamin-A for lung function and development. *Mol Aspects Med.*2003; 24: 431–40
2. Metzler MD, Snyder JM. Retinoic acid differentially regulates expression of surfactant associated proteins in human fetal lung. *Endocrinology* 1993; 133: 1990–8
3. Verma RP, McCulloch KM, Worrell L, Vidyasagar D. Vitamin A deficiency and severe bronchopulmonary dysplasia in very low birthweight infants. *Am J Perinatol* 1996; 13:389–93
4. Zachman RD, Samuels DP, Brand JM, Winston JF, Pi JT. Use of the intramuscular relative-dose-response test to predict bronchopulmonary dysplasia in premature infants. *Am J Clin Nutr* 1996; 63:123–9
5. Bancalari E, Claure N, Sosenko IR. Bronchopulmonary dysplasia: changes in pathogenesis, epidemiology and definition. *Semin Neonatol* 2003; 8:63–71
6. Silvers KM, Gibson AT, Powers HJ. The predictive value of plasma antioxidant activity at birth for morbidity and mortality of premature babies. *Proc Nutr Soc* 1993; 52: 313A

7. Frank L, Groseclose EE. Preparation for birth into an O₂- rich environment: The antioxidant enzyme in developing rabbit lung. *Pediatr* 1984; 18: 240-4
8. Shenai JP, Chytil F, Jhaveri A, Stahlman MT. Plasma vitamin A and retinol-binding protein in preterm and term neonates. *J Pediatr* 1981; 99: 302-5
9. Mactier H, Weaver LT. Vitamin A and preterm infants: what we know, what we don't know, and what we need to know. *Arch Dis Child Fetal Neonatal Ed.* 2005; 90:F103-8
10. Haas C, Genzel-Boroviczova O, Koletzko B. Losses of vitamin A and E in parenteral nutrition suitable for premature infants. *Eur J Clin Nutr* 2002; 56, 906-12
11. Massaro G, Massaro D. Postnatal treatment with retinoic acid increases the number of pulmonary alveoli in rats. *Am J Physiol* 1996; 270: 1305-10
12. Shenai JP, Kennedy KA, Chytil F, Stahlman MT. Clinical trial of vitamin A supplementation in infants susceptible to bronchopulmonary dysplasia. *J Pediatr* 1987; 111:269-77
13. Shenai JP, Rush MG, Parker RA, Chytil F. Sequential evaluation of plasma retinol-binding protein response to vitamin A administration in very-low-birth-weight neonates. *Biochem Mol Med.* 1995; 54: 67-74

- 14 Kennedy KA, Stole BJ, Ehrenkranz RA, et al. Vitamin A to prevent bronchopulmonary dysplasia in very low birth weight infants: Has the dose been too low? *Early Hum Dev* 1997; 49:19–31
15. Greene HL, Phillips BL, Frank L, Fillmore CM, Said HM, Murrell JE, et al. Persistently low blood retinol levels during and after parenteral feeding of very low birth weight infants : examination of losses into intravenous administration sets and a method of prevention by addition to a lipid emulsion. *Pediatrics* 1987; 79:894–900
16. Tyson JE, Wright LL, Oh W, Kennedy KA, Mele L, Ehrenkranz RA, et al. Vitamin A supplementation for extremely low birth weight infants. *N Eng J Med* 1999; 340: 1962–8
17. Darlow BA, Graham PJ. Vitamin A supplementation to prevent mortality and short and long–term morbidity in very low birth weight infants. *Cochrane database Systematic Reviews* 2007 Issue 4
18. Inder TE, Graham PJ, Winterbourn CC, Austin NC, Darlow BA. Plasma vitamin A levels in the very low birth weight infant: relationship to respiratory outcome. *Early Hum Dev* 1998; 52: 155–68

19. Shaul PW. Maternal vitamin A administration and the fetal ductus arteriosus.

Pediatr Res 2001; 49: 744-6

20. Ravishankar C, Nafday S, Green RS, Kamenir S, Lorber R, Stacewicz-

Sapuntzakis M, et al. A trial of vitamin A therapy to facilitate ductal closure in

premature infants. J Pediatr 2003; 143: 644-8



