

TABLE OF CONTENTS

	Page
ABSTRACT (IN THAI)	i
ABSTRACT (IN ENGLISH)	ii
ACKNOWLEDGEMENTS	iii
TABLE OF CONTENTS	iv
LIST OF TABLES	vi
LIST OF FIGURES	vii
LIST OF ABBREVIATIONS	ix
CHAPTER I INTRODUCTION	1
1. Rationale and Background	1
2. Objectives of the Study	4
3. Scope and Limitation of the Study	4
4. Anticipated Outcomes	4
CHAPTER II LITERATURE REVIEWS	5
1. Melatonin	5
2. Inflammation	12
3. Melatonin and anti-inflammatory studies	16
4. Anti-inflammatory models	19
CHAPTER III RESEARCH METHODOLOGY	26
1. Materials and reagents	26
2. Methods	26
CHAPTER IV RESULT AND DISCUSSION	31
1. Chemistry	31
2. Structure elucidation	33
3. Effect of melatonin and derivatives on Tail flick latency	35
4. Effect of melatonin and derivatives on acetic acid-induced writhing	39
5. Effect of melatonin and derivatives on croton oil induced ear edema	43

TABLE OF CONTENTS (Cont.)

	Page
6. Effect of melatonin and derivatives on cotton pellet induced granuloma	45
CHAPTER V CONCLUSIONS AND SUGGESTION	49
REFERENCES	51
APPENDICES	57
APPENDIX A RESEARCH ETHIC IN ANIMAL MODELS	58
APPENDIX B SPECTRUMS FOR STRUCTURE ELUCIDATION	60

LIST OF TABLES

	Page
Table 2.1 The expression of melatonin receptor and its function	10
Table 2.2 Limitation of <i>in vivo</i> studies anti-inflammatory models	21
Table 2.3 Limitation of <i>in vivo</i> studies analgesic models	23
Table 3.1 Summary grouping in animal models	28
Table 4.1 Condition for synthesis melatonin derivatives	32
Table 4.2 Summary ¹ H –NMR of melatonin, acetyl melatonin and benzoyl melatonin	34
Table 4.3 Summary ¹³ C –NMR of melatonin, acetyl melatonin and benzoyl melatonin	35
Table 4.4 Analgesic activity of the melatonin and derivatives in tail flick model with varies time s.c. after compound administration	37
Table 4.5 Summary results of analgesic activities of melatonin and derivatives in mice	42
Table 4.6 Summary results of croton oil induced ear edema of melatonin and derivatives in mice	45
Table 4.7 Summary result of chronic anti-inflammatory activities of melatonin and derivatives in mice	48

LIST OF FIGURES

		Page
Figure 1.1	Structure of indomethacin and melatonin derivatives	3
Figure 2.1	The structure of melatonin	6
Figure 2.2	The 24-hour cycle of melatonin production in human	6
Figure 2.3	The amount of melatonin secretion in human pineal gland in various ages	7
Figure 2.4	Biosynthesis pathway of melatonin	8
Figure 2.5	The docking of melatonin on MT ₁ and MT ₂ receptor	9
Figure 2.6	Size of substation via melatonin receptor binding studies	11
Figure 2.7	Activation of NF- κ b pathways	14
Figure 2.8	Biosynthesis pathway of PGE ₂	15
Figure 2.9	Structure of tryptophan, serotonin and melatonin	17
Figure 2.10	Melatonin derivatives in Elmegeed (2006) studies	18
Figure 2.11	Structure of indomethacin and derivatives	19
Figure 4.1	Reaction of acetyl melatonin synthesis	31
Figure 4.2	Reaction of benzoyl melatonin synthesis	32
Figure 4.3	Time-course effect of melatonin (MLT), acetyl melatonin (AcO) and benzoyl melatonin (Bn) derivatives on tail flick induced by warm water 50°C. Mice were treated by subcutaneous route with melatonin and its derivatives (50 mg/kg), Tramadol (50 mg/kg) or vehicle. The result was shown a duration before mice withdrawal or immersed tail as means \pm SE (n = 6 for each group).	38

LIST OF FIGURES (Cont.)

		Page
Figure 4.4	Effect of melatonin (MLT) and its acetyl (AcO), benzoyl (Bn) derivatives on tail flick latency time. Mice were treated by subcutaneous route with melatonin and its derivatives (10-50 mg/kg), Tramadol (50 mg/kg) or vehicle at 30 or 60 min prior to experiment. The result was shown duration before mice withdrawal or immersed tail, values are expressed as the means \pm SE, * = $P < 0.05$ when compared to vehicle treatment. (n = 6 for each group)	39
Figure 4.5	Effect of melatonin (MLT), acetyl melatonin (AcO), benzoyl melatonin (Bn) and aspirin (ASA) on acetic acid induced writhing. Mice were treated by subcutaneous route with melatonin and its derivatives (10-50 mg/kg), aspirin (300 mg/kg) or vehicle at 30 and 60 minutes before acetic acid injection. Values are expressed as the means \pm SE, * = $P < 0.05$ when compared to vehicle treatment	41
Figure 4.6	Effect of melatonin (MLT) and its acetyl (AcO), benzoyl (Bn) derivatives on croton oil induce ear edema. Melatonin and its derivatives (10-50 μ g/ear) or aspirin (600 μ g /ear) was applied to the ear 1 hr before croton oil. The result shown is relative %inhibition of ear thickness. Values are expressed as the means \pm SE, * = $P < 0.05$ when compared to vehicle treatment	44
Figure 4.7	Effect of melatonin (MLT) and its acetyl (AcO), benzoyl (Bn) derivatives and prednisolone (Pred) on granuloma tissue formation in mice. Mice were treated by subcutaneous route with melatonin and its derivatives (10-50 mg/kg), prednisolone (5 mg/kg) or vehicle daily for 7 days. Result shows as a granuloma weight. Values are expressed as the means \pm SE, * = $P < 0.05$ when compared to vehicle treatment.	47

