

## REFERENCES

- Aguilar, R., Ramírez, J.A., Garrote, G., & Vázquez, M. Kinetic study of the acid hydrolysis of sugar cane bagasse. *J. Food Eng.* 2002; 55: 309-318.
- Alani, F., & Smith, J.E. Effect of chemical pretreatment on the fermentation and ultimate digestibility of bagasse by *Phanerochaete chrysosporium*. *J. Sci. Food Agri.* 1988; 42: 19-28.
- Alexander, M.A., Chapman, T.W., & Jeffries, T.W. Xylose metabolism by *Candida shehatae* in continuous culture. *Appl. Microbiol. Biotechnol.* 1988; 28: 478-486.
- Aminoff, C., Vannan, E., & Doty, T.E. The occurrence manufacture and properties of xylitol, In counsell, J.N. (Ed). **Xylitol**. London: Applied Science Publishers; 1978.
- Amjed, M., Jury, H.G., & Donker, J.D. Effect of alkaline hydrogen peroxide treatment on cell wall Composition kinetic of sugarcane residues and wheat straw. *J. Animal Sci.* 1992; 70: 2877-2884.
- Amore, R., Kötter, P., Küster, C., Ciriacy, M., & Hollenberg, C.P. Cloning and expression in *Saccharomyces cerevisiae* of the NAD(P)H-dependent xylose reductase encoding gene (XYL1) from the xylose-assimilating yeast *Pichia stipitis*. *Gene.* 1991; 109: 89-97.
- Anderlund, M., Rådström, P., & Hahn-Hägerdal, B. Expression of biofunctional enzymes with xylose reductase and xylitol dehydrogenase activity in *Saccharomyces cerevisiae* alters product formation during xylose fermentation. *Metabolic Eng.* 2001; 3: 226-235.
- Azzam, A.M. Pre-treatment of agrocellulosic waste for microbial biomass production with a defined mixed culture. *J. Environ. Sci. Health Part B.* 1989; 27: 1643-1654.
- Bae, S.M., Park, Y.C., Lee, T.H., Kweon, D.H., Choi, J.H., Kim, S.K., Ryu, Y.W., & Seo, J.H. Production of xylitol by recombinant *Saccharomyces cerevisiae* containing xylose reductase gene in repeated fed-batch and cell recycle fermentations. *Enzyme Microb. Technol.* 2004; 35: 545-549.

- Becker, D.M., & Guarente, L. High-efficiency transformation of yeast by electroporation. In *Guide to Yeast Genetics and Molecular Biology*, pp.182-186. Edited by C. Guthrie & G. R. Fink: Academic Press. 1991.
- Barbosa, M.F.S., de Medeiros, M.B., de Mancilha, I.M., Schneider, H., & Lee, H. Screening of yeasts for production of xylitol from D-xylose and some factors which affect xylitol yield in *Candida guilliermondii*. *J. Indust. Microbiol.* 1988; 3: 241-251.
- Barnett, J.A., Payne, R.W., & Yarrow, D. **Yeast: characteristics and identification.** 2<sup>nd</sup> edition. Cambridge University, New York. Cambridge University Press. 1990.
- Bernd, N., Peter, M., Wilfried, N., & Michael, P. Structural and functional properties of aldose xylose reductase from the D-xylose-metabolizing yeast *Candida tenuis*. *Chem. Biol. Interact.* 2001; 130-132: 583-595.
- Billard, P., Ménart, S., Fler, R., & Bolotin-Fukuhara, M. Isolation and characterization of the gene encoding xylose reductase from *Kluyveromyces lactis*. *Gene.* 1995; 162: 93-97.
- Bolen, P., Hayman, G.T., & Shepherd, H.S. Sequence and analysis of an aldose (xylose) reductase gene from the xylose-fermenting yeast *Pachysolen tannophilus*. *Yeast.* 1996; 12: 1367-1375.
- Bravo, O., Ferrer, A., Aiello, C., Ledesma, A., & Davila, M. Growth of *Chaetomium cellulolyticum* in solid state fermentation of sugar cane bagasse treated with water and alkali at several liquid/solid ratios. *Biotechnol. Lett.* 1994; 16: 865-870.
- Chakravorty, M., Veiga, L.A., Bacila, M., & Horecker, B.L. Pentose metabolism in *Candida* II. The diphosphopyridine nucleotide-specific polyol dehydrogenase of *Candida utilis*. *J. Biol. Chem.* 1962; 237: 1014-1020.
- Chen, L.F., & Gong, C.S. Fermentation of sugarcane bagasse hydrolysate-acclimated yeast. *J. Food Science.* 1985; 50: 226-228.
- Chiang, C., & Knight, S.G. Metabolism of D-xylose by mould. *Nature.* 1960; 188: 79-81.
- Chin, C.S., Hwang, W.Z., & Lee, H.C. Isolation and characterization of mutants of *Clostridium saccharoperbutylacetonicum* fermenting bagasse hydrolysate. *J. Ferment. Bioeng.* 1991; 72: 249-253.

- Cruz, M.J., Domínguez J.M., Domínguez H., & Parajó, J.C. Xylitol production from barley bran hydrolysate by continuous fermentation with *Debaromyces hansenii*. **Biotechnol. Lett.** 2000; 22: 1895-1898.
- Dahiya, J.S. Xylitol production by *Peteromyces albertensis* grown on medium containing D-xylose. **Can. J. Microbiol.** 1991; 37: 14-18.
- Dahn, K.M., Davis, B.P., Pittman, P.E., Kenealy, W.R., & Jeffries, T.W. Increased xylose reductase activity in the xylose-fermenting yeast *Pichia stipitis* by overexpression of XYL1. **Appl. Biochem. Biotechnol.** 1996; 57-58: 267-276.
- Domínguez J.M., Gong C.S., & Tsao G.T. Pretreatment of sugar cane bagasse hemicellulose hydrolysate for xylitol production by yeast. **Appl. Biochem. Biotechnol.** 1996; 57/58: 49-56.
- Doran, J.B., Aldrich, H.C., & Ingram, L.O. Saccharification and fermentation of sugarcane bagasse. **Biotechnol. Bioeng.** 1994; 44: 240-247.
- du-Toit, P.J., Olivier, S.P., & van-Biljon, P.L. Sugarcane bagasse as a possible source of fermentable carbohydrates: I. Characterization of bagasse with regard to monosaccharide hemicellulose and amino acid composition. **Biotechnol. Bioeng.** 1984; 26: 1071-1078.
- Felipe, M.G.A., Vitalo, M., Mancilha, I.M., & Silva, S.S. Environmental parameters affecting xylitol production from sugarcane bagasse hemicellulose hydrolysate by *Candida guilliermondii*. **J. Indust. Microbiol. Biotechnol.** 1997; 18: 251-254.
- Fontana, J.D., Ramos, L.P., & Deschamps, F.C. Pretreated sugar cane bagasse as a model for cattle feeding. **Appl. Biochem. Biotechnol.** 1995; 51-52: 105-116.
- Gírio, F.M., Peito, A.M., & Amaral-Collaco, M.T. Enzymatic and physiological study of D-xylose metabolism by *Candida shehatae*. **Appl. Microbiol. Biotechnol.** 1989; 32: 199-204.
- Gírio, F.M., Pelica, F., & Amaral-Collaço, M.T. Characterization of xylitol dehydrogenase from *Debaromyces hansenii*. **Appl. Biochem. Biotechnol.** 1996; 56: 79-87.
- Gong, C.S., Chen, L.F., & Taso, G.T. Quantitative production of xylitol from D-xylose by a high-xylitol producing yeast mutant *Candida tropicalis* HXP2. **Biotechnol. Lett.** 1981; 3: 130-135.

- Gong, C.S., Claypool, T.A., McCracken, L.D., Maun, C.M., Ueng, P.P., & Isao, G.T. Conversion of pantoses by yeasts. *Biotechnol. Bioeng.* 1983; 25: 85-89.
- Guthrie, C., & Fink, G.K. Guide to yeast genetics and molecular biology. **In Methods in Enzymology**, Vol. 194. Edited by Abelson, J. N. and Simon, M. I. San Diego: Academic Press; CA. 1991.
- Hacker, B., Habenicht, A., Kiess, M., & Mattes, R. Xylose utilisation: cloning and characterisation of the xylose reductase from *Candida tenuis*. *Biol. Chem.* 1999; 380: 1395-1403.
- Hahn-Hägerdal, B., Jeppsson, H., Skoog, K., & Prior, B.A. Biochemistry and physiology of xylose fermentation by yeast. *Enzyme Microb. Technol.* 1994; 16: 933-944.
- Hallborn, J., Walfridsson, M., Airaksinen, U., Ojamo, H., Hahn-Hägerdal, B., Penttilä, M., & Keränen, S. Xylitol production by recombinant *Saccharomyces cerevisiae*. *Biotechnology.* 1991; 9: 1090-1095.
- Hallborn, J., Gorwa, M.F., Meinander, N., Penttilä, M., Keränen, S., & Hahn-Hägerdal, B. The influence of cosubstrate and aeration on xylitol formation by recombinant *Saccharomyces cerevisiae* expressing the XYL1 gene. *Appl. Microbiol. Biotechnol.* 1994; 42: 326-333.
- Handumrongkul, C., Ma, D.P., & Silva, J.L. Cloning and expression of *Candida guilliermondii* xylose reductase gene (xyl1) in *Pichia pastoris*. *Appl. Microbiol. Biotechnol.* 1998; 49: 399-404.
- Ho, N.W., Lin, F.P., Huang, S., Andrews, P.C., & Tsao, G.T. Purification, characterization, and amino terminal sequence of xylose reductase from *Candida shehatae*. *Enzyme Microb.* 1990; 12: 33-39.
- Horitsu, H., Yahasi, Y., Takamizawa, K., Kawai, K., Suzuki, T., & Watanabe, N. Production of xylitol from D-xylose by *Candida tropicalis*: optimization of production rate. *Biotechnol. Bioeng.* 1992; 40: 1085-1091.
- Hsiao, H.Y., Chiang, C.L., Ueng, P.P., & Tsao, G.T. Sequential utilization of mixed monosaccharides by yeasts. *Appl. Environ. Microbiol.* 1982; 43: 840-845.
- Izumori, K., & Tuzaki, K. Production of xylitol from D-xylulose by *Mycobacterium smegmatis*. *J. Ferment. Technol.* 1998; 66: 33-36.

- Jeffries, T.W. Utilization of xylose by bacteria, yeasts and fungi. *Adv. Biochem. Eng. Biotechnol.* 1983; 27: 1-32.
- Kang, M.H., Ni, H., & Jeffries, T.W. Molecular characterization of a gene for aldose reductase (CbXYL1) from *Candida boidinii* and its expression in *Saccharomyces cerevisiae*. *Appl. Biochem. Biotechnol.* 2003; 106: 265-276.
- Katzen, R., & Fowler, D.E. Ethanol from lignocellulosic wastes with utilisation of recombinant bacteria. *Appl. Biochem. Biotechnol.* 1994; 45-46: 697-707.
- Kilian, S.G., & van Uden, N. Transport of xylose and glucose in the xylose fermenting yeast *Pichia stipilis*. *Appl. Microbiol. Biotechnol.* 1988; 27: 545-548.
- Kim, S.B., & Lee, Y.Y. Diffusion of sulfuric acid within lignocellulosic biomass particles and its impact on dilute-acid pretreatment. *Bioresource Technology.* 2002; 83: 165-171.
- Kim S.Y., Kim J.H., & Oh D.K. Improvement of xylitol production by controlling oxygen supply in *Candida parapsilosis*. *J. Ferment. Bioeng.* 1997; 83: 267-270.
- Kim S.Y., Oh D.K., & Kim J.H. Evaluation of xylitol production from corn cob hemicellulose hydrolysate by *Candida parapsilosis*. *Biotechnol. Lett.* 1999; 21: 891-895.
- Kling, S.H., Neto, C.C., Ferrara, M.A., Torres, J.C.R., Magalhaes, D.B., & Ryu, D.D.Y. Enhancement of enzymatic hydrolysis of sugar cane bagasse by steam explosion pretreatment. *Biotechnol. Bioeng.* 1987; 29: 1035-1039.
- Kuhn, A., van Zyl, C., van Tonder, A., & Prior, B.A. Purification and partial characterization of an aldo-keto reductase from *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* 1995; 61: 1580-1585.
- Lache, A.H., & Jeffries, T.W. Levels of the enzyme of the pentose phosphate pathway in *Pachysolen tannophilus* Y-2460 and selected mutants. *Enzyme Microb. Technol.* 1986; 8: 353-359.
- Ladisich, M.R. Fermentable sugars from cellulosic residues. *Process Biochemistry.* 1979; 1: 21-25.
- Lavarack, B.P., Griffin, G.J., & Rodman, D. Measured kinetics of the acid-catalysed hydrolysis of sugarcane bagasse to produce xylose. *Catalysis today.* 2000; 63: 257-265.

- Lee, J.K., Koo, B.S., & Kim, S.Y. Cloning and characterization of the xyl1 gene, encoding an NADH-prefering xylose reductase from *Candida parapsilosis*, and its functional expression in *Candida tropicalis*. *Appl. Environ. Microbiol.* 2003; 69: 6179-6188.
- Ligthelm, M.E., Prior, B.A., du Preez, J C., & Brandt, V. An investigation of D-(1-<sup>13</sup>C)xylose metabolism in *Pichia stipilis* under aerobic and anaerobic conditions. *Appl. Microbiol. Biotechnol.* 1988; 28: 293-296.
- Lu J., Tsai L.B., Gong C.S., & Tsao G.T. Effect of nitrogen sources on xylitol production from D-xylose by *Candida* sp. L-102. *Biotechnol. Lett.* 1995; 17: 167-170.
- Magee, R.J., & Kosaric, N. Bioconversion of hemicellulose. *Adv. Biochem. Eng. Biotechnol.* 1985; 32: 61-93.
- Mäkinen, K.K. Dietary prevention of dental caries by xylitol-clinical effectiveness and safety. *J. Appl. Nutr.* 1992; 44: 16-28.
- Mäkinen, K.K. Can the pentitol-hexitol theory explain the clinical observation made with xylitol. *Med. Hypotheses.* 2000; 54: 603-613.
- Maleszka, R., & Schneider, H. Fermentation of D-xylose, xylitol and D-xylulose by yeasts. *Can. J. Microbiol.* 1982; 28: 360-365.
- Martin, C., Galbe, M., Wahlbom, C.F., Hahn-Hägerdal, B., & Jönsson, L.J. Ethanol production from enzyme hydrolysates of sugarcane bagasse using recombinant xylose-utilising *Saccharomyces cerevisiae*. *Enzyme Microb. Technol.* 2002; 31: 274-282.
- Martínez, E.A., Silva, S.S., & Felipe, M.G.A. Effect of the oxygen transfer coefficient on xylitol production from sugarcane bagasse hydrolysate by continuous stirred-tank reactor fermentation. *Appl. Biochem. Biotechnol.* 2000; 84-86: 633-641.
- Mayerhoff, Z.D.V.L., Roberto, I.C., & Silva, S.S. Xylitol production from rice straw hemicellulose hydrolysate using different yeast strains. *Biotechnol. Lett.* 1997; 19: 407-409.
- Mayr, P., Bruggler, K., Kulbe, K.D., & Nidetzky, B. D-Xylose metabolism by *Candida intermedia*: isolation and characterisation of two forms of aldose reductase with different coenzyme specificities. *J. Chromatogr.* 2000; 737: 195-202.

- Meinander, N., Hahn-Hägerdal, B., Linko, M., Linko, P., & Ojamo, H. Fed-batch xylitol production with recombinant XYL-1-expression *Saccharomyces cerevisiae* using ethanol as a co-substrate. *Appl. Microbiol. Biotechnol.* 1994; 42: 334-339.
- Meyer, P.S., du Preez, J.C., & Kilian, S.G. Chemostat cultivation of *Candida blankii* on sugarcane bagasse hemicellulose hydrolysate. *Biotechnol. Bioeng.* 1992; 40: 353-358.
- Meyrial, V., Delganes, J.P., Moletta, R., & Navarro, J.M. Xylitol production from D-xylose by *Candida guilliermondii*. *Biotechnol. Lett.* 1991; 13: 281-286.
- Mussatto, S.I., & Roberto, I.C. Hydrolysate detoxification with activated charcoal for xylitol production by *Candida guilliermondii*. *Biotechnol. Lett.* 2001; 23: 1681-1684.
- Mussatto, S.I., & Roberto, I.C. Alternatives for detoxification of diluted-acid lignocellulosic hydrolyzates for use in fermentative process. *Bioresource Technology.* 2004; 93: 1-10.
- Neuhauser, W., Haltrich, D., Kulbe, K.D., & Nidetzky, B. NAD(P)H-dependent aldose reductase from the xylose-assimilating yeast *Candida tenuis*, Isolation, characterization and biochemical properties of the enzyme. *Biochem. J.* 1997; 326: 683-692.
- Nidetzky, B., Bruggler, K., Kratzer, R., & Mayr, P. Multiple forms of xylose reductase of *Candida intermedia*: comparison of their functional properties using quantitative structure-activity relationships, steady-state kinetic analysis and pH studies. *J. Agric. Food. Chem.* 2003; 51: 7930-7935.
- Nigam, P., & Singh, D. Process for fermentative production of xylitol—a sugar substitute. *Process Biochemistry.* 1995; 30: 117-124.
- Nishio, N., Sugawa, K., Hayase, N., & Nagai, S. Conversion of D-xylose into xylitol by immobilized cell of *Candida pelliculosa* and *Methanobacterium* Sp HU. *J. Ferment. Bioeng.* 1989; 67: 356-360.
- Nolleau, V., Preziosi-Belloy, L., Delgenes, J.P., & Navarro, J.M. Xylitol production from xylose by two yeast strains: sugar tolerance. *Current Microbiology.* 1993; 27: 191-197.

- Olsson, L., & Hahn-Hägerdal, B. Fermentation of lignocellulosic hydrolysates for ethanol production. *Enzyme Microb. Technol.* 1996; 18: 312-331.
- Palmqvist, E., & Hahn-Hägerdal, B. Fermentation of lignocellulosic hydrolysates I: inhibition and detoxification. *Bioresour. Technol.* 2000; 74: 17-24.
- Pandey, A., Soccol, C.R., Nigam, P., & Soccol, V.T. Biotechnological potential of agro-industrial residues. I. sugarcane bagasse. *Bioresour. Technol.* 2000; 74: 69-80.
- Parajó, J.C., Domínguez, H., & Domínguez, J.M. Production of xylitol from concentrated wood hydrolysates by *Debaromyces hansenii*: effect of the initial cell concentration. *Biotechnol. Lett.* 1996; 18: 593-598.
- Parajó, J.C., Domínguez H., & Domínguez J.M. Improved xylitol production with *Debaromyces hansenii* Y-7426 from raw or detoxified wood hydrolysates. *Enzyme Microb. Technol.* 1997; 21: 18-24.
- Pessoa, A., Mancilha, I.M., & Sato, S. Cultivation of *Candida tropicalis* in sugarcane hemicellulose hydrolysate for microbial protein production. *J. Biotechnol.* 1996; 51: 83-88.
- Pfeifer M.J., Silva S.S., Felipe M.G.A., Roberto I.C., & Mancilha I.M. Effect of culture conditions on xylitol production by *Candida guilliermondii* FTI 20037. *Appl. Biochem. Biotechnol.* 1996; 57/58: 423-430.
- Preez du J.C. Process parameters and environmental factors affecting D-xylose fermentation by yeasts. *Enzyme Microb. Technol.* 1994; 16: 945-956.
- Prior, B.A., Killian, S.G., & du Preez, J.C. Fermentation of D-xylose by the yeasts *Candida shehatae* and *Pichia stipitis*. *Process Biochemistry.* 1989; 24: 21-32.
- Purchase, B.S. Products from sugarcane. *International Sugar Journal.* 1995; 97: 70.
- Roberto, I.C., Felipe, M.G.A., Lacis, L.S., Silva, S.S., & Demancilha, I.M. Utilization of sugarcane bagasse hemicellulose hydrolysate by *Candida guilliermondii* for xylitol production. *Bioresour. Technol.* 1991a; 36: 271-275.
- Roberto, I.C., Lacis, L.S., Barbosa, M.F.S., & Demancilha, I.M. Utilization of sugarcane bagasse hemicellulose hydrolysate by *Pichia stipitis* for the production of ethanol. *Process Biochemistry.* 1991b; 26: 15-21.

- Roberto, I.C., Mancilha, I.M., de Souza, C.A., Felipe, M.G.A., Sato, S., & de Castro, H.F. Evaluation of rice straw hemicellulose hydrolysate in the production of xylitol by *Candida guilliermondii*. ***Biotechnol. Lett.*** 1994; 16: 1211-1216.
- Roberto, I.C., Sato, S., & de Mancilha, I.M. Effect of inoculum level on xylitol production from rice straw hemicellulose hydrolysate by *Candida guilliermondii*. ***J. Indust. Microbiol.*** 1996; 16: 348-350.
- Roca, E., Meinander, N., & Hahn-Hägerdal, B. Xylitol production by immobilized recombinant *Saccharomyces cerevisiae* in a continuous packed-bed bioreactor. ***Biotechnol. Bioeng.*** 1996; 51: 317-326.
- Rodríguez-Chong, A., Ramírez, J.A., Garrote, G., & Vázquez, M. Hydrolysis of sugar cane bagasse using nitric acid: a kinetic assessment. ***J. Food Eng.*** 2004; 61: 143-152.
- Roseiro, J.C., Peito, F.M., & Amaral-Callaco, M. The effects of oxygen transfer coefficient and substrate concentration on xylose fermentation by *Debaromyces*. ***Arch Microbiol.*** 1991; 156: 484-450.
- Saha, B.C., & Bothast, R.J. Production of xylitol by *Candida peltata*. ***J. Indust. Microbiol. Biotechnol.*** 1999; 22: 633-636.
- Sambrook, J., Fritsch, E.F., & Maniatis, T. **Molecular cloning: A Laboratory Manual.** 2<sup>nd</sup> edition. New York: Cold Spring Harbor laboratory Press; 1989.
- Santos, J.C., Converti, A., de Carvalho, W., Mussatto, S.I., & Silva, S.S. Influence of aeration rate and carrier concentration on xylitol production from sugarcane bagasse hydrolyzate in immobilized-cell fluidized bed reactor. ***Process Biochemistry.*** 2005; 40: 113-118.
- Schmitt, M.E., Brown, T.A., & Trumppower, B.L. A rapid and simple method for preparation of RNA from *Saccharomyces cerevisiae*. ***Nucleic Acids Res.*** 1990; 18: 3091.
- Schneider, H., Lee, H., Barbosa, M.F.S., Kubicek, C.P., & James, A.P. Physiological properties of a mutant *Pachysolen tannophilus* deficient in NADHP-dependent D-xylose reductase. ***Appl. Environ. Microbiol.*** 1989; 55: 2877-2881.
- Schneider, H. Conversion of pentoses to ethanol by yeasts and fungi. ***CRC. Crit.Rev. Biotechnol.*** 1989; 9: 1-40.

- Sene, L., Felipa, M.G.A., Vitolo, M., Silva, S.S., & Mancilha, I.M. Adaptation and neutilization of *Candida guilliermondii* cells for xylitol production in bagasse hydrolysate. *J. Basic. Microbiol.* 1998; 38: 61-69.
- Sene, L., Felipa, M.G.A., Silva, S.S., & Vitolo, M. Preliminary kinetic characterization of xylose reductase and xylitol dehydrogenase extracted from *Candida guilliermondii* FTI 20037 cultivated in sugarcane bagasse hydrolysate for xylitol production. *Appl. Biochem. Biotechnol.* 2001; 91-93: 671-680.
- Silva, S.S., Roberto, I.C., Felipa, M.G.A., & Mancilha, I.M. Batch fermentation of xylose for xylitol production in stirred tank bioreactor. *Process Biochemistry.* 1996; 31: 549-553.
- Silva, S.S., Ribeiro, J.D., Felipa, M.G.A., & Vitolo, M. Maximizing xylitol production from sugarcane bagasse hydrolysate by controlling the aeration rate. *Appl. Biochem. Biotechnol.* 1997; 63-65: 557-564.
- Silva, S.S., Felipa, M.G.A., & Mancilha, I.M. Factors that affect the biosynthesis of xylitol by xylose-fermenting yeasts. *Appl. Biochem. Biotechnol.* 1998; 70-72: 331-339.
- Sirisansaneeyakul, S., Staniszewski, M., & Rizzi, M. Screening of yeasts for production of xylitol from D-xylose. *J. Ferment. Bioeng.* 1995; 6: 564-570.
- Skoog, K., & Hahn-Hägerdal, B. Xylose fermentation. *Enzyme Microb. Technol.* 1988; 10: 66-80.
- Slininger, P.J., Bolen, P.L., & Kurtzman, C.P. *Pachysolen tannophilus*: properties and process consideration for ethanol production from D-xylose. *Enzyme Microb. Technol.* 1987; 16: 933-943.
- Smiley, K.L., & Bolen, P.L. Demonstration of D-xylose reductase and D-xylitol dehydrogenase in *Pachysolen tannophilus*. *Biotechnol. Lett.* 1982; 4: 607-610.
- Takuma, S., Nakashima, N., Tangtirunkij, M., Kinoshita, S., Okada, H., Seki, T., & Yoshida, T. Isolation of xylose reductase gene of *Pichia stipitis* and its expression in *Saccharomyces cerevisiae*. *Appl. Biochem. Biotechnol.* 1991; 28: 327-340.
- Turner, P.C., McLennan, A.G., Bates, A.D., & White, M.R.H. **Insant Notes Molecular Biology**. 2<sup>nd</sup> edition. United Kingdom: Guildford, Biddles Ltd; 2001.

- van Miltenburg, R., Rüger, B., Grünewald-Janho, S., Leons, M., & Schröder, C. **The DIG system user's guide for filter hybridization.** 1<sup>st</sup> edition. Germany: Boehringer Mannheim GmbH, Biochemica; 1995.
- Vandeska, E., Amartey, S., Kuzmanova, S., & Jeffries, T.W. Effect of environmental conditions on production of xylitol by *Candida boidinii*. *J. Microbiol. Biotechnol.* 1995a; 11: 213-218.
- Vandeska, E., Kuzmanova, S., & Jeffries, T.W. Xylitol formation and key enzyme activities in *Candida boidinii* under different oxygen transfer rates. *J. ferment. Bioeng.* 1995b; 80: 513-516.
- Vandeska, E., Amartey, S., Kuzmanovat, S., & Jeffries, T.W. Fed-batch culture for xylitol production by *Candida boidinii*. *Process Biochemistry.* 1996; 31: 265-270.
- Walfridsson, M., Anderlund, M., Bao, X., & Hahn-Hägerdal, B. Expression of different levels of enzymes from *Pichia stipitis* XYL1 and XYL2 genes in *Saccharomyces cerevisiae* and its effects on product formation during xylose utilization. *Appl. Microbiol. Biotechnol.* 1997; 48: 218-224.
- Winkelhausen, E., & Kuzmanova, S. Microbial Conversion of D-Xylose to Xylitol. *J. Ferment. Bioeng.* 1998; 86: 1-14.
- Woodyer, R., Simurdiak, M., van der Donk, W.A., & Zhao, H. Heterologous expression, purification, and characterization of a highly active xylose reductase from *Neurospora crassa*. *Appl. Environ. Microbiol.* 2005; 71: 1642-1647.
- Yahashi, Y., Horitsu, H., Kawai, K., Suzuki, T., & Takamizawa, K. Production of xylitol from D-xylose by *Candida tropicalis*: the effect of D-xylose feeding. *J. Ferment. Bioeng.* 1996; 81: 148-152.
- Yokoyama, S.I., Kinoshita, Y., Suzuki, T., Kawai, K., Horitsu, H., & Takamizawa, K. Cloning and sequencing of two D-xylose reductase genes (*xyrA* and *xyrB*) from *Candida Tropicalis*. *J. Ferment. Bioeng.* 1995a; 80: 603-605.
- Yokoyama, S.I., Suzuki, T., Kawai, K., Horitsu, H., & Takamizawa, K. Purification, characterization and structure analysis of NAD(P)H-dependent D-xylose reductase from *Candida tropicalis*. *J. Ferment. Bioeng.* 1995b; 79: 217-223.