

## TABLES OF CONTENTS

	<b>Page</b>
ABSTRACT (IN THAI)	i
ABSTRACT (IN ENGLISH)	iii
ACKNOWLEDEMENTS	v
LIST OF TABLES	viii
LIST OF FIGURES	ix
CHAPTER 1 INTRODUCTION	1
1.1 Introduction	1
1.2 Objectives of the research	3
CHAPTER 2 LITERTURE REVIEW	4
2.1 Bagasse and Pretreatment	4
2.2 Acid hydrolysis processes	5
2.3 Xylitol and conventional production of xylitol	9
2.4 Xylitol-producing microorganisms	10
2.5 Conversion of D–xylose to xylitol by yeasts	12
2.6 Process conditions affecting biological production of xylitol	14
2.7 DNA cloning / recombinant DNA	16
2.8 <i>Pichia pastoris</i> expression system	16
2.9 Xylose reductase	19
CHAPTER 3 MATERIALS AND METHODS	20
3.1 Materials	20
3.2 Analytical Methods	22
3.3 Methods	23
CHAPTER 4 RESULTS AND DISCUSSION	39
4.1 Bagasse pretreatment by acid hydrolysis	39
4.2 Genetic manipulation of yeasts from selected xylitol producing yeasts	44
4.3 Growth and xylitol productivity of the isolates and recombinant yeasts	56

**TABLES OF CONTENTS (Cont.)**

	<b>Page</b>
4.4 Batch culture by recombinant <i>K. marxianus</i> (rKm6) for xylitol production	62
CHAPTER 5 CONCLUSIONS	69
REFERENCES	72
APPENDICES	83
Appendix A Chemicals and reagents	84
Appendix B Plasmid vectors	93
Appendix C Analysis of variance	101
Appendix D Chromatogram standard	103
Appendix E Primer design	107
Appendix F Research publication	114

## LIST OF TABLES

	<b>Page</b>
Table 1 Xylitol production by yeasts from hemicellulose hydrolysate	6
Table 2 HCl acid processes	7
Table 3 H <sub>2</sub> SO <sub>4</sub> acid processes	8
Table 4 Screening of yeasts for xylitol production from D-xylose	11
Table 5 Enzymes used in DNA cloning	18
Table 6 Reducing sugar concentrations (as xylose; g/l) obtained in the 1% (w/v) H <sub>2</sub> SO <sub>4</sub> hydrolysis of sugarcane bagasse at 126.7°C (1.5 kg/cm <sup>2</sup> pressure gauge) in the range of 15 - 60 min reaction time	40
Table 7 Reducing sugar concentrations (as xylose; g/l) obtained in the 2% (w/v) H <sub>2</sub> SO <sub>4</sub> hydrolysis of sugarcane bagasse at 126.7°C (1.5 kg/cm <sup>2</sup> pressure gauge) in the range of 15 - 60 min reaction time	41
Table 8 Reducing sugar concentrations (as xylose; g/l) obtained in the 3% (w/v) H <sub>2</sub> SO <sub>4</sub> hydrolysis of sugarcane bagasse at 126.7°C (1.5 kg/cm <sup>2</sup> pressure gauge) in the range of 15 - 60 min reaction time	42
Table 9 Xylitol production from bagasse hydrolysate by different yeasts during batch fermentation	62
Table 10 Xylitol production during batch fermentation in sugarcane bagasse hydrolysate by recombinant <i>K. marxianus</i> (rKm6) under three different aeration rate of 0.5, 1.0 and 1.5 vvm and stirring rate of 200 and 300 rpm at 30°C for approximately 168 h	68

## LIST OF FIGURES

	<b>Page</b>
Figure 1 Metabolism of xylose via a xylose reductase and a xylitol dehydrogenase	14
Figure 2 Reducing sugar concentrations (as xylose; g/l) obtained in 1-3%(w/v) H <sub>2</sub> SO <sub>4</sub> hydrolysis of sugarcane bagasse at 126.7°C (1.5 kg/cm <sup>2</sup> pressure gauge) in the range of 15, 30, 45 and 60 min reaction times	43
Figure 3 Growth of <i>C. guilliermondii</i> 5068 on different carbon sources media.	44
Figure 4 Growth of <i>H. anomala</i> 5302 on different carbon sources media.	45
Figure 5 Growth of <i>K. marxianus</i> 5057 on different carbon sources media.	45
Figure 6 Genomic DNA extraction from <i>C. guilliermondii</i> , <i>H. anomala</i> and <i>K. marxianus</i> .	46
Figure 7 PCR production of xylose reductase (XR) gene from <i>C. guilliermondii</i> .	46
Figure 8 PCR production of xylose reductase (XR) gene from <i>K. marxianus</i> 5057	47
Figure 9 DNA Dot blotting analysis showing xylose reductase (XR) gene in <i>K. marxianus</i> 5057	48
Figure 10 Northern - blot analysis showing xylose reductase (XR) gene in <i>K. marxianus</i> 5057 grown on various carbon sources and period of time; (YPDX Glucose-Xylose medium), YPX (Xylose medium) and sugarcane bagasse hydrolysate medium and for 2, 3, 4 and 5 days, respectively	49
Figure 11 Southern – blot analysis showing the 4.5 kb <i>SacI</i> fragment	50
Figure 12 Positive colonies hybridisation with blue colour	51
Figure 13 Sequences of <i>K. marxianus</i> 5057 XR gene (colony 1)	51
Figure 14 Sequences of <i>K. marxianus</i> 5057 XR gene (colony 2)	52
Figure 15 Sequences blast of <i>K. marxianus</i> 5057 XR gene (colony 1) with <i>K. lactis</i> xylose reductase (XYL1) gene from database GenBank	53

## LIST OF FIGURES (Cont.)

	<b>Page</b>
Figure 16 Sequences blast of <i>K. marxianus</i> 5057 XR gene (colony 2) with <i>K. lactis</i> xylose reductase (XYL1) gene from database GenBank	54
Figure 17 SDS-PAGE showing XR protein and the molecular mass in <i>P. pastoris</i> GS115	55
Figure 18 Amino acid sequences of XR gene of <i>K. marxianus</i> in <i>P. pastoris</i> GS115	56
Figure 19 Glucose utilisation (g/l) during batch fermentation in sugarcane bagasse hydrolysate by different yeasts	58
Figure 20 Xylose consumption (g/l) during batch fermentation in sugarcane bagasse hydrolysate by different yeasts	59
Figure 21 Cell mass (as dry weight; g/l) during batch fermentation in sugarcane bagasse hydrolysed bagasse by different yeasts	59
Figure 22 Xylitol production (g/l) during batch fermentation in sugarcane bagasse hydrolysate by different yeasts	60
Figure 23 Arabinose concentrations (g/l) during batch fermentation in sugarcane bagasse hydrolysate by different yeasts	60
Figure 24 Acetic acid concentrations (g/l) during batch fermentation in sugarcane bagasse hydrolysate by different yeasts	61
Figure 25 Ethanol concentrations (g/l) during batch fermentation in sugarcane bagasse hydrolysate by different yeasts	61
Figure 26 Cell mass (as dry weight; g/l) during batch fermentation in sugarcane bagasse hydrolysate by recombinant <i>K. marxianus</i> (rKm6) under different aeration rate of 0.5, 1.0 and 1.5 vvm and stirring rate of 200 rpm at 30°C for approximately 168 h	64
Figure 27 Composition of sugarcane bagasse hydrolysate during batch fermentation by recombinant <i>K. marxianus</i> (rKm6) under aeration rate of 0.5 vvm and stirring rate of 200 rpm at 30°C for approximately 168 h	65

**LIST OF FIGURES (Cont.)**

	<b>Page</b>
Figure 28 Composition of sugarcane bagasse hydrolysate during batch fermentation by recombinant <i>K. marxianus</i> (rKm6) under aeration rate of 1.0 vvm and stirring rate of 200 rpm at 30°C for approximately 168 h	65
Figure 29 Composition of sugarcane bagasse hydrolysate during batch fermentation by recombinant <i>K. marxianus</i> (rKm6) under aeration rate of 1.5 vvm and stirring rate of 200 rpm at 30°C for approximately 168 h	66
Figure 30 Xylitol production (g/l) during batch fermentation in sugarcane bagasse hydrolysate by recombinant <i>K. marxianus</i> (rKm6) under three different aeration rate of 0.5, 1.0 and 1.5 vvm and stirring rate of 200 rpm at 30°C for approximately 168 h	66
Figure 31 Composition of sugarcane bagasse hydrolysate during batch fermentation by recombinant <i>K. marxianus</i> (rKm6) under aeration rate of 0.5 vvm and stirring rate of 300 rpm at 30°C for approximately 168 h	67
Figure 32 Xylitol production (g/l) during batch fermentation in sugarcane bagasse hydrolysate by recombinant different yeasts under different aeration rate of 0.5, 1.0 and 1.5 vvm and stirring rate of 200 rpm at 30°C for approximately 168 h	67