

เอกสารอ้างอิง

- ธีระ เอกสมทราเมษฐ์, ชัยรัตน์ นิลนนท์, ชีรพงศ์ จันทรมนิยม, ประกิจ ทองคำ และสมเกียรติ สีสนอง. 2548. เส้นทางสู่ความสำเร็จการผลิตปาล์มน้ำมัน. พิมพ์ครั้งที่ 2. ศูนย์วิจัยและพัฒนาการผลิตปาล์มน้ำมันคณะทรัพยากรธรรมชาติ มหาวิทยาลัยสงขลานครินทร์, สงขลา
- ศูนย์วิจัยปาล์มน้ำมันสุราษฎร์ธานี. 2532. ปาล์มน้ำมันพันธุ์ลูกผสมสุราษฎร์ธานี. โครงการวิจัยและพัฒนาพันธุ์ปาล์มน้ำมัน สถาบันวิจัยพืชสวน กรมวิชาการเกษตร.
- Altunkaya, A. and Gokmen, V. 2009. Effect of various anti-browning agents on phenolic compounds profile of fresh lettuce (*L. sativa*). Food Chem. Doi: 10.1016/j.foodchem.2009.03.085.
- Barcelos, E., Amblard, P., Berthaud, J., and Seguin, M. 2002. Genetic diversity and relationship in American and African oil palm as revealed by RFLP and AFLP molecular marker. Pesq Agropec Bras. 37(8): 1105-1114.
- Bennett, MD and Leitch, I. 2005. Genome size evolution in plants. In: Gregory T, ed. The evolution of the genome. London: Elsevier Academic Press, 89–162.
- Dolezel, J., Binarova, P. and Lucretti, S. 1989. Analysis of nuclear DNA content in plant cell by flow cytometry. Biol. Plantarum (Praha). 31(2): 113-120.
- Dolezel, J. 1991. Flow cytometric analysis of nuclear DNA content in higher plants. Phytochem. Analysis. 2: 143-154.
- Dolezel, J., Bartos. J., Voglmayr, H. and Greilhuber, J. 2003. Nuclear DNA content and genome size of trout and human. Cytometry Part A. 51A: 127-128.
- Dolezel, J. and Bartos, J. 2005. Plant DNA flow cytometry and estimation of nuclear genome size. Ann. Bot. 95: 99-110.
- Fang, D.Q. and Roose, M.L. 1997. Identification of closely related citrus cultivars with intersimple sequence repeat markers. Theor Appl Genet. 95: 408-417.
- Galbraith, D. W., Harkins, K. R., Maddox, J. M., Ayres, N. M., Sharma, D. P. and Firoozabady, E. 1983. Rapid flow cytometric analysis of the cell cycle in intact plant tissues. Science. 220: 1049-1051.
- Greihuber, J., Tensch, E. M. and Loureiro, J. 2007 Nuclear DNA content measurement. In: *Flow cytometry with plant cells*. (Eds) Dolezel, J., Greihuber, J. and Suda J. WILEY-VSH Verlag GmbH&Co. KGaA. Weinheim, Germany. pp. 67-101.
- Gupta, M., Chyi, Y-S., Romeo-Severson, J. and Owen, J.L. 1994. Amplification of DNA markers from evolutionarily diverse genomes using single primers of simple-sequence repeats. Theor Appl Genet. 89: 998-1006.

- Loureiro, J., Rodriguez, E., Dolezel, J. and Santos, C. 2006a. Flow cytometric and microscopic analysis of the effect of tannic acid on plant nuclei and estimation of DNA content. *Ann. Bot.* 98: 515-527.
- Loureiro, J., Rodriguez, E., Dolezel, J. and Santos, C. 2006b. Comparison of four nuclear isolation buffers for plant DNA flow cytometry. *Ann. Bot.* 98: 679-689.
- Loureiro, J., Rodriguez, E., Dolezel, J. and Santos, C. 2007. Two new nuclear isolation buffers for plant DNA flow cytometry: A test with 37 species. *Ann. Bot.* 1-14. doi:10.1093/annbot/mcm152.
- Madon, M., Phoon, L. Q., Clyde, M. M. and Mohd, D. 2008. Application of flow cytometry for estimation of nuclear DNA content in *Elaeis*. *J Oil Palm Res.* 20: 447-452.
- Mayes, S., Jack, P. L. and Corley, R. H. V. 2000. The use of molecular markers to investigate the genetic structure of an oil palm breeding programme. *Heredity.* 85(3): 288-293.
- Otto, E. J. 1990. DAPI staining of fixed cells for high-resolution flow cytometry of nuclear DNA. In: *Methods in cell biology.* (Eds) Darzynkiewicz, Z. and Crissman, H. A. Vol.33. San Diego: Academic Press. pp. 105-110.
- Pfossen, A., Amon, A., Lelley, T. and Heberle-bors, E. 1995. Evaluation of sensitivity of flow cytometry in detecting aneuploidy in wheat using disomic and ditelosomic wheat-rye addition lines. *Cytometry.* 21: 387-393.
- Rival, A., Beule, T., Barre, P., Hamon, S., Duval, Y. and Noirot, M. 1997. Comparative flow cytometric estimation of nuclear DNA content in oil palm (*Elaeis guineensis* Jacq.) tissue cultures and seed-derived plants. *Plant Cell Rep.* 16: 884-887.
- Sathish, D. K. and Mohankumar, C. 2007. RAPD markers for identifying oil palm (*Elaeis guineensis* Jacq.) parental varieties (*dura* & *pisifera*) and the hybrid *Tenera*. *Indian J Biotechnol.* 6: 354-358.
- Srisawat, T., Kanchanapoom, K., Pattanapanyasat, K., Srikul, S. and Chuthammathat, W. 2005. Flow cytometric analysis of oil palm: a preliminary analysis for cultivars and genomic DNA alteration. *Songklanakarin J Sci. Technol.* 27(Suppl. 3): 645-652.
- Srisawat, T. and Kanchanapoom, K. 2005. The influence of physical conditions on embryo and protoplast culture in oil palm (*Elaeis guineensis* Jacq.). *ScienceAsia.* 31: 23-28.
- Te-Chato, S. and Thawaro, S. 2008. RAPD (random amplified polymorphic DNA) marker as a tool for hybrid oil palm verification from half mature zygotic embryo culture. *J Agri. Technol.* 2: 165-176.

Winkelmann, T., Sangwan, R. S. and Schwenke, H. G. 1998. Flow cytometric analyses in embryogenic and non-embryogenic callus lines of *Cyclamen persicum* Mill.: relation between ploidy level and competence for somatic embryogenesis. *Plant Cell Rep.* 17: 400-404.

Output ที่ได้รับจากโครงการวิจัย