


เอกสารอ้างอิง

- Chantrapromma, S., Jindawong, B., Fun, H.-K., Patil, P.-S. and Karalai, C. (2006) "2-[(E)-2-(3-Hydroxy-4-methoxyphenyl)ethenyl]-1-methylquinolinium 4-chloro" *Acta Cryst.*, E62: o1802-o1804.
- Crasta, V., Ravindrachary, V., Bhajantri, R. F. and Gonsalves, R. (2004) "Growth and characterization of an organic NLO crystal: 1-(4-methylphenyl)-3-(4-methoxyphenyl)-2-propen-1-one" *J. Cryst. Growth* 267: 129-133.
- Hong, H.-K., Park, J. W., Lee, K.-S. and Yoon, C.-S. (2005) "Growth of highly nonlinear optical organic crystal, 3-methyl-4-methoxy-4'-nitrostilbene (MMONS)" *J. Cryst. Growth* 277: 509-517.
- Jindawong, B., Chantrapromma, S., Fun, H.-K., Yu, X.-L. and Karalai, C. (2005) "4(4'-Hydroxy-3'-methoxystyryl)-1-Methylpyridinium 4-bromobenzenesulfonate", *Acta Cryst.*, E61, o1340-o1342.
- Lakshmanaperumal, C. K., Arulchakkaravarthi, A., Balamurugan, N., Santhanaraghavan, P. and Ramasamy, P. (2004) "Synthesis, Crystal growth and Characterization of novel NLO material: 4-hydroxy benzaldehyde-N-methyl-4-stilbazolium tosylate", *Journal of Crystal Growth* 265: 260-265.
- Lakshmanaperumal, C. K., Arulchakkaravarthi, A., Rajesh, N. P., Raghavan, P.S., Huang, Y. C., Ichimura, M. and Ramasamy, P. (2002) "Synthesis, Crystal growth and FTIR, NMR, SHG studies of 4-methoxy benzaldehyde-N-methyl-4-stilbazolium tosylate (MBST)", *Journal of Crystal Growth* 240: 212-217.
- Nogi, K., Usman, A., Tsuji, K., Duan, X.-M., Okada, S., Oikawa, H., Matsuda, H. and Nakanishi, H. (2000) "Preparation of Polyene Analogues of Stilbazolium", *Nonlinear Optics* 24: 35-40.

Umezawa, H., Tsuji, K., Usman, A., Duan, X.-M., Okada, S., Oikawa, H., Matsuda, H. and Nakanishi, H. (2000) "Synthesis of Stilbazolium Derivatives having 2-(6-Dimethylamino)Naphthyl Group for Nonlinear Optics", *Nonlinear Optics* 24: 73-78.

Usman, A., Duan, X.-M., Komatsu, K., Okada, S., Matsuda, H., Oikawa, H. and Nakanishi, H. (1997) "Second-Order Hyperpolarizability of Pyridinium Cations" *Chemistry Letters*. 247-248.



Usman, A., Duan, X.-M., Komatsu, K., Okada, S., Oikawa, H., Matsuda, H. and Nakanishi, H. (1999) "Synthesis of Substituted pyridinium-benzenesulfonates crystals for second-harmonic generation", *Nonlinear Optics*, 22: 251-254.

Usman, A., Duan, X.-M., Okada, S., Matsuda, H., Oikawa, H. and Nakanishi, H. (1998) "First Hyperpolarizability of benzoate anions and esters", *J. Chem. Soc. Perkin. Trans 2*: 2451-2457.

Usman, A., Komatsu, K., Okada, S., Oikawa, H., Matsuda, H. and Nakanishi, H. (1999) "Synthesis, Crystal Structures and Second-Order Nonlinear Optic Properties of New Colorless 4-Carbamoylpyridinium Benzenesulfonate Salts", *Organic Nonlinear Optical Materials* 3796: 218-228.

Usman, A., Kosuge, H., Okada, S., Oikawa, H. and Nakanishi, H. (2001) "4-(Dimethylamino)-1-ethylpyridinium Iodide: A New Colorless Organic Ionic Crystal for Second-Order Nonlinear Optics", *Jpn. J. Appl. Phys* 40: 4213-4216.

Usman, A., Okada, S., Oikawa, H. and Nakanishi, H. (2000) "Preparation and Crystal Structures of New Colorless 4-Amino-1-methylpyridinium Benzenesulfonate Salts for Second-Order Nonlinear Optics", *Chem. Mater* 12: 1162-1170.

Rahman, A. A., Razak, I. A., Fun, H.-K., Saeneey, P., Boonwasana, J., Chantrapromma, S. and Kararai, C. (2003) "4-(4'-hydroxy-3'-methoxyphenyl)ethenyl-N-methyl pyridinium p-toluenesulfonate", *Acta Cryst.*, E59, o1798-o1800.

Williams, D. J. (1984). *Angew. Chem. Int. Ed. Engl.* 23, 690-703.