



บทที่ 6

เอกสารอ้างอิง

- Ahmad M. S. and Ahmad N. (2006). Antiglycation properties of aged garlic extract: possible role in prevention of diabetic complications. *J. Nutr.* 136: 796S-799S.
- Ali, H., Houghton, P.J. and Soumyanath, A. (2006). α -Amylase inhibitory activity of some Malaysian plants used treat diabetes; with particular reference to *Phyllanthus amarus*. *J. Ethanopharmacology* 107(3): 449-455.
- Bernfeld, P. (1955). Amylases, α and β . *Methods in Enzymology* 1: 149-158.
- Bradford, M. (1976). A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of dye binding. *Anal. Biochem.* 72; 248-254.
- Brunkhorst, C., Andersen, C., and Schneider, E. (1999). Acarbose, a pseudooligosaccharide, is transported but not metabolized by maltose-maltodextrin system of *Escherichia coli*. *J. Bacteriol.* 181(8): 2612-9.
- Cha, H.J., Yoon, H.G., Kim, Y.W., Lee, H.S., Kim, J.W., Kweon, K.S., Oh, B.H., and Park, K.H. (1998). Molecular and enzymatic characterization of a maltogenic amylase that hydrolyzes and transglycosylates acarbose. *European J. Biochem.* 253(1): 251-262.
- Chagolla-Lopez, A., Blanco-Labra, A., Patthy, A., Sa' nchez, R., and Pongor, S. (1994). A Novel α -Amylase Inhibitor from Amaranth (*Amaranthus hypocondriacus*) Seeds. *J. Biol. Chem.* 269(38): 23,675-23,680.
- Chang, H.B., Kim, S.H., Kwon, Y.I., Choung, D.H., Choi, W.K., Kang, T.W., Lee, S., Kim, J.G., Chun, H.S., Ahn, S.K., Hong, C.I. and Han, K.H. (2002). Novel alpha-glucosidase inhibitors, CKD-711 and CKD-711a produced by *Streptomyces* sp. CK-4416. III. Physico-chemical properties and structure elucidation. *J. Antibiotics* 55(5): 467-471.
- Cheah, J.S. (1998). Management of obesity in NIDDM (non-insulin-dependent diabetes mellitus). *Singapore Med. J.* 39(8): 380-384.
- Clark, J.M. and Switzer, R.L. (1997). *Experimental Biochemistry* 2nd ed. Freeman & Company, USA.
- Correia, R.T.P., McCue, P., Vattem, D.A., Magalhaes, M.M.A., Macedo, G.R. and Shetty, K. (2004). Amylase and *Helicobacter pylori* inhibition by phenolic extract of pineapple wastes bioprocessed by *Rhizopus oligosporus*. *J. Food Biochemistry* 28: 419-434.
- Davis, B.J. (1964). Disc electrophoresis II. Method and application tonhuman serum proteins. *Ann. N.Y. Acad. Sci.* 121:404-427.
- Endoa, K., Miyashitaa, Y., Sasakib, H., Ohiraa, M., Saikia, A., Koidea, N., Otsukaa, M., Oyamaa, T.,

- Takeyoshic, M., Itoa, Y. and Shiraid, K. (2006). Probuocol delays progression of diabetic nephropathy. *Diabetes Research and Clinical Practice* 71 (2):156-163.
- Farias, L.R., Costa, F.T., Sonza, L.A., Pelegrini, P.B., Grossi-de-Sa, M.F., Neto, S.M., Bloch Jr., C., Laumann, R.A., Noronha, E.F. and Franco, O.L. (2007). Isolation of a novel *Carica papaya* α -amylase inhibitor with deleterious activity toward *Callosobruchus maculatus*. *Pesticide Biochemistry and Physiology* 87: 255-260.
- Gin, H., Rigalleau, V., Caubet, O., Aubertin, J. (1999). Effects of red wine, tannic acid, or ethanol on glucose tolerance in non-insulin dependent diabetic patients and on starch digestibility *in vitro*. *Metabolism* 48: 1179-1183.
- Giri, A.P. and Kachole, M.S. (1996). Detection of electrophoretically separated amylase inhibitors in starch-polyacrylamide gels. *J. Chromatography A* 752; 261-264.
- Hansawasdi, C., Kawabata, J., and Kasai, T. (2000). Alpha-amylase inhibitors from roselle (*Hibiscus sabdariffa* Linn.) tea. *Bioscience, Biotechnology, and Biochemistry* 64(5): 1041-1043.
- Hara, T., Nakamura, J., Koh, N., Sakakibara, F., Takeuchi, N. and Hotta, N. (1996). An importance of carbohydrate ingestion for the expression of the effect of alpha-glucosidase inhibitor in NIDDM. *Diabetes-Care*. 19: 642-647.
- Huang, C.F., Chen, Y.W., Yang, C.Y., Lin, H.Y., Way, T.D., Chiang, W., and Liu, S.H. (2011). Extract of Lotus Leaf (*Nelumbo nucifera*) and Its Active Constituent Catechin with Insulin Secretagogue Activity. *J. Agric. Food Chem.* 59(4): 1087-1094
- Heidari, R., Zareae, S., and Heidarizadeh, M. (2005). Extraction, purification and inhibitory effect of alpha-amylase inhibitor from wheat (*Triticum aestivum* var. Zarrin). *Pakistan J. Nutrition* 4(2): 101-105.
- Jamaluddin, F., Mohammed, S., and Lajis, M.N. (1994). Hypoglycaemic effect of *Parkia speciosa* seeds due to the synergistic action of beta-sitosterol and stigmasterol. *Food Chem.* 49: 339-345.
- Kandra, L., Gyemant, G., Zajacz, A., and Battab, G. (2004). Inhibitory effects of tannin on human salivary α -amylase. *Biochem Biophys Res Commun.* 319: 1265-1271.
- Kim, J.G., Chang, H.B., Kwon, Y.I., Moon, S.K., Chun, H.S., Ahn, S.K. and Hong, C.I. (2002a). Novel alpha-glucosidase inhibitors, CKD-711 and CKD-711a produced by *Streptomyces* sp. CK-4416. I. Taxonomy, fermentation and isolation. *J. Antibiotics* 55(5): 457-461.
- Kim, M.J., Lee, H.S., Cho, J.-S., Kim, T.J., Moon, T.W., Oh, S.T., Kim, J.W., Oh, B.H., and Park, K.H. (2002b). Preparation and characterization of α -D-glucopyranosyl- α -acarviosinyl-D-glucopyranose, a novel inhibitor specific for maltose-producing amylase. *Biochemistry* 41: 9099-9108.
- Kluh, I., Hom, M., Hyblova, J., Hubert, J., Doleckova-Maresova, L., Voburk, Z., Kudlikova, I, Kocourek,

- F., and Mares, M. (2005). Inhibitory specificity and insecticidal selectivity of α -amylase inhibitor from *Phaseolus vulgaris*. *Phytochem.* 66: 31-39.
- Kwon, Y.I., Son, H.J., Moon, K.S., Kim, J.K., Kim, J.G., Chun, H.S., Ahn, S.K. and Hong, C.I. (2002). Novel alpha-glucosidase inhibitors, CKD-711 and CKD-711a produced by *Streptomyces* sp. CK-4416. II. Biological properties. *J. Antibiotics* 55(5): 462-466.
- Lajolo, F. and Genovese, M.I. (2002). Nutritional significance of lectins and enzyme inhibitors from legumes. *J. Agric. Food Chem.* 50:6592-6598.
- Lankisch, M., Layer, P., Rizza, R.A., and DiMagno, E.P. (1998). Acute postprandial gastrointestinal and metabolic effects of wheat amylase inhibitor (WAI) in normal, obese, and diabetic humans. *Pancreas* 17(2): 176-181.
- Lee, S.C., Gepts, P., and Whitaker, J.R. (2002). Protein structures of common bean (*Phaseolus vulgaris*) α -amylase inhibitors. *J. Agric. Food Chem.* 50: 6618-6627.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., and Randall, R.J. (1951). Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193: 265-275
- Martin, A.E., and Montgomery, P.A. (1996). Acarbose: an alpha-glucosidase inhibitor. *American Journal of Health-System Pharm.* 53: 2277-2290.
- Matsuda, H., Murakami, T., Yashiro, K., Yamahara, J., and Yoshikawa, M. (1999). Antidiabetic principles of natural medicines IV. Aldose reductase and alpha-glucosidase inhibitors from the roots of *Salacia oblonga* Wall (Celastraceae) structure of the new friedelands type triterpane, kotalagenin 16 acetate. *Chemical and Pharmaceutical Bulletin.* 17: 1725-1729.
- McCue, P., Kwon, Y.I. and Shetty, K. (2005). Anti-amylase, anti-glucosidase and anti-angiotensin I-converting enzyme potential of selected foods. *J. Food Chem.* 29:278-294.
- Mukherjee, P.K., S.K. Pal, K. Saha, and B.P. Saha. (1995). Hypoglycaemic activity of *Nelumbo nucifera* Gaertn. (Fam. Nymphaeaceae) rhizome (methanolic extract) in streptozotocin-induced diabetic rats. *Phytotherapy Research* 9:522-524.
- Nakamura, T., Sasaki, I., Masada, M., Tamai, T., and Nakai, T. (1998). Effects of alpha-glucosidase inhibitors by once a day administration on the blood glucose level. *Japanese Journal of Hospital Pharmacy.* 24: 473-478.
- Noor, H., and Ashcroft, S.J.H. (1989). Pharmacological characterisation of the antihyperglycaemic properties of *Tinospora crispa* extract. *Journal of Ethnopharmacology.* 62: 7-13.
- Noor, H., and Ashcroft, S.J.H. (1998). Insulinotropic activity of *Tinospora crispa* extract: effect of beta cell Ca^{2+} handling. *Phytotherapy Research* 12: 98-102.

- de Sales, P.M., de Souza, P.M., Simeoni, L.A., de Oliveira, P., and Silveira, D. (2012). α -Amylase inhibitors: A review of raw material and isolated compounds from plant source. *J Pharmaceul Sci* 15(1):141-183.
- Sawada, S., Takeda, Y., and Tashiro, M. (2002). Primary structure of alpha- and beta-subunits of alpha-amylase inhibitors from seeds of three cultivars of Phaseolus beans. *J. Protein Chem.* 21(1):9-17.
- Sevensson, B., Fukuda, K., Nielson, P.K., and Bonsager, B.C. (2004). Review proteinaceous α -amylase inhibitors. *Biophysica Acta* 1696: 145-146.
- Sharma, A. and Gupta, M.N. (2001). Three phase partitioning as a large-scale separation method for purification of a wheat germ bifunctional protease/amylase inhibitor. *Process Biochemistry* 37:193-196.
- Shibib, B.A., Khan, L.A., and Rahman, R. (1993). Hypoglycaemic activity of *Coccinia indica* and *Momordica charantia* in diabetic rats : depression of the hepatic gluconeogenic enzyme glucose-6-phosphatase and fructose-1, 6-bisphosphatase and elevation of both liver and red cell shunt enzyme glucose-6-phosphate dehydrogenase. *Biochem. J.* 292: 267-270.
- Sidenius, U., Olsen, K., Svensson, B., and Christensen, U. (1995). Stopped-flow Kinetic Studies of the Reaction of Barley α -Amylase/Subtilisin Inhibitor and the High pI Barley α -Amylase. *FEBS Letters* 361: 250-254.
- Silva, B.M., Andrade P.B., Mendes, G.C., Seabra, R.M. and Ferreira, M.A. (2002). Study of the organic acids composition of quince (*Cydonia oblonga* Miller) fruit and jam. *J. Agri. Food Chem.* 50: 2313-2317.
- Slinkkard, K. and Singleton, K. (1997). Total phenol analysis: Automation and comparison with manual methods. *Am. J. Enol. Viticult.* 28:49-55.
- Yamagata, H., Kunitatsu, K., Kamasaka, H., Kuramoto, T. and Iwasaki, T. (1998). Rice Bifunctional α -Amylase/Subtilisin Inhibitor: Characterization, localization, and changes in developing and maturing Seeds. *Biosci. Biotechnol. Biochem.* 62(5): 978-985.
- Ye, F., Shen, Z., and Xie, M. (2002). Alpha-glucosidase inhibition from a Chinese medical herb (*Ramulus mori*) in normal and diabetic rats and mice. *Phytomedicine* 9:161-166.
- Yuasa, H., Takada, J., and Hashimoto, H. (2000). Synthesis of salacinol. *Tetrahedron letters.* 41: 6615-6618.
- Zang Y, Sato H, and Igarashi K. (2011). Anti-diabetic effects of a kaempferol glycoside-rich fraction from unripe soybean (Edamame, *Glycine max* L. Merrill. 'Jindai') leaves on KK-A(y) mice. *Biosci. Biotechnol. Biochem.* 75(9):1677-84.

www.nutriceps.com/ATW.htm

www.progenebio.in/DMP

กรรณา จีนถนอม (2550). เหตุเกิดจากความอ้วน. วารสารโภชนาการ 42(1):39-41.

เพียววี เหมือนวงษ์ญาติ (2534). คู่มือการใช้สมุนไพร. พิมพ์ครั้งที่ 5: หจก.ภาพพิมพ์ กรุงเทพฯ.

ภูมิพิชญ์ สุชาวรรณ และปรีชา กาบแตก (2535). สมุนไพรใช้เป็นยา: โรงพิมพ์ทิพย์วิสุทธิ กรุงเทพฯ.

มาโนช วามานนท์ และแพทย์หญิงเพ็ญนภา ทรัพย์เจริญ (2537). ยาสมุนไพรสำหรับงานสาธารณสุขมูลฐาน: โรงพิมพ์องค์การสงเคราะห์ทหารผ่านศึก กรุงเทพฯ.

วิทยา ศรีดามา และศักดิ์ชัย ปาละวัฒน์ (2541). ยาเมล็ดระดับน้ำตาลในเลือด: การดูแลรักษาผู้ป่วยเบาหวาน: ยูนิตี้ พับลิเคชั่น กรุงเทพฯ.

วุฒิ วุฒิชรรมเวช (2541). สรรพคุณสมุนไพร: ส่วนที่นิยมใช้ทำยา: เพชร 69 การพิมพ์ กรุงเทพฯ.

สำนักงานคณะกรรมการสาธารณสุขมูลฐาน (2541). สมุนไพรในงานสาธารณสุขมูลฐาน: สำนักพิมพ์ดอกหญ้า กรุงเทพฯ.



