



บรรณานุกรม

- [1] WEAVER, A.L. (2008). Epidemiology of gout. *Cleve Clin J Med*, 75(2), 9-12.
- [2] ไพจิตร ธีศรณบดี และอัจฉรา กุลวิสุทธิ. (2549). โรคข้อสำหรับบุคลากรทางการแพทย์และประชาชน. กรุงเทพฯ: สมาคมรูมาติสซั่มแห่งประเทศไทย.
- [3] Kramer, H.M., and Curhan, G. (2002). The association between gout and nephrolithiasis: the national health and nutrition examination survey III, 1988-1994. *Am J Kidney Dis*, 40(1), 37-42.
- [4] Meisenberg, G. and Simmons, W.H. (2006). *Principles of medical biochemistry* (2nd ed.). Mosby: Louis.
- [5] ศุภราภรณ์ วังแก้ว. (2549). โรคข้ออักเสบจากผลึกเกลือ. กรุงเทพฯ: ซีดีพีริ้น.
- [6] พงษ์ศักดิ์ วัฒนมา, ธวัช ประสาทฤทธา และวัลลภ สำราญเวทย์ (บรรณาธิการ). (2533). *การวินิจฉัยและการรักษาโรคข้อที่พบบ่อย*. กรุงเทพฯ: โครงการควบคุมโรคข้อแผนงานควบคุมโรคไม่ติดต่อ กรมการแพทย์ กระทรวงสาธารณสุข.
- [7] Alexander C.B., and John, C. K. (2005). *Medical biochemistry: A core text with self-assessment* (2nd ed.). Edinburgh: Elsevier/Churchill Livingstone.
- [8] Becker, M.A. (1997). *Clinical gout and the pathogenesis of hyperuricemia*. N.p.: n.p.
- [9] Gilani, A. H., and Atta-ur-Rahman. (2005). Trends in ethnopharmacology. *Journal of Ethnopharmacology*, 100(1-2), 43-49.
- [10] กัญญา ดิวีเศษ. (2542). *เภสัชกรรมแผนไทย*. กรุงเทพฯ: โรงพิมพ์องค์การสงเคราะห์ทหารผ่านศึก.
- [11] นิจศิริ เรืองรังษี และธวัชชัย มังคละคุปต์. (2547). *สมุนไพรไทย* (เล่ม 1). กรุงเทพฯ: บี เฮลท์ดี.
- [12] Umamaheswari, M., AsokKumar, K., Somasundaram, A., Sivashanmugam, T., Subhadradevi, V., and Ravi, T.K. (2007). Xanthine oxidase inhibitory activity of some Indian medical plants. *Journal of Ethnopharmacology*, 109(3), 547-551.
- [13] สุธี วรศิรินิมิต. (2545). *มหัศจรรย์พันธุ์พืชสมุนไพร เครื่องดื่มป้องกันโรค*. กรุงเทพฯ: พิมพ์ทอง.

- [14] Merry, R. S., Mark, G.P.S., and Fernando, Perez-Ruiz. (2007). PEG-uricase in the management of treatment-resistant gout and hyperuricemia. *Advanced Drug Delivery Reviews*, 60(1), 59-68.
- [15] Choi, H. K., Mount, D.B., and Reginato, A.M. (2005). Pathogenesis of gout. *AnnIntern Med*, 143(7), 499-516.
- [16] Bomalaski, J.S., Holtsberg, F.W., Ensor, C.M., and Clark, M.A. (2002). Uricase formulated with polyethylene glycol (uricase-PEG 20): biochemical rationale and preclinical studies. *Journal of rheumatology*, 29(9), 1942 -1949.
- [17] CHOI, H.K., FORD, E.S., CHAOYANG, L., and CURHAN, G. (2007). Prevalence of the metabolic syndrome in patients with gout: the third national health and nutrition examination survey. *Arthritis care and research*, 57(1), 109-115.
- [18] Mikuls, T.R., Farra, J.T., Bilke, W.B., Fernandes, S., Schumacher Jr, H.R., and Saag, K.G. (2005). Gout epidemiology: results from the UK general practice research database, 1990–1999. *Ann Rheum Dis*, 64(2), 267-272.
- [19] Ronenn Roubenoff, R., Klag, M.J., Mead, L.A., Liang, K.Y., Seidler, A.J., and Hochberg, M.C. (1991). Incidence and risk factors for gout in white men. *JAMA*, 266(21), 3004-3007.
- [20] CASPI, D., LUBART, E., GRAF, E., HABOT, B., YARON, M., and SEGAL, R. (2000). The effect of mini-dose aspirin on renal function and uric acid handling in elderly patients. *Arthritis & Rheumatism*, 43(1), 103-108.
- [21] Choi, H.K., Atkinson, K., Karlson, E.W., and Curhan, G. (2005). Obesity, weight change, hypertension, diuretic use, and risk of gout in men: the health professionals follow-up study. *Arch Intern Med*, 165(7), 742-748.
- [22] Choi, H.K., Atkinson, K., Karlson, E.W., Willett, W., and Curhan, G. (2004). Purine-rich foods, dairy and protein intake, and the risk of gout in men. *N Engl J Med*, 350(11), 1093-1103.
- [23] Choi, H.K. (2005). Diet, alcohol, and gout: how do we advise patients given recent developments?. *Current Rheumatology Reports*, 7(3), 220-226.
- [24] วิจิตร บุญยะโหดระ. (2534). *อาหารรักษาโรค (พิมพ์ครั้งที่ 2)*. กรุงเทพฯ: ดวงกมลสมัย.

- [25] Kang, D.H., Nakagawa, Feng, L., Watanabe, S., Han, L., Mazzali, M., Truong, L., et al. (2002). A role for uric acid in the progression of renal disease. *J Am Soc Nephrol*, 13(12), 2888-2897.
- [26] Johnson, R. J., Segal, M. S., Srinivas, T., Ejaz, A., Mu, W., Roncal, C., et al. (2005). Essential hypertension, progressive renal disease, and uric acid: a pathogenetic link?. *J Am Soc Nephrol*, 16(7), 1909-1919.
- [27] Krishnan, E., Baker, J.F., Furst, D.E., and Schumacher, H.R. (2006). Gout and the risk of acute myocardial infarction. *Arthritis Rheum*, 54(8), 2688-2696.
- [28] Krishnan, E., Kwoh, C.K., Schumacher, H.R., and Kuller, L. (2007). Hyperuricemia and incidence of hypertension among men without metabolic syndrome. *Hypertension*, 49(2), 298-303.
- [29] Jerome, J.T., Varghese, M., Sankaran, B., Thomas, S., and Thirumagal, S.K. (2008). Tibialis anterior tendon rupture in gout-case report and literature review. *Foot Ankle Surg*, 14(3), 166-169.
- [30] Choi, H.K., and Curhan, G. (2007). Alcohol and gout. *The American Journal of Medicine*, 120(10), 1.
- [31] Faller, J., and Fox, I.H. (1982). Ethanol-induced hyperuricemia: evidence for increased urate production by activation of adenine nucleotide turnover. *N Engl J Med*, 307(26), 1598 – 1602.
- [32] Owen, P.L. and Johns, T. (1999). Xanthine oxidase inhibitory activity of northeastern north american plant remedies used for gout. *Journal of Ethnopharmacology*, 64(2), 149-160.
- [33] Schardinger, F. (1902). Ueber das verhalten der kuhmilch gegen methylenblau und seine verwendung zur unterscheidung von ungekochter und gekochter milch. *Zeitschrift für Lebensmitteluntersuchung und -Forschung A*, 5(22), 1113-1121.
- [34] Akdemir, H., Asik, Z., Pasaoglu, H., Karakucuk, I., Oktem, I.S., and Koc, R.K. (2001). The effect of allopurinol on focal cerebral ischaemia: an experimental study in rabbits. *Neurosurg Rev*, 24(2-3), 131-135.

- [35] Warner, D.S., Sheng, H. and Batinic, H.I. (2004). Oxidants, antioxidants and the ischemic brain. *Journal of Experimental Biology*, 207(18), 3221 – 3231.
- [36] Patt, A., Harken, A.H., Burton, L. K., Rodell, T.C., Piermattei, D., Schorr, W.J., et al. (1988). Xanthine oxidase-derived hydrogen peroxide contributes to ischemia reperfusion-induced edema in gerbil brains. *J Clin Invest*, 81(5), 1556-1562.
- [37] Anup, R., and Balasubramanian, K. A. (2000). Surgical stress and the gastrointestinal tract. *Journal of Surgical Research*, 92(2), 291-300.
- [38] Harrison, R. (2002). Structure and function of xanthine oxidoreductase: where are we now?. *Free Radical Biology and Medicine*, 33(6), 774-797.
- [39] Cheng, H.Y., Lin, T.C., Yu, K.H., Yang, C.M., and Lin, C.C. (2003). Antioxidant and free radical scavenging activities of terminalia chebula. *Biol Pharm Bull*, 26(9), 1331-1335.
- [40] Borges, F., Fernandes, E., and Roleira, F. (2002). Progress towards the discovery of xanthine oxidase inhibitors. *Curr Med Chem*, 9(2), 195-217.
- [41] Enroth, C., Eger, B.T., Okamoto, K., Nishino, T., and Pai, E.F. (2000). Crystal structures of bovine milk xanthine dehydrogenase and xanthine oxidase: structure-based mechanism of conversion. *Proc Natl Acad Sci U S A*, 97(20), 10723-10728.
- [42] Cos, P., Ying, L., Calomme, M., Hu, J.P., Cimanga, K., Poel, B.V., Pieters, L., et al. (n.d.) Structure-activity relationship and classification of flavonoids as inhibitors of xanthine oxidase and superoxide scavengers. *J Nat Prod*, 61 (1), 71-76.
- [43] Berry, C.E., and Hare, J.M. (2004). Xanthine oxidoreductase and cardiovascular disease: molecular mechanisms and pathophysiological implications. *J Physiol*, 555(3), 589-606.
- [44] Saksela, M., Lapatto, R., and Raivio, K.O. (1999). Irreversible conversion of xanthine dehydrogenase into xanthine oxidase by a mitochondrial protease. *FEBS Letters*, 44(2), 117-120.

- [45] Terada, L.S., Leff, J.A., Repine, E., Lester, P., and Alexander, N. G. (1990). Measurement of xanthine oxidase in biological tissues, in *Methods in Enzymology*. Academic Press, 651-656.
- [46] ดวงมณี วิเศษกุล. (2525). *โรคและอาหารเฉพาะโรค (พิมพ์ครั้งที่ 2)*. กรุงเทพฯ: โครงการตำราศิริราช.
- [47] วรวิทย์ เล่าห์เรณู (บรรณารักษกร). (2548). *โรคข้อจากผลึกเกลือ*. เชียงใหม่: หน่วยโรคข้อและรูมาติสซั่ม ภาควิชาอายุรศาสตร์ คณะแพทยศาสตร์ มหาวิทยาลัยเชียงใหม่.
- [48] Nelson, D.V., and Cox, M.M. (2004). *Lehninger principles of biochemistry* (4th ed.). New York: W.H. Freeman.
- [49] Emmerson, B.T. (1996). The management of gout. *N Engl J Med*, 334(7), 445-451.
- [50] Schlesinger, N., and Schumacher, H.R.J. (2001). Gout: can management be improved?. *Current Opinion in Rheumatology*, 13(3), 240-244.
- [51] Fields, M., Lewis, C.G., and Lure, M.D. (1996). Allopurinol an inhibitor of xanthine oxidase reduces uric acid levels and modifies the signs associated with copper deficiency in rats fed fructose. *Free Radical Biology and Medicine*, 20(4), 595-600.
- [52] Ishibuchi, S., Morimoto, H., Oe, T., Ikebe, T., Inoue, H., Fukunari, A., et al. (2001). Synthesis and structure-activity relationships of 1-Phenylpyrazoles as xanthine oxidase inhibitors. *Bioorganic & Medicinal Chemistry Letters*, 11(7), 879-882.
- [53] Taylor, M.D., Palmer, G.C., and Callahan, A. S. (1984). Protective action by methylprednisolone, allopurinol and indomethacin against stroke-induced damage to adenylate cyclase in gerbil cerebral cortex. *Stroke*, 15(2), 329-335.
- [54] Hammer, B., Link, A., Wagner, A., and Bohm, M. (2001). Hypersensitivity syndrome during therapy with allopurinol in asymptomatic hyperuricemia with a fatal outcome. *Dtsch Med Wochenschr*, 126(47), 1331-1334.
- [55] Chen, I.H., Kuo, M.C., Hwang, S.J., Chang, J.M., and Chen, H.C. (2005). Allopurinol-induced severe hypersensitivity with acute renal failure. *Kaohsiung J Med Sci*, 21(5), 228-232.

- [56] ศุภลักษณ์ ศรีจรรย์. (2544). โครมาโทกราฟีของเหลวสมรรถนะสูง: ทฤษฎีและปฏิบัติ. ขอนแก่น: ภาควิชาเคมี คณะวิทยาศาสตร์ มหาวิทยาลัยขอนแก่น.
- [57] วชิร ชาติกิตติคุณวงศ์. (2542). เคมีวิเคราะห์ 2 เรื่องโครมาโทกราฟีของเหลวที่มีสมรรถนะสูง CH334. กรุงเทพฯ: สำนักพิมพ์แห่งมหาวิทยาลัยรามคำแหง.
- [58] เย็นหทัย แนนหนา. (2549). สเปกโทรสโกปีสำหรับเคมีอินทรีย์. กรุงเทพฯ: สำนักพิมพ์แห่งจุฬาลงกรณ์มหาวิทยาลัย.
- [59] ธิติมา รุกชไชยศิริกุล. (2538). เคมีอินทรีย์ 2 CH 323 (พิมพ์ครั้งที่ 3). กรุงเทพฯ: สำนักพิมพ์แห่งมหาวิทยาลัยรามคำแหง.
- [60] สมเดช กนกเมธากุล. (2547). สเปกโทรสโกปี ในการพิสูจน์โครงสร้างของสารอินทรีย์. ขอนแก่น: ขอนแก่นการพิมพ์.
- [61] Hsieh, J.F., Wu, S.H., Yang, Y.L., Choong, K.F., and Chen, S.T. (2007). The screening and characterization of 6-aminopurine-based xanthine oxidase inhibitors. *Bioorganic & Medicinal Chemistry*, 15(10), 3450-3456.
- [62] Kumar, U.S., Tiwari, A.K., Reddy, S.V., Aparna, P., Rao, R.J., Ali, A.Z., and Rao, J.M. (2005). Free-radical-scavenging and xanthine oxidase inhibitory constituents from *stereospermum personatum*. *J. Nat. Prod*, 68(10), 1615-1621.
- [63] Kong, L., Zhou, J., Wen, Y., Li, J., and Cheng, C.H.K. (2002). Aesculin possesses potent hypouricemic action in rodents but is devoid of xanthine oxidase/dehydrogenase inhibitory activity. *Planta Med*, 68(2), 175-178.
- [64] Liu, X., Chen, R., Shang, Y., Jiao, B., and Huang, C. (2008). Lithospermic acid as a novel xanthine oxidase inhibitor has anti-inflammatory and hypouricemic effects in rats. *Chemico-Biological Interactions*, 176(2-3), 137-142.
- [65] Eduardo Vivot, E., Juan de Dios, J.D., Josefina, M., Tapia, A., (2001). Inhibitory activity of xanthine-oxidase and superoxide scavenger properties of *Inga verna* subsp. *affinis*. Its morphological and micrographic characteristics. *Journal of Ethnopharmacology*, 65-71.

- [66] Nguyen, M.T.T., Awale, S., Tezuka, Y., Tran, Q.L., and Kadota, S. (2005). Xanthine oxidase inhibitors from the heartwood of vietnamese caesalpinia sappan. *Chem Pharm Bull (Tokyo)*, 53(8), 984-988.
- [67] Kong, L.D., Cai, Y., Huang, W.W., Cheng, C.H.K., and Tan, R.X. (2000). Inhibition of xanthine oxidase by some Chinese medicinal plants used to treat gout. *Journal of Ethnopharmacology*, 73(1-2), 199-207.
- [68] สมพร ภูติยานันต์. (2542). การตรวจเอกลักษณ์พืชสมุนไพร. นนทบุรี: โครงการพัฒนาตำรา สถาบันการแพทย์แผนไทย กรมการแพทย์ กระทรวงสาธารณสุข.
- [69] Grases, F., March, J.G., Ramis, M., and Costa, B.A. (1993). The influence of zea mays on urinary risk factors for kidney stones in rats. *Phytotherapy Research*, 7(2), 146-149.
- [70] Newall, C., Anderson, L.A., and Phillipson, J.O. (1996). *Herbal medicine, a guide for health - care professionals*. London: U.K.
- [71] Maksimovic, Z.A., and Kovacevic, N. (2003). Preliminary assay on the antioxidative activity of maydis stigma extracts. *Fitoterapia*, 74(1-2), 144-147.
- [72] Ebrahimzadeh, M.A., Pourmorad, F., and Hafezi, S. (2008). Antioxidant activities of iranian corn silk. *Turk Journal of Bioogy*, 32(1), 43-49.
- [73] Leung, Albert Y. (1980). *Encyclopedia of common natural ingredients: Used in food, drygs, and cosmetics*. New York: n.p.
- [74] Zeringue, H.J. (2000). Identification and effects of maize silk volatiles on cultures of aspergillus flavus. *J Agric Food Chem*, 48(3), 921-925.
- [75] Habtemariam, S. (1998). Extract of corn silk (stigma of Zea mays) inhibits the tumour necrosis factor-alpha- and bacterial lipopolysaccharide-induced cell adhesion and ICAM-1 expression. *Planta Med*, 64(4), 314-318.
- [76] Saousse, H., Sadok, B., Habib, K., and Mayer, P.M. (2009). Fatty acid accumulation in the different fractions of the developing corn kernel. *Food Chemistry*, 117(3), 432-437.

- [77] Pourmorad, F., Hosseinimehr, S.J., and Shahabimajd, N. (2006). Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *African Journal of Biotechnology*, 5(11), 1142-1145.
- [78] Benzie, I.F.F., and Strain, J.J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Analytical Biochemistry*, 231(1), 70-76.
- [79] Griffin, S.P. and Bhagooli, R. (2004). Measuring antioxidant potential in corals using the FRAP assay. *Journal of Experimental Marine Biology and Ecology*, 302(2), 201-211.
- [80] Saeed, M.K., Yulin, D., Rongji, D. and Yuhong, Y. (2007). Optimization of condition for maximum recovery of antioxidants from cephalotaxus sinensis and their FRAP assay. *IEEE/ICME International Conference*, 23-27, 1664-1667.
- [81] มหาวิทยาลัยเชียงใหม่. (2545). คู่มือปฏิบัติการชีวเคมี. เชียงใหม่: ภาควิชาชีวเคมี คณะแพทยศาสตร์ มหาวิทยาลัยเชียงใหม่.
- [82] ตรีเพชร กาญจนภูมิ. (2552). เคมีของสมุนไพร: การหาโครงสร้างเคมีของสาร แอลโรแมติกไกลโคไซด์ด้วยนิวเคลียร์แมกเนติกเรโซแนนซ์ สเปกโทรสโกปี. กรุงเทพฯ: สำนักพิมพ์แห่งจุฬาลงกรณ์มหาวิทยาลัย.
- [83] Pazmino-Duran, E.A., Monica Giusti, M., Wrolstad, R.E., Gloria, M., and Beatriz, A. (2001). Anthocyanins from oxalis triangularis as potential food colorants. *Food Chemistry*, 75(2), 211-216.
- [84] บุปผาชาติ พตด้วง, ธัญวรรณ์ กาจสงคราม, มณีรัตน์ มีพลอย และพงศธร หลิมศิริวงษ์. การวิจัยข้ามหลามดง: การวิเคราะห์หากลุ่มสารหลัก. ม.ป.ท.: ม.ป.พ.
- [85] Chen, X., and Xiao, J. (1996). RP-HPLC-DAD determination of flavonoids: separation of quercetin, luteolin and apigenin in *marchantia*. *Nigerian Journal of Natural Products and Medicine*, (3), 175-181.
- [86] Zhou, X., Peng, J., Fan, G., and Wu, Y. (2005). Isolation and purification of flavonoid glycosides from *trollius ledebouri* using high-speed counter-current chromatography by stepwise increasing the flow-rate of the mobile phase. *Journal of Chromatography A*, 1092(2), 216-221.

- [87] Zhao, G., Qin, G.W., Wang, J., Chu, W.J., and Guo, L. (2009). Functional activation of monoamine transporters by luteolin and apigenin isolated from the fruit of *Perilla frutescens* (L.) Britt. *Neurochemistry International*, 56(1), 168-176.
- [88] Ahmad, N.S., Farman, M., Najmi, M.H., Mian, K.B., and Hasan, A. (2008). Pharmacological basis for use of *Pistacia integerrima* leaves in hyperuricemia and gout. *Journal of Ethnopharmacology*, 117(3), 478-482.
- [89] Orech, F.O., Akenga, T., Ochora, J., Friis, H., and Hansen, J.A. (2005). Potential Toxicity of Some Traditional Leafy Vegetables Consumed in Nyang' oma Division, Western Kenya. *African Journal of Food and Nutritional Sciences*, 5(1), 1-13.
- [90] พัชรา วีระกะลัศ. (2541). เอนไซม์. กรุงเทพฯ: สำนักพิมพ์แห่งจุฬาลงกรณ์มหาวิทยาลัย.
- [91] Adom, K.K., and Rui Hai Liu, R.H. (2002). Antioxidant activity of grains. *J. Agric. Food Chem*, 50(21), 6182- 6187.
- [92] Cakir, A., Mavi, A., Yildirim, A., Duru, M.E., Harmandar, M., and Kazaz, C. (2003). Isolation and characterization of antioxidant phenolic compounds from the aerial parts of *Hypericum hyssopifolium* L. by activity-guided fractionation. *Journal of Ethnopharmacology*, 87(1), 73-83.
- [93] Wang, C.K., and Lee, W.H. (1996). Separation, characteristics, and biological activities of phenolics in areca fruit. *J. Agric. Food Chem*, 44(8), 2014-2019.
- [94] Wange, S.Y., and Lint, H.S. (2000). Antioxidant activity in fruits and leaves of blackberry, raspberry, and strawberry varies with cultivar and developmental stage. *Journal of agricultural and food chemistry*, 28, 140-146.
- [95] Yang, C.S. (2001). Inhibition of carcinogenesis and toxicity by dietary constituents. *Adv Exp Med Biol*, 541-50.
- [96] Ozyurek, M., Bektasoglu, B., Guclu, K., and Apak, R. (2009). Measurement of xanthine oxidase inhibition activity of phenolics and flavonoids with a modified cupric reducing antioxidant capacity (CUPRAC) method. *Analytica Chimica Acta*, 636(1), 42-50.

- [97] Dewanto, V., Wu, X., and Liu, R.H. (2002). Processed sweet corn has higher antioxidant activity. *J Agric Food Chem*, 50(17), 4959- 4964.
- [98] Pedreschi, R., and Cisneros-Zevallos, L. (2007). Phenolic profiles of Andean purple corn (*Zea mays* L.). *Food Chemistry*, 100(3), 956-963.
- [99] ระวีวรรณ แก้วอมตวงศ์ และทรงพร จิ่งมันคง. (2549).ฤทธิ์ต้านอนุมูลอิสระ DPPH และปริมาณสารฟีนอลรวมของสารสกัดพืชสมุนไพรไทยบางชนิด. *วารสารวิชาการมหาวิทยาลัยอุบลราชธานี*. 8(2), 76-78.
- [100] Wagner, H., and Sabine Bladt, S. (1996). *Plant drug analysis: a thin layer chromatography atlas* (2nd ed.). Berlin: Springer.
- [101] Fleming, T. (2000). *Herbal Medicines* (2nd ed.). New Jersey: Medical Economics Company.
- [102] Wahab, A., Tanbouly, E., Kassem, H.A., and Mohamed, E.A. (2002). Phytochemical and biological study of corn silk (styles and stigmas of *Zea mays* L). *Bull Fac Pharm*, 41(1), 93-102.
- [103] Chang, Y. C., Lee, F.W., Chen, C.S., Huang, S.T., Tsai, S.H., Huang, S.H., et al. (2007). Structure-activity relationship of C6-C3 phenylpropanoids on xanthine oxidase-inhibiting and free radical-scavenging activities. *Free Radical Biology and Medicine*, 43(11), 1541-1555.
- [104] Mo, S.F., Zhou, F., Lv, Y.Z., Hu, Q.H., Zhang, D.M., and Kong, L.D. (2007). Hypouricemic action of selected flavonoids in mice: structure-activity relationships. *Biol Pharm Bull*, 30(8), 1551-1556.
- [105] Pauff, J.M., and Hille, R. (2009). Inhibition studies of bovine xanthine oxidase by luteolin, silibinin, quercetin, and curcumin. *J Nat Prod*, 72(4), 725-731.
- [106] Lin, C.M., Chen, C.S., Chen, C.T., Liang, Y.C., and Lin, J.K. (2002). Molecular modeling of flavonoids that inhibits xanthine oxidase. *Biochemical and Biophysical Research Communications*, 294(1), 167-172.

- [107] Da Silva, S.L., Da Silva, A., Honorio, K.M., Marangoni, S., Toyama, M.H., and Da Silva, A.B.F. (2004). The influence of electronic, steric and hydrophobic properties of flavonoid compounds in the inhibition of the xanthine oxidase. **Journal of Molecular Structure: Theochem**, 684(1-3), 1-7.
- [108] Wang, S.P., and Huang, K.J. (2004). Determination of flavonoids by high-performance liquid chromatography and capillary electrophoresis. **Journal of Chromatography**, 1032(1-2), 273-279.
- [109] Prabhakar, M.C., Bano, H., Kumar, I., Shamsi, M.A., and Khan, S.Y. (1981). Pharmacological Investigations on Vitexin. **Planta Med**, 43(12), 396-403.
- [110] Agnese, A.M., Perez, C., and Cabrera, J.L. (2001). Adesmia aegiceras: antimicrobial activity and chemical study. **Phytomedicine: international journal of phytotherapy and phytopharmacology**, 8(5), 389-394.
- [111] Picerno, P., Mencherini, T., Lauro, M., Barbato, F., and Aquino, R. (2003). Phenolic Constituents and Antioxidant Properties of Xanthosoma violaceum Leaves. **Journal Of Agricultural And Food Chemistry**, 51(22), 6423-6428.
- [112] Bramati, L., Aquilano, F., and Pietta, P. (2003). Unfermented Rooibos Tea: Quantitative Characterization of Flavonoids by HPLC-UV and Determination of the Total Antioxidant Activity. **Journal Of Agricultural And Food Chemistry**, 51(25), 7472-7474.