

Original article

Indoor air microbial counts in dormitory of nursing students, Bangkok

Pipat Luksamijarulkul^{a*}, Maneenut Wutthikarn^b, Poolsuk Janepanish Visudtibhan^c
Pisit Vatanasomboon^d

^aFaculty of Health Sciences, Siam Technology College /Faculty of Public Health, Mahidol University, Bangkok, Thailand

^bMinistry of Public Health, Nonthaburi Province, Thailand

^cFaculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand

^dFaculty of Public Health, Mahidol University, Bangkok, Thailand

Background: Most medical and nursing students live in institute dormitories located in a university hospital where a variety of biological and chemical hazards are used and probably diffused into the dormitory environments.

Objectives: This study aimed to investigate bacterial and fungal counts, and microbial groups in a dormitory of nursing students in Bangkok, Thailand.

Methods: Two-time assessments of bacterial and fungal counts in indoor air samples collected from a dormitory of nursing students including 8 floors with 95 rooms were carried out. A total of 380 indoor air samples (190 for bacteria and 190 for fungi) were collected from all dormitory rooms using BioStage Impactor and 48 outdoor air samples were collected for comparison. Air samples were collected two times, i.e., in the first month and the third month of the study.

Results: The means bacterial counts were 151 ± 109 cfu/m³ and 158 ± 92 cfu/m³ and those of fungal counts were 374 ± 273 cfu/m³ and 363 ± 257 cfu/m³ from the first and the second assessment. The lower floors of the dormitory (1 - 4 floors) had significantly higher levels of bacterial and fungal counts in both assessments when compared with the higher floors of the dormitory (5 - 8 floors) ($P < 0.001$). The percentages of bacterial and fungal counts with > 300 cfu/m³ in the lower floors were significantly higher than those in the higher floors in the first and the second assessment ($P < 0.05$ and $P < 0.001$, respectively). Additionally, the most common isolated bacteria and fungi were *Staphylococcus spp.*, and *Aspergillus spp.*, respectively.

Conclusion: This study reveals that the lower floors of the dormitory had significantly higher levels of bacterial and fungal counts when compared with the higher floors of the dormitory from both assessments. Most isolated microbes do not generally present a health hazard but high loads may trigger allergic reactions in susceptible hosts.

Keywords: Indoor air, bacterial counts, fungal counts, nursing student dormitory.

Indoor air quality (IAQ) is a major concern to businesses, schools, building managers, tenants, and workers because it can impact the health, comfort, well-being, and productivity of the building occupants.^(1, 2) Good indoor air quality leads to improve productivity at the workplace. On the other hand, poor indoor air quality causes productivity to

drop because of comfort problems, illnesses of health and sickness absenteeism.⁽¹⁾ At present, most people spend up to 90% of their time indoors and in an office environment.⁽³⁾ The World Health Organization (WHO) estimates that 30 % of the buildings may have problems with indoor air quality which will lead to sick building syndrome.⁽⁴⁾

Microbial indoor air comes from hundreds of species of bacteria, fungi and moulds that grow indoors when sufficient moisture is available. Exposure to microbial contaminants is clinically associated with respiratory symptoms, allergies and asthma, and can affect the immunological system.⁽⁴⁻⁷⁾ A previous study reported that dampness

*Correspondence to: Pipat Luksamijarulkul, Faculty of Health Sciences, Siam Technology College, Bangkok 10600, Thailand.

E-mail: pipatl@siamtechno.ac.th, pipat.luk@mahidol.ac.th

Received: February 11, 2019

Revised: April 2, 2019

Accepted: April 18, 2019

problem at dorms of Chinese students was a risk factor in irritating allergic symptoms; hence, there is a need for dorm environment improvement.⁽⁸⁾ Another previous study demonstrated that dorm rooms is a kind of residential environment for students, and may be more polluted than the home environment, especially when dorms tend to be more crowded due to a low ventilation rate.⁽⁹⁾ Furthermore, healthcare workers, such as doctors, nurses and others are exposed to infectious diseases and demanding work conditions in healthcare settings.^(10,11) In addition to the danger of air-borne microbes and diseases being spread in hospitals, there are also a number of chemicals released from typical hospital activities that occupants should be aware of.⁽¹²⁾ A study in a Thai university dormitory reported that most students spent time in their university dormitory. The dormitory should have good air quality, convenient, clean, hygienic and safe which will enhance not only the quality of life, physical and mental health, but also the academic performance achievement.⁽¹³⁾ However, this study assessed only physical environment and sanitation of the dormitory; the biological contaminants were not included.

Medical and nursing students living in their dormitories might be at risk for acquiring biological hazards from the dormitories because most of the dormitories are located in the university hospitals. There was a variety of infectious hazards in hospital environments which might probably affect dormitory environments. In a recent period, there have some complaints from some nursing students living in the institute dormitory about symptoms related to indoor air quality. The 2-time assessments of microbial indoor air quality in this nursing student dormitory were carried out to investigate bacterial and fungal counts, and microbial groups.

Materials and methods

Study design

This study design was a cross-sectional study on 2-time assessments of microbial indoor air quality in dormitory rooms of nursing students in 2015. The first time of air sample collection was performed in the first month after complaints from some nursing students about symptoms related to indoor air quality, and the second time was performed again in the third month of the study to confirm the association between indoor air quality and related symptoms (data are not presented in this study). All air samples were collected by using BioStage Impactor QuickTake 30 sample

pump (SKC Inc, USA) to assess total bacterial and fungal counts. A nursing student dormitory included 8 floors with 95 rooms (approximately 15 m² for each room). In each assessment, at least 2 air sampling points in each room were collected and some outdoor air samples were collected in the same period of indoor air collection day for comparison following the guidelines for good indoor air quality in office premises, Ministry of the Environment, Singapore (1996).⁽¹⁴⁾ As for the outdoor air samples, 2 - 4 air sampling points in each floor of the dormitory were collected. Totally, 380 indoor air samples (190 for bacteria and 190 for fungi) and 48 outdoor air samples (24 for bacteria and 24 for fungi) were included in each assessment time. This study was a part of the research protocol approved by the Ethics Committee for Human Research, the Faculty of Medicine, Ramathibodi Hospital, Mahidol University (Ref. No. 10-56-01).

The microorganisms collected by BioStage Impactor are impacted onto an agar surface in accordance with the USP reference method (USP 24, p. 2099). A sample pump draws the air through the sampler where multiple jets of air direct airborne particles toward the surface of the agar collection medium. BioStage Impactor contains a 400 hole jet classification stage and is operated at 28.3 L/min for four minutes. In this study, 113.2 liters of air volume were collected in this study. The air collection technique was followed the active air sampling method of Pasquarella C, *et al*, 2008.⁽¹⁵⁾ Details were described in Luksamijarulkul P, *et al*, 2015.⁽¹⁶⁾ The plate count method was used to estimate bacterial or fungal counts. General bacteria were cultured in plate count agar at 37°C for 48 hours, and general fungi were cultured in a Sabouraud 4% dextrose agar, at room temperature, for 5 days with daily observation. After incubation, the bacterial and fungal colonies were counted and calculated to express as colony forming unit/m³ (cfu/m³) by the following:

$$\text{Microbial counts (colony forming unit/m}^3\text{)} = \frac{\text{Total colonies} \times 10^3}{28.3 \times 4}$$

The isolated colonies of bacteria and fungi were identified group or genus by gram staining and with lacto-phenol cotton blue dye following Larone's guide.⁽¹⁷⁾

Interpretation for microbial indoor air quality

Following the recommended guideline of the American Conference of Governmental Industrial Hygienists (ACGIH)⁽¹⁸⁾ and the guidelines for good indoor air quality in office premises, Ministry of the Environment, Singapore (1996)⁽¹⁴⁾, if the microbial count was more than 500 cfu/m³, it was an indication of overcrowding or poor ventilation. However, for general offices, bacterial counts or fungal counts should be less than 300 cfu/m³.^(4,5)

Statistical analysis

Statistical analysis was performed using the Statistical Package for the Social Sciences (SPSS) version 22.0. Comparison of means between 2 groups was employed using student's unpaired *t* - test. *P* < 0.05 was considered for statistically significant difference.

Results

Two-time assessments on microbial indoor air quality were performed in a nursing student dormitory including 8 floors with 95 rooms to investigate bacterial and fungal counts. In each assessment, 2 points of air samples were collected from each dormitory room in

the morning (190 samples for bacterial count and 190 samples for fungal count). The first-time assessment found that the mean ± standard deviation (SD) of bacterial counts in dormitory rooms was 151 ± 109 cfu/m³ (outdoor air samples = 175 ± 75 cfu/m³). The highest mean level was found at Floor 1st (257 ± 111 cfu/m³) and the lowest level was found at Floor 7th (85 ± 66 cfu/m³). The second-time assessment found that the mean ± SD of bacterial counts in dormitory rooms was 158 ± 92 cfu/m³ (outdoor air samples = 162 ± 53 cfu/m³). The highest mean level was found at Floor 1st (239 ± 160 cfu/m³) and the lowest level was found at Floor 6th (112 ± 35 cfu/m³). Details are shown in Table 1. As for fungal count, the first-time assessment found that the mean ± SD was 374 ± 273 cfu/m³ (outdoor air samples = 409 ± 136 cfu/m³). The highest mean level was found at Floor 2nd (674 ± 333 cfu/m³) and the lowest level was found at Floor 6th (135 ± 66 cfu/m³). The second-time assessment found that the mean of fungal counts in dormitory rooms was 363 ± 258 cfu/m³ (outdoor air samples = 364 ± 153 cfu/m³). The highest mean level was found at floor 2nd (695 ± 317 cfu/m³) and the lowest level was found at floor 8th (126 ± 27 cfu/m³), as shown in Table 1.

Table 1. Means ± SD of bacterial and fungal counts (cfu/m³) in air samples collected from the dormitory rooms (the first and the second round assessment).

Floors	The first round		The second round	
	Bacterial counts (mean ± SD)	Fungal counts (mean ± SD)	Bacterial counts (mean ± SD)	Fungal counts (mean ± SD)
Floor 1 (n = 10)	257 ± 111 (88 - 442)	210 ± 106 (62 - 380)	239 ± 160 (80 - 504)	199 ± 63 (133 - 309)
Floor 2 (n = 26)	112 ± 56 (27 - 265)	674 ± 333 (88 - 1228)	170 ± 112 (88 - 530)	694 ± 317 (194 - 1,237)
Floor 3 (n = 32)	214 ± 138 (35 - 689)	293 ± 179 (62 - 769)	222 ± 102 (50 - 477)	224 ± 105 (72 - 592)
Floor 4 (n = 28)	183 ± 132 (53 - 574)	595 ± 267 (256 - 1,060)	124 ± 41 (88 - 212)	601 ± 266 (265 - 1,067)
Floor 5 (n = 28)	136 ± 68 (44 - 327)	340 ± 202 (71 - 936)	122 ± 70 (88 - 459)	255 ± 86 (88 - 468)
Floor 6 (n = 28)	100 ± 65 (18 - 274)	134 ± 66 (18 - 292)	112 ± 34 (62 - 212)	256 ± 79 (115 - 424)
Floor 7 (n = 28)	85 ± 66 (17 - 283)	383 ± 147 (177 - 839)	168 ± 86 (88 - 530)	343 ± 193 (106 - 848)
Floor 8 (n = 10)	210 ± 107 (62 - 380)	141 ± 79 (44 - 283)	148 ± 46 (97 - 239)	126 ± 27 (88 - 168)
Total (n = 190)	151 ± 109 (17 - 689)	374 ± 273 (18 - 1,228)	158 ± 92 (50 - 530)	363 ± 257 (72 - 1,237)
Outdoor (n = 24)	175 ± 74 (80 - 345)	409 ± 136 (88 - 553)	162 ± 53 (97 - 292)	364 ± 153 (88 - 675)

From both assessments, it was found that the lower floors of the dormitory (1 - 4 floors) had significantly higher levels of microbial counts when compared with the higher floors of the dormitory (5 - 8 floors) ($P < 0.001$). Details are shown in Table 2. As well as, the percentages of bacterial and fungal counts with > 300 cfu/m³ in the lower floors (1 - 4 floors) were significantly higher than those in the higher floors (5 - 8 floors) in the first and the second assessment ($P < 0.05$ and $P < 0.001$, respectively) (Table 3). However, the percentages of only fungal counts with > 500 cfu/m³ (higher the recommended indoor air guideline of American Conference of Governmental Industrial Hygienists, ACGIH) in the

lower floors (1 - 4 floors) were significantly higher than those in the higher floors (5 - 8 floors) ($P < 0.05$) (Table 3).

Isolated bacteria were preliminarily identified and found that 64.5% and 22.1% were *Staphylococcus spp.* and *Streptococcus spp.*, respectively. Whereas, 62.5% of *Staphylococcus spp.* and 12.5% of *Streptococcus spp.* were found in outdoor air samples. Isolated fungi were found 62.1% of *Aspergillus spp.*, and 21.3% of *Penicillium spp.*, respectively. Moreover, 58.3% of *Aspergillus spp.* and 20.8% of *Penicillium spp.* were found in outdoor air samples. Details are shown in Table 4 and Table 5.

Table 2. Comparison of bacterial and fungal counts (means \pm SD) between dormitory rooms in lower floors and dormitory rooms in higher floors (the first and the second round assessment).

Floors	The first round		The second round	
	Bacterial counts (n = 190) (mean \pm SD)	Fungal counts (n = 190) (mean \pm SD)	Bacterial counts (n = 190) (mean \pm SD)	Fungal counts (n = 190) (mean \pm SD)
1 - 4 (n = 96)	183 \pm 92	469 \pm 189	179 \pm 82	448 \pm 194
5 - 8 (n = 94)	120 \pm 67	274 \pm 129	137 \pm 58	271 \pm 128
P - value from t-test	<0.001	<0.001	<0.001	<0.001

Table 3. Comparison of number and percentage of bacterial and fungal counts with >300 cfu/m³ and >500 cfu/m³ between dormitory rooms in lower floors and dormitory rooms in higher floors (the first and the second round assessment).

Floors	The first round		The second round	
	Bacterial counts (n = 190) > 300 cfu/m ³	Fungal counts (n = 190) > 300 cfu/m ³	Bacterial counts (n = 190) > 300 cfu/m ³	Fungal counts (n = 190) > 300 cfu/m ³
1 - 4 (n = 96)	24 (25.0%)	63 (65.6%)	22 (22.9%)	55 (57.3%)
5 - 8 (n = 94)	6 (6.4%)	33 (35.1%)	4 (4.3%)	28 (29.8%)
P - value	< 0.05*	< 0.001*	< 0.05*	< 0.001*
Floors	> 500 cfu/m ³		> 500 cfu/m ³	
1 - 4 (n = 96)	3 (3.1%)	33 (34.4%)	2 (2.1%)	32 (33.3%)
5 - 8 (n = 94)	0 (0.0%)	7 (7.4%)	0 (0.0%)	3 (3.2%)
p-value	NC	< 0.05*	NC	< 0.05*

*Statistical significance at $\alpha = 0.05$ by Proportional Z test
 NC = Not calculation

Table 4. Number and percentage of isolated bacterial group in air samples collected from the dormitory room (n = 428 colonies).

Air sampling site	No (%) of Bacterial groups				
	<i>Staphylococcus spp.</i>	<i>Streptococcus spp.</i>	<i>Micrococcus spp.</i>	<i>Bacillus spp.</i>	Gram-negative rods
Floor 1 (n = 20)	10 (50.0)	5 (25.0)	3 (15.0)	2 (10.0)	0 (0.0)
Floor 2 (n = 52)	35 (67.3)	10 (19.3)	5 (9.6)	2 (3.8)	0 (0.0)
Floor 3 (n = 64)	42 (65.6)	16 (25.0)	3 (4.7)	2 (3.1)	1 (1.6)
Floor 4 (n = 56)	40 (71.4)	11 (19.6)	2 (3.6)	3 (5.4)	0 (0.0)
Floor 5 (n = 56)	30 (53.6)	16 (28.6)	5 (8.9)	3 (5.4)	2 (3.5)
Floor 6 (n = 56)	42 (75.0)	11 (19.6)	2 (3.6)	1 (1.8)	0 (0.0)
Floor 7 (n = 56)	36 (64.3)	12 (21.4)	5 (8.9)	2 (3.6)	1 (1.8)
Floor 8 (n = 20)	10 (50.0)	3 (15.0)	2 (10.0)	3 (15.0)	2 (10.0)
Total (n = 380)	245 (64.5)	84 (22.1)	27 (7.1)	18 (4.7)	6 (1.6)
Outdoor (n = 48)	30 (62.5)	6 (12.5)	6 (12.5)	4 (8.3)	2 (4.2)

Table 5. Number and percentage of isolated fungal group in air samples collected from the dormitory room (n= 362 colonies).

Air sampling site	No (%) of fungal culture			
	<i>Aspergillus spp.</i>	<i>Penicillium spp.</i>	<i>Fusarium spp.</i>	Others*
Floor 1 (n = 16)	5 (31.2)	4 (25.0)	4 (25.0)	3 (18.8)
Floor 2 (n = 40)	25 (62.5)	10 (25.0)	5 (12.5)	0 (0.0)
Floor 3 (n = 56)	35 (62.5)	12 (21.4)	6 (10.7)	3 (5.4)
Floor 4 (n = 50)	30 (60.0)	10 (20.0)	8 (16.0)	2 (4.0)
Floor 5 (n = 50)	34 (68.0)	7 (14.0)	6 (12.0)	3 (6.0)
Floor 6 (n = 46)	30 (65.2)	9 (19.6)	5 (10.9)	2 (4.3)
Floor 7 (n = 46)	32 (69.6)	11 (23.9)	3 (6.5)	0 (0.0)
Floor 8 (n = 10)	4 (40.0)	4 (40.0)	2 (20.0)	0 (0.0)
Total (n = 314)	195 (62.1)	67 (21.3)	39 (12.4)	13 (4.2)
Outdoor (n = 48)	28 (58.3)	10 (20.8)	7 (14.6)	3 (6.3)

* Such as *Cladosporium spp.*, *Alternaria spp.*, and *Rhizopus spp.*

Discussion

Indoor air quality is one of the most significant factors affecting the health and well-being of individuals who spend more than 90% of their lives indoors.^(1,3) According to the World Health Organization (WHO) roughly 3 billion people around the world are suffering from diseases caused by indoor air pollution; dormitory room is a kind of residential environment which, may be more polluted from the crowded dormitory environments.⁽²⁾ In dorms, a low ventilation rate is a risk factor for asthma, allergy and adverse health.^(2,9) However, a part of students who were susceptible for illnesses related to indoor air quality might be associated with physiological and personal health condition, personal hygiene and environmental factors.⁽¹⁹⁾ Dormitory is one of important indoor places for students in a university and its environments probably affect student's health. Ventilation systems and indoor air quality (IAQ) are also affecting health of students.⁽⁹⁾ Many studies have shown that indoor air pollutants, especially in the hospital were higher than outdoor air pollutants and that occurrence, concentration, and duration of indoor volatile organic compounds exposure in residences may contribute to the occupant's short and long-term adverse health effects.^(11,12,20) Previous studies in China reported that dormitory environment factors, such as dampness and poor ventilation were related to asthma and respiratory infections among college students.^(8,9)

This short-time follow-up study of microbial indoor air quality in a nursing institute dormitory was conducted 2 times during 3-month observation due to the students' complaints before the study was done. Indoor air samples were collected from dormitory rooms using BioStage Impactor (QuickTake 30 sample pump) to assess total bacterial and fungal counts. After that, the isolated bacteria and fungi were preliminarily identified group or genus by Gram's staining and lactophenol cotton blue. Results showed that the higher mean levels of bacteria and fungi were found at the lower floors (especially, floor 1 or floor 2) and the lower levels of bacterial and/or fungal counts were found at the higher floors (floor 6 or over) in the both assessments. It might be due to the lower floors, especially in the first and the second floors, to be inadequate ventilation and the effects of surrounding environment air. Data from observation and interviews showed that most of dorm rooms in the lower floors, especially, floor 1 and floor 2 did not regularly open the windows due to the smell from the surrounding

environment. Additionally, the common bathroom facilities were regularly wet and the exhaust fans were out of order. The higher relative humidity supported the growth of microbial organisms.⁽⁷⁾ Whereas, most of dorm rooms in the higher floors regularly opened the windows and the common bathroom facilities were regularly dry due to the exhaust fans with well-work. Moreover, in the lower floors, each dorm room was shared by 4 - 5 students with crowded space and poor room hygiene practices tended to have higher bacterial and fungal counts than other higher floors of dormitory with lower density of students (2 - 3 students in each room). A previous study showed the lower ventilation in studied rooms, perceived the higher stuffy smell.⁽⁹⁾ Another study in China explained that cleaning routines and crowded spaces may be important factors for the propagation of respiratory infection in students.⁽⁸⁾ Many reviews have shown that dampness with a low ventilation rate in the building is strongly associated to health problems like asthma and respiratory symptoms due to higher fungal contamination in indoor air.⁽⁷⁻⁹⁾

The isolated bacterial and fungal from air samples were identified by group or genus; it was found that the most common bacteria were *Staphylococcus spp.* and *Streptococcus spp.*, and the most common fungi were *Aspergillus spp.*, and *Penicillium spp.* Although, these isolated micro-organisms do not generally present a health hazard, high loads of bacteria and fungi suggested overcrowding and poor air hygiene. They may trigger allergic reactions, such as, allergic rhinitis and cough, allergic skin problems, and non-specific symptoms particularly among children, elderly and immune-compromised hosts.⁽⁴⁻⁶⁾ The fact that these groups are normally present in air, environments and on human skin.⁽²¹⁾ They can survive for a long period in air and the environment. A previous study demonstrated that most *Staphylococcus spp.* found in air and the environment was *S.epidermidis*, the normal flora of the human skin and respiratory tract.⁽²²⁾ This present study of airborne isolated fungi was similar to the study in an academic dormitory in several studies^(9,22,23) which found that *Aspergillus spp.* was the most common fungi. This fungal genus could compromise the health and well-being of humans.⁽²²⁾ Mold or fungi can live practically anywhere and have particularly favorable growth condition inside residential houses and dormitory rooms.

The preventive measures to reduce the probable effects from poor indoor air quality in the dormitory should be considered. Increase the amount of air ventilation in dormitory rooms, especially dormitory rooms at lower floor (1 - 4) should be done, probably by opening the window or exhaust fans. Minimizing moisture accumulation or protect stored materials from moisture in the room, such as waste paper. Indoor environment should be sufficiently cleaned, especially dorm room cleaning. Additionally, the environmental surveillance, especially biological and chemical hazards contaminated in outdoor air of the dormitory surroundings should be done.

Conclusion

This study reveals that the lower floors of the dormitory had significantly higher levels of bacterial and fungal counts when compared with the higher floors of the dormitory from both assessments. Most isolated bacteria and fungi do not generally present a health hazard, but high loads may trigger allergic reactions in susceptible hosts.

Acknowledgments

Researchers would like to thank heads of studied nursing student dormitory for her help and cooperate during this study.

Conflict of interest

The authors, hereby, declare no conflict of interest.

References

1. World Health Organization. Preventing disease through healthy environments: Towards an estimate of environmental burden of disease. Geneva: WHO Press; 2006.
2. Turunen M, Putus T, Nevalainen A, Shaughnessy R, Haverinen-Shaughnessy U. Indoor environmental quality in school buildings, and the health and well-being of students. *Inter J Hygiene Environ Health* 2014;217:733-9.
3. Reijula K. Buildings with moisture problems—a new challenge to occupational health care. *Scand J Work Environ Health* 1996;22:1-3.
4. World Health Organization. Indoor air quality: Biological contaminants. WHO regional publications. European Ser 1990;30:385-74.
5. Zweers T. Health and indoor climates complaints of 7043 office workers in 61 buildings in the Netherlands. *Indoor Air* 1992;2:127-36.
6. Gorny RL, Dutkiewicz J. Bacterial and fungal aerosols in indoor environment in Central and Eastern European countries. *Ann Agric Environ Med* 2002;9:17-23.
7. World Health Organization. Development of WHO guidelines for indoor air quality: Dampness and mould. Report on a working group meeting. Bonn, Germany 17-18 October 2007. WHO Regional Office for Europe; 2008.
8. Sun Y, Zhang Y, Sundell J, Fan Z, Bao L. Dampness at dorm and its association with allergy and airways infection among college students in China: a cross-sectional study. *Indoor Air* 2009;19:174-82.
9. Sun Y, Zhang Y, Bao L, Fan Z, Sundell J. Ventilation and dampness in dorms and their associations with allergy among college students in China: a case-control study. *Indoor Air* 2011;21:277-83.
10. Reynolds MG, Huy B, Thu VH, Montgomery JM, Bausch DG, Shah JJ, et al. Factors associated with nosocomial SARS-CoV transmission among healthcare workers in Hanoi, Vietnam. *BMC Public Health* 2006; 6:207.
11. Siegel JD, Rhinehart E, Jackson M, Chiarello L. Healthcare infection control transmission of infectious agents in healthcare settings. *Am J Infect Control* 2007;35(10 Suppl 2):S65-164.
12. Yau YH, Chandrasegaran D, Badarudin A. The ventilation of multiple-bed hospital wards in the tropics: A review. *Build Environ* 2011;46:1125-32.
13. Pentamwa P, Jongprasom T, Chungchom N, Prompol B. Environmental health assessment in university dormitories. *KKU J Public Health Res* 2012;5:35-44.
14. Institute of Environmental Epidemiology, Ministry of the Environment. Guidelines for Good Indoor Air Quality in Office Premises. 1st ed. Ministry of the Environment, Singapore, 1996:1-47.
15. Pasquarella C, Albertini R, Dall'aglio P, Saccani E, Sansebastiano GE, Signorelli C. Air microbial sampling: the state of the art. *Ig Sanita Pubbl* 2008;64:79-120.
16. Luksamijarulkul P, Pipitsangjan S. Microbial air quality and bacterial surface contamination in ambulances during patient services. *Oman Med J* 2015;30:104-10.
17. Larone DH. Medically important fungi: a guide to identification. 3rd ed. Washington DC: American Society of Microbiology Press; 1995.
18. Seitz TA. NIOSH indoor air quality investigations 1971-1988. In: Weekes DM, Gammage RB, editors. Proceedings of the indoor air quality, international symposium: The practitioner's approach to indoor air quality investigations. American Industrial Hygiene Association Ohio 1989:163-71.

19. Sun Y, Zhang Y, Bao L, Fan Z, Wang D, Sundell J. Effects of gender and dormitory environment on sick building syndrome symptoms among college students in Tianjin, China. *Build Environ* 2013;68:134-9.
20. El Sharkawy MF, Noweir ME. Indoor air quality levels in a university hospital in the eastern province of Saudi Arabia. *J Family Community Med* 2014;21:39-47.
21. Sun Y, Sundell J, Zhang Y. Validity of building characteristics and dorm dampness obtained in a self-administrated questionnaire. *Sci Total Environ* 2007;387:276-82.
22. Sun Y, Wang Z, Zhang Y, Sundell J. In China, students in crowded dormitories with a low ventilation rate have more common colds: evidence for airborne transmission. *PLoS ONE* 2011; 6: 271-340.
23. Luksamijarulkul P, Aiempradit N, Vatanasomboon P. Microbial contamination on used surgical masks among hospital personnel and microbial air quality in their working wards: A hospital in Bangkok. *Oman Med J* 2014;29:249-53.