

Original article

Lung function and exercise performance assessment after inhalation of naturally essential oils: A descriptive observational study

Hamzeh Al Zabadi^{a*}, Azmi Hussein^b, Jamal Shaker^c, Belal Rahhal^d, Nidal Amin Jaradat^e,
Malik Al Qub^b

^aPublic Health Department, Faculty of Medicine and Health Sciences, An-Najah National University, P.O. Box 7, Nablus, Palestine

^bBiochemistry, Molecular Biology and Genetics Department, Faculty of Medicine and Health Sciences, An-Najah National University, P.O. Box 7, Nablus, Palestine

^cFaculty of Physical Education, An-Najah National University, P.O. Box 7, Nablus, Palestine

^dPhysiology, Pharmacology and Toxicology Department, Faculty of Medicine and Health Sciences, An-Najah National University, P.O. Box 7, Nablus, Palestine

^eDepartment of Pharmacy, Faculty of Medicine and Health Sciences, An-Najah National University, P.O. Box 7, Nablus, Palestine

Background: Essential oils have gained massive attention. We aimed to assess the effects of *Clinopodium serpyllifolium* and *Cymbopogon citratus* volatile oils inhalation on exercise performance and lung function.

Methods: Twenty physical education students were randomly assigned into two groups: *Clinopodium serpyllifolium* and *Cymbopogon citratus* (ten participants each). One group was nebulized by *Clinopodium serpyllifolium* oil and the other by *Cymbopogon citratus* oil in a concentration of (0.02 ml/kg of body mass) which was mixed with 2 ml of normal saline before 800 m running tests. Lung function tests were measured using a spirometer pre and post nebulization.

Results: There was significant increase in Forced Expiratory Volume in the first second and Forced Vital Capacity after inhalation of both oils. The normal spirometry after inhalation of *C. serpyllifolium* oil increased by 30%. A sharp decrease in mild spirometry (50% before to 10% after inhalation of the *C. citratus* oil) was observed. Significant reductions in the means of the running time were observed.

Conclusion: Our findings support the effectiveness of *C. serpyllifolium* and *C. citratus* essential oils on exercise performance and respiratory function parameters.

Keywords: *Clinopodium serpyllifolium*, *Cymbopogon citratus*, essential oil, lung function, athletic performance.

The Plant Kingdom is considered an essential part of human being life and it is impossible to live without them. Recently huge number of researchers are focusing on the phytochemical products due to their importance for solving many pharmacological problems and for investigation of new drugs and other none therapeutic applications.^(1,2) Volatile oils (essential

oils) are odorous principles with aromatic odor and most of them are phytochemical compounds that evaporated at ordinary temperature and have been used since ancient times in manufacturing of perfumes, flavoring agents, food preservatives and medicament.^(3, 4) They are considered secondary metabolic compounds which are biosynthesized in specialized structures for various purposes such as pollination, protection, communication and signaling between other plants. They have been used pharmaceutically for various aims such as local analgesics (eugenol), decongestants (eucalyptol), expectorants (carvacrol), antiseptics (menthol), anthelmintic (ascaridol) and many of them used as

*Correspondence to: Hamzeh Al Zabadi, Public Health Department, Faculty of Medicine and Health Sciences, An-Najah National University, P.O. Box 7, Nablus, Palestine.

E-mail: halzabadi@gmail.com

Received: April 9, 2019

Revised:

Accepted:

flavoring agents in the pharmaceutical industry to get rid of unpleasant odors and tastes of drugs.^(5, 6) Many of the volatile oils have been used in the folk medicine for treatment and prevention of various upper respiratory system diseases such as *Clinopodium serpyllifolium* and *Cymbopogon citratus* volatile oils.^(7, 8) *Clinopodium serpyllifolium* subsp. *fruticosum* (L.) Bräuchler, is a synonym name for *Micromeria fruticosa* (L.) Druce which is a member of the Lamiaceae family and commonly known as White leafed Savory or White Hyssop or White micromeria. It is a perennial herbaceous plant which endemic to Palestine, Syria, Lebanon, Jordan and Turkey and has grey, egg shaped, opposite leaves and small white flowers on its stems.^(9, 10) Its volatile oil contents have a mixture of monoterpene (+)-pulegone, isomenthol, isomenthone, menthol, pinene and limonene, piperitone, piperitenone oxide as well as it contains sesquiterpenoids such as caryophyllene and germacrene-D.⁽¹¹⁾ The aerial parts of the plants are widely used in various Mediterranean regions as infusions and decoctions for treatment of various diseases including gastrointestinal spasms, diarrhea, ophthalmic inflammations, cardiac disorders, hypertension, open wounds, colds and other respiratory system inflammations.^(12, 13) It has been approved that these plant volatile oils exhibited activity against 14 bacteria, three fungi and a yeast and its methanolic extracts exhibited significant antioxidant activity in the DPPH, 1,1-Diphenyl-2-picryl-hydrazyl and inhibiting linoleic acid oxidation assays.⁽¹⁴⁾ *Cymbopogon citratus* (DC.) Stapf, commonly known as lemon grass or citronella grass belongs to the Poaceae family which is a perennial herbaceous plant widely distributed throughout the world especially in the semi-tropical and tropical regions. Its decoction used as antibacterial, antifungal, antipyretic, stomachic, carminative and sedative. Various studies have approved its sedative, CNS depressant, analgesic, antibacterial and antifungal activities.^(15 - 19) In the light of this, the present study looks at two species used in folk medicine; *Clinopodium serpyllifolium* and *Cymbopogon citratus* volatile oils inhalation in two different groups of male athlete students to assess their exercise performance and lung function using a spirometric pre- and post test assessment design.

Materials and methods

Instrumentations

During this study, the following instruments were used: ultrasonic-microwave cooperative extractor/

reactor (CW-2000, China), balance (Radwag, AS 220/c/2, Poland), Philips respironics inhalators nebulizers (75644321, China) and Care-fusion spirometer (ME44QY 08563848, UK)

Collection and preparing plant materials

Clinopodium serpyllifolium and *Cymbopogon citratus* plants were collected during its flowering time from Jerusalem region (Palestine) in April, 2016. Botanical identification was carried out by Pharmacognosist Dr. Nidal Jaradat from the Pharmacognosy and Herbal Products Laboratory, Faculty of Medicine and Health Sciences, An-Najah National University, Nablus. The identification process was conducted using live herbal specimens and photographs from books. Voucher specimens were deposited in the Pharmacognosy and Herbal Products Laboratory under the code numbers: Pharm-PCT-1575 for *C. serpyllifolium* and was Pharm-PCT-2791 for *C. citratus*. To extract volatile oil, the aerial parts of *C. serpyllifolium* and *C. citratus* were separated carefully and then washed twice with distilled water.

Essential oils extraction

The essential oil of *C. serpyllifolium* was extracted using a microwave oven as described by Jaradat, 2016 with some modifications.⁽¹⁵⁾ The power of the microwave oven was set at 1000 W. Clevenger apparatus with a 1L round-bottom flask containing 100g of *C. serpyllifolium* was placed inside the microwave oven. About 500 ml distilled water was then added into the flask containing the powder. The flask was then connected to Clevenger apparatus. Microwave distillation was carried out three times for 15 min each at 100°C. The obtained volatile oil was collected into a clean beaker and chemically dried then the purified essential oil was weighed and stored in tightly-closed amber-colored bottles at 4°C in the refrigerator. The same procedure was repeated for extraction of *C. citratus* essential oil.

Subjects and study design

Twenty male university students from the Faculty of Physical Education, An-Najah National University in Nablus-Palestine volunteered and were randomly assigned into two different groups (10 subjects each) to take part in the experiments using a non-randomized quasi-experimental uncontrolled before-and-after study design for each group. The study was single blind at the level of participants. One group (10 students) was nebulized with *C. serpyllifolium* oil, while the other

group (10 students) was nebulized with *C. citratus* oil (0.02 ml/kg of body mass of each oil) mixed with 2ml of normal saline 5 min before a 800 m running test according to the method validated by Spencer and Gastin, 2001.⁽¹⁶⁾ Traditionally, it was thought that in the middle distance track running the relative energy system was 75% aerobic and 22% anaerobic and a 2 - 3% of the energy coming from the creatine-phosphate system. But recently, the aerobic energy system contributes significantly to the energy supply during middle distance running and long sprint event. Moreover, the relative anaerobic and aerobic system contributions of the four events (200 m, 400 m, 800 m, and 1500 m). In the middle distance running such as 800 m event depends on the aerobic 84% and anaerobic energy system which is greater than it was traditionally thought, so during this event the need of oxygen increases with the event duration from 200 to 1500 m running race.⁽¹⁶⁾ Moreover, the 800 m distance was required to ensure the duration of effect of the oils (if any) as inhaled route of administration and the greater distances could have been taken into consideration if the interventions were given orally or by another route of administration, however, our estimation was that this distance could be representable of such route of administration given at the same time that the same group conducted the same running pre and post intervention. A washout period of three days (from Sunday to Wednesday) between pre and after test for each group was given to rule out any fatigue or significant impact on our results pre and post test. Lung function tests were measured using a spirometer for each student pre and post nebulization. In each stage, two measurements were assessed and the average was taken. The lung function tests were assessed 3 min after inhalation for both Nacl and the oil. The study was carried out in October, 2016. All subjects we first aligned at a straight horizontal line as the same starting point for all and at the end of the 800 m exactly another straight horizontal line was drawn. The running moment was given to all at the same point in time and when the subject arrive to the end exactly his exact running time in seconds was calculated depending on the starting time. This running procedure was adapted for all subjects at the same time.

Methodology

The subjects were familiarized with the spirometry. Lung function tests (LFTs) were measured using a spirometer for each student pre and post

inhalations with *C. serpyllifolium* and *C. citratus* oils. The spirometer is used to differentiate between obstructive and restrictive diseases and assess the degree of associated changes.^(17, 18) Such parameters include Forced Expiratory Volume in the first second (FEV1) and Forced Vital Capacity (FVC). The specificity and sensitivity of spirometer in diagnosis of obstructive lung disease are reported as 84% and 92%, respectively.⁽¹⁹⁾ While in the diagnosis of restrictive lung disease, it has a sensitivity and specificity of 42.2% and 94.3%, respectively.⁽²⁰⁾ FEV1 is the maximum air volume exhaled with maximal effort in the first second from a position of full inspiration. This value declines less severely with restrictive diseases than obstructive diseases. FVC, on the other hand, is the maximal air volume exhaled with maximal effort from a position of full inhalation.⁽²¹⁾ and is reduced by an airflow obstruction and ventilation restriction, which results from lung-exterior factors such as skeletal pains or intrinsic lung disease, especially restrictive ones.⁽²²⁾ In the latter, there is a decline in the lung compliance associated with the presence of partial or diffuse lung fibrosis. These fibrotic changes render the lung smaller and stiffer, leading to a decrease in the FVC. The FEV1/FVC ratio is reduced in obstructive patterns, but it is normal or even increased in restrictive patterns as both nominator and denominator proportionally change.

The classification criteria for normal, mild and moderate restriction were based on the American thorax society depending on the FEV1 value. Therefore, the severity of any spirometric abnormality was based on the FEV1 (%predicted) as follows: mild >70%, moderate 60-69%, moderately severe 50-59%, severe 35-49% and very severe <35.

Data collection form

Different sections were included. The first was the demographic section, which contained questions regarding age, gender, education level, health history, smoking status and body mass index. Then, spirometry was performed using a spirometer apparatus. Regular guidelines for spirometer testing were followed.⁽²³⁾ The subjects were seated during the test, with the nose clipped to prevent air leakage through the nares. Three attempts for each subject were allowed, and the best spirograph was selected automatically by the spirometer. Forced spirometry was measured for each subject. As for each subject, FEV1, FVC, FEV1/FVC ratios were assessed.

Ethical considerations

The study aims, protocols and the informed consent forms were approved by the Institutional Review Board (IRB) at An-Najah National University (IRB archived number: 11/August/2016). The study was conducted in accordance with the requirements of the declarations of Helsinki. All subjects gave their signed informed consent before they entered in the study. Subjects were not offered any incentives and they were able to withdraw from the study at any time. Data obtained were kept confidential. Subjects were assured that refusal to participate in the study would not affect their future healthcare carrier in any way. Two trained research assistants conducted the face to face interviews with the subjects.

Data analysis

All data were entered and analyzed using the Statistical Package for the Social Sciences (SPSS) version 16. The main study outcomes were tested for normality using Shapiro-Wilk significant test and found not to be normally distributed. Qualitative variables have been expressed as frequency tables and bar charts. The non-parametric Two-related samples Wilcoxon test was used to test for the differences in the means between pre- and post-inhalers of the continuous outcomes. *P* - value of less than 0.05 was considered statistically significant.

Results

Socio-demographic characteristics

All subjects were single. The mean age (standard

deviation; SD) of *C. serpyllifolium* group subjects was 19.20 (1.23) years; 30% of them smokers, 1-10 cigarettes per day. In the *C. citratus* group, however, the mean age (SD) was 20.70 (1.25) years and the majority (60%) were smokers 1-10 cigarettes per day. Table 1 shows the socio-demographic characteristics details of the studied subjects in each group.

Respiratory parameters assessment

A lung function test showed an improvement on the lung status for the students after their inhalation of the oils. Figure 1 shows that the normal spirometry results were 30%, while after inhalation with *C. serpyllifolium* oil the ratio were 60%. The mild spirometry was dropped from 70% before to 40% after inhalation of the *C. serpyllifolium* oil. On the other hand, there was also an increase in normal lung status of the students who inhaled *C. citratus* oil (40% before and 60% after). There was also a sharp decrease in mild spirometry (50% before inhalation to 10% after inhalation of the *C. citratus* oil (Figure 2). Interestingly, Table 2 shows that there is a statistically significant increase in FEV1 and FVC in the post-test students after their inhalation of the *C. serpyllifolium* and *C. citratus* oils. Regarding the FEV1/FVC ratios, there was a strong significant decrease toward the normal in *C. serpyllifolium* oil participants but only a weak and borderline decrease (not significant) toward the normal in *C. citratus* oil participants.

Table 1. Socio-demographic characteristics of the study population given *C. serpyllifolium* and *C. citratus* oils (N=10 for each group).

Variables	<i>C. serpyllifolium</i> oil group	<i>C. citratus</i> oil group
	n (%)*	
Gender		
-Male	10(100)	10(100)
Marital status		
-Single	10(100)	10(100)
Smoking status (cigarette/day)		
-Non-smokers	4 (40)	4 (40)
-1-10	3(30)	6 (60)
-11-20	3(30)	0 (0)
Age (year)	19.20 ± 1.23**	20.70 ± 1.25**
Weight (kg)	72.80 ± 5.88**	71.90 ± 9.07**
Height (meter; m)	1.79 ± 0.036**	1.74 ± 0.044**
BMI (kg/m²)	22.59 ± 2.08**	23.40 ± 2.12**

*Data is presented as frequency (percent) (N=10). ** Mean ± standard deviation.




Figure 1. Respiratory parameters for participants before and after inhalation of *C. serpyllifolium* oil (as percent)




Figure 2. Respiratory parameters for participants before and after oil inhalation of *C. citratus* oil (as percent)

Table 2. Respiratory parameters in the post- and pre-inhalers measurements for the study outcomes of the participants given *C. serpyllifolium* (n=10) and *C. serpyllifolium* oils (n=10) when running equal distance of 800m before and after.

Outcome Variables	<i>C. serpyllifolium</i> oil group			<i>C. citratus</i> oil group		
	Mean \pm SD*	Mean difference \pm SD	P - value [§]	Mean \pm SD*	Mean difference \pm SD	P - value [§]
FEV1						
- Post-inhaler	4.76 \pm 0.24		0.005 [!]	4.62 \pm 0.22		0.009 [!]
- Pre-inhalers	4.06 \pm 0.57	0.70 \pm 0.69		3.89 \pm 0.64	0.73 \pm 0.57	
FVC						
- Post-inhaler	5.64 \pm 0.31		0.005 [!]	5.54 \pm 0.29		0.005 [!]
- Pre-inhalers	4.29 \pm 0.63	1.35 \pm 0.75		4.37 \pm 0.47	1.17 \pm 0.38	
FEV1/FVC						
- Post-inhaler	84.00 \pm 1.88		0.005 [!]	83.90 \pm 0.32		0.092
- Pre-inhalers	96.90 \pm 8.90	-12.90 \pm 10.16		89.80 \pm 10.83	-5.90 \pm 10.91	

*SD, standard deviation; FEV1, Forced Expiratory Volume in the first second; FVC, Forced Vital Capacity. FEV1/FVC, Ratio of FEV1 over FVC. §P - value for the mean differences of the Two-related samples Wilcoxon test. [!]Significant P - values.

Running performance

There was a statistically significant reduction in the mean running time after inhalation of *C. serpyllifolium* oil and *C. citratus* oil among the study participants (P - values 0.005 and 0.025; respectively) as shown in Table 3.

Discussion

Our findings showed that the essential oils isolated from *C. serpyllifolium* and *C. citratus* enhanced athletic performance and lung function. The results showed that there is a significant increase in FEV1 and FVC in the post-test students after their inhalation of the *C. serpyllifolium* and *C. citratus* oils. Moreover, the running times for each group were significantly decreased in both *C. serpyllifolium* and *C. citratus* nebulized groups (P-values, 0.005 and 0.025; respectively). As previously mentioned, natural plants essential oils have been traditionally used in the

treatment of various physiological and psychological disorders.⁽²⁴⁾ Most of the essential oils are considered to be safe and their safety had been monitored in different ways, and they have been used from the ancient times in perfumery, cosmetics and food industries.⁽²⁵⁾ Most of these studies focused on *peppermint plant* but unfortunately and according to the best of authors' knowledge there were no previous studies about *C. serpyllifolium* and *C. citratus* oils inhalations on the lung function and on the athletic performance. We have found no previous studies were conducted regarding *C. citratus* in relation to exercise and/or lung function performance. In the meanwhile, all previous studies conducted on these two plants were in-vitro tested and even though (to the best of our knowledge) no single previous study tested the oils of these plants against exercise and lung functions. However, *C. citratus* has shown an effect against dental caries⁽²⁶⁾, anti-malarial effect⁽²⁷⁾, anti-bacterial

Table 3. Changes in exercise performance (time in seconds) for the participants given *C. serpyllifolium* (n=10) and *C. citratus* oils (n=10) when running equal distance of 800 m before and after oil inhalations.

Outcome Variables	<i>C. serpyllifolium</i> oil group			<i>C. citratus</i> oil group		
	Mean \pm SD*	Mean difference \pm SD	P - value [§]	Mean \pm SD*	Mean difference \pm SD	P - value [§]
Time (sec)**						
- Post-inhaler	188.40 \pm 13.29			171.90 \pm 14.36		
- Pre-inhalers	206.60 \pm 13.79	-18.20 \pm 13.60	0.005 [!]	179.00 \pm 16.15	-7.10 \pm 7.75	0.025 [!]

* SD, standard deviation; §P-value for the mean differences of the Two-related samples Wilcoxon test. [!]Significant P-values. **Data in the table is expressed as seconds.

effect.⁽²⁸⁾ Another study showed a positive effect of *C. citratus* leaves on some serum components and the antioxidant status without any beneficial effect on growth performance and carcass traits in broiler chickens.⁽²⁹⁾ Regarding *C. serpyllifolium*, however, results from mice models suggested that the aqueous extract of *Micromeria fruticosa* (*C. serpyllifolium*) has both anti-inflammatory as well as, gastro-protective activities. The authors concluded that this plant could be used as an alternative or supplementary herbal remedy for the treatment of inflammatory diseases especially when combined with strong anti-inflammatory medications that have ulcerogenic side effects such as NSAIDs (Non-steroidal anti-inflammatory drugs).⁽³⁰⁾ Another study found that both the oil and the aqueous extract of *M. fruticosa* exhibited marked anti-tumor activities against Human Colon Tumor cells (HCT) and Mammary Carcinoma F7 (MCF7). The study concluded that extract of *M. fruticosa* has a remarkable inhibitory activity in non-inflammatory reactions as well as inflammatory pain.⁽³¹⁾ The use of essential oils in medicine began in the ancient Egyptian Era, and has continued ever since^(32, 33). One of the most popular parts of complementary and alternative medicine is considered to be aromatherapy, which depends only on essential oils utilization for treatment of various diseases and this branch has spread worldwide, despite the lack of scientific basis for the effectiveness of essential oils.⁽³⁴⁾ On the other hand, the long history of essential oils usages in medicine and pharmacy suggests that they may indeed be effective. The odor of essential oils is believed to be important for their effectiveness in treating various illnesses.⁽³⁵⁾ Future study is needed to demonstrate oral supplementation of *C. serpyllifolium* and *C. citratus* essential oils to improve athletic performance and post-exercise recovery; another future objective would be to clarify their mechanism of action and to explain their physiological and pharmacological effects.

Our study has some limitations. This study is limited by the lack of control group of nebulizer (0.9% NaCl only). There was neither placebo/control experiment within subjects nor randomized (subjects/sequence) placebo control experiment. However, no control groups have been proposed in order to ensure that the intervention reaches the maximum number of students agreed to participate. In addition, the design could also be another source of limitations. However, we are aware that in the design we used, secular

trends or sudden changes make it difficult to attribute observed changes to the intervention. Further limitations of this design could be that the occurrence of over-estimation of the effects of intervention. By using this design, however, we were able to measure performance before and after the introduction of intervention in the same study group and the observed differences in performance were assumed to be due to the intervention. This study design has the advantages in that it is relatively simple to conduct and participants are having same characteristics pre-and post-test and it is superior to observational studies. Furthermore, due to lack of resources and difficulty of circumstances in our country and university we found this the most robust design possible to minimize bias and maximize generalizability. Small sample size might also be a source of limitation that could have affected the study statistical power, we are fully aware of this but we were not able to recruit more due to voluntary participation. Therefore, we found it fairly enough to divide the twenty students ten by ten in each group in order to run this study. We also did not conduct between group changes as we did not aim to perform comparison between the two groups with the different oils intervention but we only aimed to assess the outcome for each intervention in the same group pre-and post-test. As we have indicated above, we used a quasi-experimental uncontrolled before-and-after study design and we did not aim to compare between the two groups; therefore, differences in smoking or any other variations between groups were not taken into consideration in the analysis as we have measured the changes for each intervention separately among the same group pre-and post-test. There was no dietary control in this study. However, we believe that those are informative groups of students who might have same weekly dietary intake level and we asked them to restrict any energy stimulator intakes during the week of the study. Therefore, dietary intake could have minimal effects on our findings.

Conclusions

This is the first study that explored the effect of inhalation of *C. serpyllifolium* and *C. citratus* essential oils on exercise performance and lung function. Our findings support the effectiveness of *C. serpyllifolium* and *C. citratus* essential oils on the exercise performance and respiratory function parameters. However, our conclusion and generalizability of our results should be interpreted

with caution due to small sample size and lack of control groups, randomization or masking. We recommend further investigations to explain the mechanism of actions for these two essential oils on exercise performance and respiratory gas exchange parameters.

Acknowledgements

The authors acknowledge the assistance of the technicians: Mohamad Arar and Linda Esa and An-Najah National University for its support.

Conflict of interest

The authors, hereby, declare no conflict of interest.

References

- Berlin B. Ethnobiological classification: Principles of categorization of plants and animals in traditional societies. New Jersey: Princeton University Press; 2014.
- Muhammad G, Hussain MA, Anwar F, Ashraf M, Gilani AH. Alhagi: a plant genus rich in bioactives for pharmaceuticals. *Phytother Res* 2015;29:1-13.
- Burdock GA. Fenaroli's handbook of flavor ingredients. 6th ed. Boca Raton: CRC press; 2016.
- Ali B, Al-Wabel NA, Shams S, Ahamad A, Khan SA, Anwar F. Essential oils used in aromatherapy: A systemic review. *Asian Pac J Trop Biomed* 2015;5: 601-11.
- Seigler DS. Plant secondary metabolism. Berlin: Springer Science & Business Media; 2012.
- Efferth T, Olbrich A, Sauerbrey A, Ross DD, Gebhart E, Neugebauer M. Activity of ascaridol from the anthelmintic herb *Chenopodium anthelminticum* L. against sensitive and multidrug-resistant tumor cells. *Anticancer Res* 2002;22:4221-4.
- Said O, Khalil K, Fulder S, Azaizeh H. Ethnopharmacological survey of medicinal herbs in Israel, the Golan Heights and the West Bank region. *J Ethnopharmacol* 2002;83:251-65.
- Fagbemi JF, Ugoji E, Adenipekun T, Adelowotan O. Evaluation of the antimicrobial properties of unripe banana (*Musa sapientum* L.), lemon grass (*Cymbopogon citratus* S.) and turmeric (*Curcuma longa* L.) on pathogens. *Afr J Biotechnol* 2009;8:1176-82.
- Dudai N, Larkov O, Ravid U, Putievsky E, Lewinsohn E. Developmental Control of Monoterpene Content and Composition in *Micromeria fruticosa*(L.) Druce. *Ann Bot* 2001;88:349-54.
- Dudai N, Yaniv Z. Endemic aromatic medicinal plants in the holy land vicinity. In: Yaniv Z, Dudai N, editors. Medicinal and aromatic plants of the Middle-East. New York: Springer; 2014. p. 37-58.
- Baser KHC, Kirimer N, Özek T, Tümen G, Karaer F. Essential oil composition of three Labiatae endemic to Turkey (*Micromeria fruticosa* (L.) Druce subsp. *giresunica* PH Davis, *Sideritis lycia* Boiss. et Heldr. and *S. arguta* Boiss. et Heldr.). *J Essent Oil Res* 1996; 8:699-701.
- Dafni A, Yaniv Z, Palevitch D. Ethnobotanical survey of medicinal plants in northern Israel. *J Ethnopharmacol* 1984;10:295-310.
- Jaradat N. Ethnopharmacological survey of natural products in palestine. *An-Najah Univ J Res (N Sc)* 2005;19:13-67.
- Güllüce M, Sökmen M, Şahin F, Sökmen A, Adigüzel A, Özer H. Biological activities of the essential oil and methanolic extract of *Micromeria fruticosa* (L) Druce ssp *serpyllifolia* (Bieb) PH Davis plants from the eastern Anatolia region of Turkey. *J Sci Food Agric* 2004;84:735-41.
- Jaradat N. Quantitative estimations for the volatile Oil by using hydrodistillation and microwave accelerated distillation methods from *Ruta graveolens* L. and *Ruta chalepensis* L. leaves from Jerusalem area/ Palestine. *Mor J Chem* 2016;4:1-6.
- Spencer MR, Gastin PB. Energy system contribution during 200-to 1500-m running in highly trained athletes. *Med Sci Sports Exer* 2001;33:157-62.
- Walker BR, Colledge NR, Ralston SH. Davidson's principles and practice of medicine. 22nd ed. New York: Churchill Livingstone Elsevier; 2013.
- Holguin F. The metabolic syndrome as a risk factor for lung function decline. *Am J Respir Crit Care Med* 2012;185:352-3
- Schneider A, Gindner L, Tilemann L, Schermer T, Dinant GJ, Meyer FJ, et al. Diagnostic accuracy of spirometry in primary care. *BMC Pulm Med* 2009;9: 31.
- Quadrelli S, Bosio M, Salvado A, Chertcoff J. Valor de la espirometría para el diagnostico de restricción pulmonary. *Medicina (BAires)* 2007;67:685-90.
- Pellegrino R, Viegi G, Brusasco V, Crapo RO, Burgos F, Casaburi R, et al. Interpretative strategies for lung function tests. *Eur Respir J* 2005;26:948-68.
- Derom E, van Weel C, Liistro G, Buffels J, Schermer T, Lammers E, et al. Primary care spirometry. *Eur Respir J* 2008;31:197-203.
- Broekhuizen BD, Sachs AP, Hoes AW, Moons KG, van den Berg JW, Dalinghaus WH, et al. Undetected chronic obstructive pulmonary disease and asthma

- in people over 50 years with persistent cough. *Br J Gen Pract* 2010;60:489-94.
24. Jaradat NA, Al Zabadi H, Rahhal B, Hussein AM, Mahmoud JS, Mansour B, et al. The effect of inhalation of *Citrus sinensis* flowers and *Mentha spicata* leave essential oils on lung function and exercise performance: a quasi-experimental uncontrolled before-and-after study. *J Int Soc Sports Nutr* 2016;13: 36-44.
 25. Kunnumakkara AB, Koca C, Dey S, Gehlot P, Yodkeeree S, Danda D, et al. Traditional uses of spices: an overview. In: Aggarwal AB, Kunnumakkara AB, editors. *Molecular targets and therapeutic uses of spices*. New Jersey: World Scientific; 2009. p. 1-24.
 26. Tofino-Rivera A, Ortega-Cuadros M, Galvis-Pareja D, Jimenez-Rios H, Merini LJ, Martinez-Pabon MC. Effect of *Lippia alba* and *Cymbopogon citratus* essential oils on biofilms of *Streptococcus mutans* and cytotoxicity in CHO cells. *J Ethnopharmacol* 2016;194:749-54.
 27. Chukwuocha UM, Fernandez-Rivera O, Legorreta-Herrera M. Exploring the antimalarial potential of whole *Cymbopogon citratus* plant therapy. *J Ethnopharmacol* 2016;193:517-23.
 28. Ortega-Ramirez LA, Silva-Espinoza BA, Vargas-Arispuro I, Gonzalez-Aguilar GA, Cruz-Valenzuela MR, Nazzaro F, et al. Combination of *Cymbopogon citratus* and *Allium cepa* essential oils increased antibacterial activity in leafy vegetables. *J Sci Food Agric* 2017;97: 2166-73.
 29. Alzawqari MH, Al-Baddany AA, Al-Baadani HH, Alhidary IA, Khan RU, Aqil GM, et al. Effect of feeding dried sweet orange (*Citrus sinensis*) peel and lemon grass (*Cymbopogon citratus*) leaves on growth performance, carcass traits, serum metabolites and antioxidant status in broiler during the finisher phase. *Environ Sci Pollut Res Int* 2016;23:17077-82.
 30. Abu-Gharbieh E, Shehab NG, Khan SA. Anti-inflammatory and gastroprotective activities of the aqueous extract of *Micromeria fruticosa* (L.) Druce ssp *Serpyllifolia* in mice. *Pak J Pharm Sci* 2013;26: 799-803.
 31. Shehab NG, Abu-Gharbieh E. Constituents and biological activity of the essential oil and the aqueous extract of *Micromeria fruticosa* (L.) Druce subsp. *serpyllifolia*. *Pak J Pharm Sci* 2012;25:687-92.
 32. Lima NG, De Sousa DP, Pimenta FC, Alves MF, De Souza FS, Macedo RO, et al. Anxiolytic-like activity and GC-MS analysis of (R)-(+)-limonene fragrance, a natural compound found in foods and plants. *Pharmacol Biochem Behav* 2013;103:450-4.
 33. Liu P, Kong M, Yuan S, Liu J, Wang P. History and experience: a survey of traditional chinese medicine treatment for Alzheimer's disease. *Evid Based Complement Alternat Med* 2014; 2014:642128.
 34. Goldbeck-Wood S, Dorozynski A, Lie LG, Yamauchi M, Zinn C, Josefson D. Complementary medicine is booming worldwide. *BMJ* 1996;313:131-3.
 35. Lv XN, Liu ZJ, Zhang HJ, Tzeng CM. Aromatherapy and the central nerve system (CNS): therapeutic mechanism and its associated genes. *Curr Drug Targets* 2013;14:872-9.