

เอกสารอ้างอิง

- Bradbury PC** (1989) Evidence for hymenostome affinities in an apostome ciliate. *J Protozool* **36**: 95–103
- Bradbury PC, Song W, Zhang L** (1997) Stomatogenesis during the formation of the tomite of *Hyalophysa chattoni* (Hymenostomatida: Ciliophora). *Eur J Protistol* **33**: 409–419
- Chatton E, Lwoff A** (1935) Les ciliés apostomes. Morphologie, cytologie, éthologie, évolution, systématique. Première partie. Aperçu historique et général. Étude monographique des genres et des espèces. *Arch Zool Exp Gén* **77**: 1–453
- Clamp JC, Bradbury PC, Strüder-Kypke MC, Lynn DH** (2008) Phylogenetic position of the apostome ciliates (phylum Ciliophora, subclass Apostomatia) tested using small subunit rRNA gene sequences. *Denisia* **23**: 395–402
- Corliss JO** (1979) The ciliated protozoa: characterization, classification, and guide to the literature. 2nd ed. Pergamon Press, London and New York, pp 455
- de Puytorac P, Grain J** (1975) Étude de la tomitogenèse et de l'ultrastructure de *Collinia orchestiae*, cilié apostome sanguicole, endoparasite du crustacé *Orchestia gammarella* Pallas. *Protistologica* **11**: 61–74
- de Puytorac P, Lom J** (1962) La tomitogenèse des *Collinia* ciliés apostomes sanguicoles endoparasites des crustacés. *Ann Parasitol Humaine Comp* **37**: 195–209
- Gómez-Gutiérrez J, Peterson WT, De Robertis A, Brodeur RD** (2003) Mass mortality of krill caused by parasitoid ciliates. *Science* **301**: 339
- Gómez-Gutiérrez J, Strüder-Kypke MC, Lynn DH, Shaw TC, Aguilar-Méndez MJ, López-Cortés A, Martínez-Gómez S, Robinson CJ** (2012) *Pseudocollinia brintoni* gen. nov., sp. nov. (Apostomatida: Colliniidae), a parasitoid ciliate infecting the euphausiid *Nyctiphanes simplex*. *Dis Aquat Org* **99**: 57–78
- Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O** (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* **59**: 307–321
- Guo Z, Liu S, Hu S, Li T, Huang Y, Liu G, Zhang H, Lin S** (2012) Prevalent ciliate symbiosis on copepods: high genetic diversity and wide distribution detected using small subunit ribosomal RNA gene. *PLoS ONE* **7**: e44847. doi:10.1371/journal.pone.0044847
- Huelsenbeck JP, Ronquist F** (2001) MrBayes: Bayesian inference of phylogenetic trees. *Bioinformatics* **17**: 754–755
- Jankowski AW** (1980) Conspectus of a new system of the phylum Ciliophora. *Trudy Zool Inst Leningrad* **94**: 103–121 (in Russian)
- Jankowski AW** (2007) Phylum Ciliophora Doflein, 1901. In Alimov AF (ed), *Protista, Part 2, Handbook on Zoology*, Russian Academy of Sciences, Zoological Institute, St. Petersburg, pp 415–993 (in Russian with English summary)
- Jukes TH, Cantor CR** (1969) Evolution of protein molecules. In Munro HN (ed) *Mammalian protein metabolism*. Academic Press, New York, pp 21–132
- Katoh K, Toh H** (2008) Recent developments in the MAFFT multiple sequence alignment program. *Brief Bioinform* **9**: 286–298

- Kimura M** (1980) A simple model for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* **16**: 111–120
- Lynn DH** (2008) *The ciliated protozoa: characterization, classification, and guide to the literature*. 3rd ed. Springer, Dordrecht, Second Printing 2010, pp 606
- Montagnes DJS, Lynn DH** (1993) A quantitative protargol stain (QPS) for ciliates and other protists. In Kemp P, Sherr B, Sherr E, Cole J (eds) *Aquatic microbial ecology*. Lewis Publishers, London, UK, pp 229–240
- Posada D** (2008) jModelTest: phylogenetic model averaging. *Mol Biol Evol* **25**: 1253–1256
- Prokopowicz AJ, Rueckert S, Leander BS, Michaud J, Fortier L** (2010) Parasitic infection of the hyperiid amphipod *Themisto libellula* in the Canadian Beaufort Sea (Arctic Ocean), with a description of *Ganymedes themistos* sp. n. (Apicomplexa, Eugregarinorida). *Polar Biol* **33**: 1339–1350
- Ronquist F, Huelsenbeck JP** (2003) MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**: 1572–1574
- Small EB, Lynn DH** (1981) A new macrosystem for the Phylum Ciliophora Doflein, 1901. *BioSystems* **14**: 387–401
- Small EB, Lynn DH** (1985) Phylum Ciliophora Doflein, 1901. In Lee JJ, Hutner SH, Bovee EC (eds) *An illustrated guide to the protozoa*. Society of Protozoologists, Lawrence, Kansas, pp 393–575
- Tajima F, Nei M** (1984) Estimation of evolutionary distance between nucleotide sequences. *Mol Biol Evol* **1**: 269–285
- Tamura K** (1992) Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G + C-content biases. *Mol Biol Evol* **9**: 678–687
- Tamura K, Kumar S** (2002) Evolutionary distance estimation under heterogeneous substitution pattern among lineages *Mol Biol Evol* **19**: 1727–1736
- Tamura K, Nei M** (1993) Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol* **10**: 512–526
- Tamura K, Nei M, Kumar S** (2004) Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc Natl Acad Sci USA* **101**: 11030–11035
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S** (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* **28**: 2731–2739