

# ภาคผนวก

## ภาคผนวก ก

การสังเคราะห์สารตั้งต้น  $[\text{Cu}(\text{II})(\text{bipyam})(\text{O}_2\text{CCH}_2\text{CH}_3)_2] \cdot 2\text{H}_2\text{O}$ 

การสังเคราะห์สารประกอบเชิงซ้อน  $[\text{Cu}(\text{II})(\text{bipyam})(\text{O}_2\text{CCH}_2\text{CH}_3)_2] \cdot 2\text{H}_2\text{O}$  โดยวิธีเตรียมโดยตรงเพื่อใช้เป็นสารตั้งต้นในการสังเคราะห์ reoxygenation products โดยวิธีรีดิวซ์ด้วยโลหะคอปเปอร์ในบรรยากาศปกติ และภายใต้บรรยากาศก๊าซไนโตรเจน มีขั้นตอนดังนี้

1. ละลาย  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2 \cdot \text{H}_2\text{O}$  0.19 กรัม (1.5 มิลลิโมล) ในน้ำ 20 มล. และเติมกรดโพรพอออนิกเพื่อช่วยในการละลาย อุณหภูมิห้อง
2. ละลาย bipyam 0.27 กรัม (1.57 มิลลิโมล) ในเอทานอล 30 มล. อุณหภูมิห้อง
3. ผสมสารละลายจากข้อ 1 และข้อ 2 เข้าด้วยกันแล้วเติมโซเดียมโพรพอเนต 0.31 กรัม (3.2 มิลลิโมล) อุณหภูมิห้อง ตั้งสารละลายที่ได้ให้ระเหยอย่างช้า ๆ ที่อุณหภูมิห้อง ประมาณ 1 สัปดาห์จะได้ผลึกสีเขียวใสของ  $[\text{Cu}(\text{II})(\text{bipyam})(\text{O}_2\text{CCH}_2\text{CH}_3)_2] \cdot 2\text{H}_2\text{O}$

## ภาคผนวก ข

พิกัดของอะตอม ความยาวและมุมพันธะและเทอร์มอลพารามิเตอร์ของอะตอม

ในโมเลกุลของ  $[\text{Cu}(\text{II})(\text{bipyam})_2(\text{O}_2\text{CCH}_2\text{CH}_3)][\text{NO}_3]$ ตารางที่ พ1 พิกัดของอะตอมในโมเลกุล  $[\text{Cu}(\text{bipyam})_2(\text{O}_2\text{CCH}_2\text{CH}_3)][\text{NO}_3]$  (I)

อะตอม	x/a	y/b	z/c	U
Cu	0.18056	0.57010	0.12839	* 0.0449
N(1)	0.04522	0.67502	0.12219	* 0.0499
C(1)	-0.01349	0.65625	0.06980	* 0.0629
C(2)	-0.11508	0.68420	0.06490	* 0.0777
C(3)	-0.16285	0.73476	0.11634	* 0.0849
C(4)	-0.10694	0.75745	0.16935	* 0.0733
C(5)	-0.00062	0.72755	0.17136	* 0.0525
N(3)	0.05548	0.75596	0.22503	* 0.0579
C(6)	0.15913	0.77856	0.23470	* 0.0490
C(7)	0.19363	0.86652	0.28485	* 0.0668
C(8)	0.29575	0.89443	0.29440	* 0.0754
C(9)	0.36347	0.83659	0.25398	* 0.0685
C(10)	0.32610	0.74859	0.20631	* 0.0562
N(2)	0.22467	0.71713	0.19607	* 0.0484
N(4)	0.30398	0.43286	0.13748	* 0.0478
C(11)	0.32195	0.36036	0.19264	* 0.0593
C(12)	0.37563	0.22867	0.20005	* 0.0678
C(13)	0.41406	0.16058	0.14746	* 0.0718
C(14)	0.40003	0.23288	0.09252	* 0.0624
C(15)	0.34605	0.37155	0.08847	* 0.0467
N(6)	0.33750	0.44455	0.03208	* 0.0535
C(16)	0.31580	0.59677	0.01835	* 0.0507
C(17)	0.35694	0.66266	-0.03370	* 0.0681
C(18)	0.33838	0.81170	-0.04649	* 0.0811
C(19)	0.28063	0.89906	-0.00798	* 0.0819
C(20)	0.24114	0.82714	0.04088	* 0.0709
N(5)	0.25531	0.67724	0.05405	* 0.0540
O(1)	0.11225	0.38290	0.08958	* 0.0636
O(2)	0.06859	0.36196	0.18490	* 0.0817
C(21)	0.07003	0.30923	0.13189	* 0.0776
C(22)	0.05042	0.13796	0.11821	* 0.1312
C(22')	0.00611	0.16136	0.12748	* 0.1776
C(23)	-0.05478	0.14728	0.08484	* 0.1705
C(23')	0.01382	0.09127	0.06401	* 0.1170
N(7)	0.56164	0.74544	0.10821	* 0.0620
O(3)	0.52705	0.61855	0.12193	* 0.1115
O(3')	0.56215	0.61501	0.08566	* 0.1061
O(4)	0.56946	0.84622	0.14696	* 0.1105
O(4')	0.56881	0.77153	0.16277	* 0.1258
O(5)	0.59591	0.76903	0.05782	* 0.0903
O(5')	0.56245	0.85404	0.07093	* 0.1792

ตารางที่ ๗2 ความยาวพันธะในโมเลกุล  $[\text{Cu}(\text{bipyam})_2(\text{O}_2\text{CCH}_2\text{CH}_3)][\text{NO}_3]$  (I)

พันธะ	ความยาวพันธะ (Å)	พันธะ	ความยาวพันธะ (Å)
N(7)-O(3)	1.2326(1)	C(7)-C(8)	1.3613(5)
N(7)-O(3')	1.2344(1)	C(8)-H(8)	.9600(0)
N(7)-O(4)	1.2178(1)	C(8)-C(9)	1.3745(2)
N(7)-O(4')	1.2121(2)	C(9)-H(9)	.9601(2)
N(7)-O(5)	1.2253(1)	C(9)-C(10)	1.3623(1)
N(7)-O(5')	1.2458(1)	C(10)-H(10)	.9599(1)
O(3)-O(3')	.9349(1)	C(10)-N(2)	1.3601(3)
O(3)-O(4)	2.1163(2)	N(4)-C(11)	1.3692(1)
O(3)-O(4')	1.6756(1)	N(4)-C(15)	1.3371(1)
O(3)-O(5)	2.1440(2)	C(11)-H(11)	.9599(0)
O(3)-O(5')	2.3833(2)	C(11)-C(12)	1.3462(1)
O(3')-O(4')	2.1620(2)	C(12)-H(12)	.9601(1)
O(3')-O(5)	1.5415(1)	C(12)-C(13)	1.4059(1)
O(3')-O(5')	2.0989(2)	C(13)-H(13)	.9600(1)
O(4)-O(4')	.7346(0)	C(13)-C(14)	1.3586(1)
O(4)-O(5)	2.1034(3)	C(14)-H(14)	.9600(0)
O(4)-O(5')	1.6601(2)	C(14)-C(15)	1.3951(1)
O(4')-O(5)	2.3365(3)	C(15)-N(6)	1.3847(2)
O(4')-O(5')	2.1277(3)	N(6)-H(6B)	.9000(0)
O(5)-O(5')	.9106(0)	N(6)-C(16)	1.3808(1)
Cu-N(1)	1.9867(4)	C(16)-C(17)	1.4026(1)
Cu-N(2)	2.0154(2)	C(16)-N(5)	1.3340(1)
Cu-N(4)	2.0059(3)	C(17)-H(17)	.9601(0)
Cu-N(5)	2.1466(2)	C(17)-C(18)	1.3426(1)
Cu-O(1)	2.0193(2)	C(18)-H(18)	.9600(0)
Cu-O(2)	2.6643(3)	C(18)-C(19)	1.3829(1)
N(1)-C(1)	1.3558(2)	C(19)-H(19)	.9601(1)
N(1)-C(5)	1.3349(1)	C(19)-C(20)	1.3594(1)
C(1)-H(1)	.9600(1)	C(20)-H(20)	.9600(0)
C(1)-C(2)	1.3477(3)	C(20)-N(5)	1.3430(1)
C(2)-H(2)	.9599(1)	O(1)-O(2)	2.1928(3)
C(2)-C(3)	1.3828(1)	O(1)-C(21)	1.2716(1)
C(3)-H(3)	.9600(2)	O(2)-C(21)	1.2460(1)
C(3)-C(4)	1.3528(2)	C(21)-C(22)	1.5349(2)
C(4)-H(4)	.9600(1)	C(21)-C(22')	1.5311(2)
C(4)-C(5)	1.4109(3)	C(22)-C(22')	.6545(1)
C(5)-N(3)	1.3740(2)	C(22)-C(23)	1.5256(3)
N(3)-H(3A)	.8999(1)	C(22)-C(23')	1.3187(1)
N(3)-C(6)	1.3737(3)	C(22')-C(23)	1.2010(1)
C(6)-C(7)	1.3919(1)	C(22')-C(23')	1.5217(2)
C(6)-N(2)	1.3412(2)	C(23)-C(23')	1.1337(2)
C(7)-H(7)	.9600(1)		

ตารางที่ ๓3 มุมพันธะในโมเลกุล  $[\text{Cu}(\text{bipyam})_2(\text{O}_2\text{CCH}_2\text{CH}_3)][\text{NO}_3]$  (I)

มุมพันธะ	ขนาดมุมพันธะ (°)	มุมพันธะ	ขนาดมุมพันธะ (°)
O(3)-N(7)-O(3')	44.537(6)	O(3')-O(4')-O(4)	100.891(9)
O(3)-N(7)-O(4)	119.457(8)	O(3')-O(4')-O(5)	39.841(5)
O(3)-N(7)-O(4')	86.531(5)	O(3')-O(4')-O(5')	58.580(8)
O(3)-N(7)-O(5)	121.451(2)	O(4)-O(4')-O(5)	62.692(4)
O(3)-N(7)-O(5')	148.154(4)	O(4)-O(4')-O(5')	42.333(1)
O(3')-N(7)-O(4)	159.026(1)	O(5)-O(4')-O(5')	22.928(4)
O(3')-N(7)-O(4')	124.183(2)	N(7)-O(5)-O(3)	29.309(3)
O(3')-N(7)-O(5)	77.612(5)	N(7)-O(5)-O(3')	51.457(3)
O(3')-N(7)-O(5')	115.615(9)	N(7)-O(5)-O(4)	30.475(4)
O(4)-N(7)-O(4')	35.191(3)	N(7)-O(5)-O(4')	16.456(3)
O(4)-N(7)-O(5)	118.841(7)	N(7)-O(5)-O(5')	69.578(6)
O(4)-N(7)-O(5')	84.72(1)	O(3)-O(5)-O(3')	22.6757(9)
O(4')-N(7)-O(5)	146.904(6)	O(3)-O(5)-O(4)	59.760(7)
O(4')-N(7)-O(5')	119.909(8)	O(3)-O(5)-O(4')	43.659(4)
O(5)-N(7)-O(5')	43.236(4)	O(3)-O(5)-O(5')	93.729(9)
N(7)-O(3)-O(3')	67.836(5)	O(3')-O(5)-O(4)	81.253(7)
N(7)-O(3)-O(4)	30.070(5)	O(3')-O(5)-O(4')	63.970(4)
N(7)-O(3)-O(4')	46.226(7)	O(3')-O(5)-O(5')	115.305(9)
N(7)-O(3)-O(5)	29.180(2)	O(4)-O(5)-O(4')	18.079(3)
N(7)-O(3)-O(5')	16.010(2)	O(4)-O(5)-O(5')	49.311(2)
O(3')-O(3)-O(4)	96.699(5)	O(4')-O(5)-O(5')	65.548(3)
O(3')-O(3)-O(4')	108.475(8)	N(7)-O(5')-O(3)	15.836(2)
O(3')-O(3)-O(5)	39.471(5)	N(7)-O(5')-O(3')	32.028(4)
O(3')-O(3)-O(5')	61.174(5)	N(7)-O(5')-O(4)	46.926(5)
O(4)-O(3)-O(4')	17.956(2)	N(7)-O(5')-O(4')	29.592(2)
O(4)-O(3)-O(5)	59.166(7)	N(7)-O(5')-O(5)	67.187(8)
O(4)-O(3)-O(5')	42.789(7)	O(3)-O(5')-O(3')	22.968(3)
O(4')-O(3)-O(5)	74.293(9)	O(3)-O(5')-O(4)	59.994(4)
O(4')-O(3)-O(5')	60.273(9)	O(3)-O(5')-O(4')	43.148(2)
O(5)-O(3)-O(5')	22.412(2)	O(3)-O(5')-O(5)	63.858(9)
N(7)-O(3')-O(3)	67.627(4)	O(3')-O(5')-O(4)	78.854(1)
N(7)-O(3')-O(4')	27.632(1)	O(3')-O(5')-O(4')	61.527(2)
N(7)-O(3')-O(5)	50.931(7)	O(3')-O(5')-O(5)	41.603(6)
N(7)-O(3')-O(5')	32.358(5)	O(4)-O(5')-O(4')	17.337(3)
O(3)-O(3')-O(4')	47.314(6)	O(4)-O(5')-O(5)	106.112(4)
O(3)-O(3')-O(5)	117.853(4)	O(4')-O(5')-O(5)	91.524(6)
O(3)-O(3')-O(5')	95.859(2)	N(1)-Cu-N(2)	88.937(8)
O(4')-O(3')-O(5)	76.188(9)	N(1)-Cu-N(4)	170.5814
O(4')-O(3')-O(5')	59.892(7)	N(1)-Cu-N(5)	100.70(1)
O(5)-O(3')-O(5')	23.092(3)	N(1)-Cu-O(1)	88.08(1)
N(7)-O(4)-O(3)	30.473(3)	N(1)-Cu-O(2)	80.55(1)
N(7)-O(4)-O(4')	71.983(9)	N(2)-Cu-N(4)	95.949(9)
N(7)-O(4)-O(5)	30.684(3)	N(2)-Cu-N(5)	99.000(9)
N(7)-O(4)-O(5')	48.352(6)	N(2)-Cu-O(1)	157.163(4)
O(3)-O(4)-O(4')	44.686(7)	N(2)-Cu-O(2)	103.473(9)
O(3)-O(4)-O(5)	61.074(3)	N(4)-Cu-N(5)	86.52(1)
O(3)-O(4)-O(5')	77.217(3)	N(4)-Cu-O(1)	34.29(1)
O(4')-O(4)-O(5)	99.229(8)	N(4)-Cu-O(2)	90.46(1)
O(4')-O(4)-O(5')	120.329(4)	N(5)-Cu-O(1)	103.802(9)
O(5)-O(4)-O(5')	24.577(5)	N(5)-Cu-O(2)	157.517(2)
N(7)-O(4')-O(3)	47.243(3)	O(1)-Cu-O(2)	53.716(7)
N(7)-O(4')-O(3')	28.185(3)	Cu-N(1)-C(1)	117.842(8)
N(7)-O(4')-O(4)	72.825(6)	Cu-N(1)-C(5)	122.265(7)
N(7)-O(4')-O(5)	16.640(3)	C(1)-N(1)-C(5)	117.56(1)
N(7)-O(4')-O(5')	30.499(5)	N(1)-C(1)-H(1)	117.59(1)
O(3)-O(4')-O(3')	24.212(3)	N(1)-C(1)-C(2)	124.012(8)
O(3)-O(4')-O(4)	117.358(8)	H(1)-C(1)-C(2)	118.392(8)
O(3)-O(4')-O(5)	62.048(5)	C(1)-C(2)-H(2)	120.812(7)
O(3)-O(4')-O(5')	76.580(7)	C(1)-C(2)-C(3)	118.280(8)

ตารางที่ ๒3 มุมพันธะในโมเลกุล  $[\text{Cu}(\text{bipyam})_2(\text{O}_2\text{CCH}_2\text{CH}_3)][\text{NO}_3]$  (I) (ต่อ)

มุมพันธะ	ขนาดมุมพันธะ (°)	มุมพันธะ	ขนาดมุมพันธะ (°)
H(2)-C(2)-C(3)	120.90(1)	C(15)-N(6)-C(16)	129.613(2)
C(2)-C(3)-H(3)	120.167(7)	H(6B)-N(6)-C(16)	115.315(7)
C(2)-C(3)-C(4)	119.71(1)	N(6)-C(16)-C(17)	118.901(5)
H(3)-C(3)-C(4)	120.124(7)	N(6)-C(16)-N(5)	119.660(5)
C(3)-C(4)-H(4)	120.80(1)	C(17)-C(16)-N(5)	121.431(7)
C(3)-C(4)-C(5)	119.280(8)	C(16)-C(17)-H(17)	119.786(8)
H(4)-C(4)-C(5)	119.921(6)	C(16)-C(17)-C(18)	119.267(5)
N(1)-C(5)-C(4)	121.119(7)	H(17)-C(17)-C(18)	120.947(6)
N(1)-C(5)-N(3)	120.42(1)	C(17)-C(18)-H(18)	119.389(4)
C(4)-C(5)-N(3)	118.456(8)	C(17)-C(18)-C(19)	119.977(5)
C(5)-N(3)-H(3A)	115.33(1)	H(18)-C(18)-C(19)	120.634(8)
C(5)-N(3)-C(6)	129.580(7)	C(18)-C(19)-H(19)	121.048(5)
H(3A)-N(3)-C(6)	115.09(1)	C(18)-C(19)-C(20)	117.697(7)
N(3)-C(6)-C(7)	118.556(3)	H(19)-C(19)-C(20)	121.255(5)
N(3)-C(6)-N(2)	120.04(1)	C(19)-C(20)-H(20)	119.060(7)
C(7)-C(6)-N(2)	121.40(1)	C(19)-C(20)-N(5)	124.001(4)
C(6)-C(7)-H(7)	120.14(1)	H(20)-C(20)-N(5)	116.939(6)
C(6)-C(7)-C(8)	119.686(4)	Cu-N(5)-C(16)	121.297(8)
H(7)-C(7)-C(8)	120.18(1)	Cu-N(5)-C(20)	121.231(5)
C(7)-C(8)-H(8)	120.603(3)	C(16)-N(5)-C(20)	117.456(5)
C(7)-C(8)-C(9)	119.51(1)	Cu-O(1)-O(2)	78.357(7)
H(8)-C(8)-C(9)	119.86(1)	Cu-O(1)-C(21)	107.325(8)
C(8)-C(9)-H(9)	120.98(1)	O(2)-O(1)-C(21)	29.096(3)
C(8)-C(9)-C(10)	118.55(1)	Cu-O(2)-O(1)	47.927(3)
H(9)-C(9)-C(10)	120.434(4)	Cu-O(2)-C(21)	77.563(7)
C(9)-C(10)-H(10)	118.76(1)	O(1)-O(2)-C(21)	29.753(5)
C(9)-C(10)-N(2)	123.223(3)	O(1)-C(21)-O(2)	121.152(7)
H(10)-C(10)-N(2)	118.01(1)	O(1)-C(21)-C(22)	114.821(6)
Cu-N(2)-C(6)	123.00(1)	O(1)-C(21)-C(22')	129.065(5)
Cu-N(2)-C(10)	119.269(3)	O(2)-C(21)-C(22)	121.771(2)
C(6)-N(2)-C(10)	117.57(1)	O(2)-C(21)-C(22')	109.472(4)
Cu-N(4)-C(11)	117.535(5)	C(22)-C(21)-C(22')	24.652(4)
Cu-N(4)-C(15)	121.088(7)	C(21)-C(22)-C(22')	77.352(7)
C(11)-N(4)-C(15)	117.526(8)	C(21)-C(22)-C(23)	100.340(3)
N(4)-C(11)-H(11)	117.812(8)	C(21)-C(22)-C(23')	121.464(1)
N(4)-C(11)-C(12)	124.174(4)	C(22')-C(22)-C(23)	48.88(1)
H(11)-C(11)-C(12)	118.910(6)	C(22')-C(22)-C(23')	94.92(1)
C(11)-C(12)-H(12)	121.106(4)	C(23)-C(22)-C(23')	46.275(4)
C(11)-C(12)-C(13)	117.579(6)	C(21)-C(22')-C(22)	78.00(1)
H(12)-C(12)-C(13)	121.304(8)	C(21)-C(22')-C(23)	118.185(7)
C(12)-C(13)-H(13)	120.479(6)	C(21)-C(22')-C(23')	109.402(5)
C(12)-C(13)-C(14)	119.306(8)	C(22)-C(22')-C(23)	106.88(1)
H(13)-C(13)-C(14)	120.202(5)	C(22)-C(22')-C(23')	59.702(6)
C(13)-C(14)-H(14)	120.350(8)	C(23)-C(22')-C(23')	47.430(8)
C(13)-C(14)-C(15)	120.017(4)	C(22)-C(23)-C(22')	24.239(3)
H(14)-C(14)-C(15)	119.632(6)	C(22)-C(23)-C(23')	57.20(1)
N(4)-C(15)-C(14)	121.222(6)	C(22')-C(23)-C(23')	81.29(1)
N(4)-C(15)-N(6)	120.772(8)	C(22)-C(23')-C(22')	25.376(7)
C(14)-C(15)-N(6)	118.005(5)	C(22)-C(23')-C(23)	76.52(1)
C(15)-N(6)-H(6B)	115.07(1)	C(22')-C(23')-C(23)	51.277(6)

ตารางที่ ๗4 เทอร์มอลพารามิเตอร์ของอะตอมในโมเลกุล  $[\text{Cu}(\text{bipyam})_2(\text{O}_2\text{CCH}_2\text{CH}_3)]$   
 $[\text{NO}_3] (\text{I})$

อะตอม	U11	U22	U33	U23	U13	U12
Cu	.0405	.0573	.0382	-.0043	.0134	-.0062
N(1)	.0422	.0657	.0420	-.0029	.0053	-.0064
C(1)	.0617	.0741	.0524	-.0078	-.0014	-.0039
H(1)	.080					
C(2)	.0650	.0941	.0721	-.0096	-.0157	.0028
H(2)	.080					
C(3)	.0412	.1094	.1025	.0004	-.0115	.0118
H(3)	.080					
C(4)	.0461	.0989	.0765	.0067	.0188	.0001
H(4)	.080					
C(5)	.0415	.0656	.0511	-.0023	.0094	-.0034
N(3)	.0440	.0871	.0441	.0013	.0149	-.0162
H(3A)	.080					
C(6)	.0531	.0554	.0390	.0029	.0073	-.0055
C(7)	.0748	.0778	.0479	-.0034	.0045	-.0150
H(7)	.080					
C(8)	.0922	.0752	.0566	-.0104	-.0201	-.0134
H(8)	.080					
C(9)	.0577	.0661	.0799	-.0109	-.0142	.0024
H(9)	.080					
C(10)	.0410	.0605	.0673	-.0044	.0036	-.0016
H(10)	.080					
N(2)	.0415	.0593	.0449	-.0016	.0067	-.0054
N(4)	.0451	.0555	.0439	-.0001	.0116	-.0029
C(11)	.0582	.0680	.0528	-.0045	.0141	.0032
H(11)	.080					
C(12)	.0726	.0641	.0671	.0011	.0082	.0130
H(12)	.080					
C(13)	.0658	.0599	.0900	.0105	.0060	.0021
H(13)	.080					
C(14)	.0559	.0590	.0740	.0013	.0188	-.0134
H(14)	.080					
C(15)	.0355	.0543	.0509	-.0063	.0107	-.0052
N(6)	.0495	.0678	.0443	.0018	.0130	-.0122
H(6B)	.080					
C(16)	.0384	.0742	.0402	-.0065	.0082	-.0013
C(17)	.0567	.0995	.0501	.0003	.0218	.0078
H(17)	.080					
C(18)	.0674	.1140	.0635	-.0038	.0196	.0340
H(18)	.080					
C(19)	.0821	.0760	.0897	-.0026	.0242	.0298
H(19)	.080					
C(20)	.0764	.0646	.0737	.0039	.0233	.0083
H(20)	.080					
N(5)	.0503	.0630	.0507	-.0028	.0212	.0014
O(1)	.0613	.0788	.0520	-.0145	.0170	-.0189
O(2)	.0889	.0935	.0658	-.0270	.0352	.0026
C(21)	.0733	.0740	.0875	-.0352	.0233	-.0100
C(22)	.0955	.1037	.2025	-.0540	.0891	-.0237
C(22')	.1493	.2126	.1810	-.1409	.1103	-.1389
C(23)	.1670	.1644	.1808	-.0060	.0139	-.1034
C(23')	.1558	.0668	.1196	-.0114	.0182	-.0333
N(7)	.0480	.0795	.0598	.0142	.0155	-.0047
O(3)	.1128	.0860	.1406	-.0004	.0582	.0124
O(3')	.1728	.0628	.0871	.0049	.0513	-.0249
O(4)	.1457	.1148	.0734	.0216	.0303	-.0300
O(4')	.1535	.1816	.0431	-.0064	.0130	.0099
O(5)	.1063	.1080	.0592	.0047	.0310	-.0063
O(5')	.3397	.1356	.0660	.0754	.0487	.0401

ตารางที่ ๗5 พิกัดของอะตอมไฮโดรเจนในโมเลกุล  $[\text{Cu}(\text{bipyam})_2(\text{O}_2\text{CCH}_2\text{CH}_3)]$   
 $[\text{NO}_3]$  (I)

อะตอม	x	y	z	U (eq)
H(22A)	5249 (7)	9061 (7)	1630 (7)	150
H(22B)	6081 (7)	9183 (7)	1133 (7)	150
H(23A)	4467 (13)	9829 (23)	690 (9)	258
H(23B)	4044 (13)	8212 (23)	874 (9)	258
H(23C)	4872 (13)	8333 (23)	380 (9)	258
H(22C)	4271 (9)	7751 (15)	1043 (9)	150
H(22D)	4840 (9)	8816 (15)	1535 (9)	150
H(23D)	4885 (17)	9959 (28)	565 (10)	258
H(23E)	5437 (17)	8499 (28)	317 (10)	258
H(23F)	6003 (17)	9559 (28)	807 (10)	258
H(14)	9209 (23)	8032 (37)	567 (14)	60 (9)
H(13)	9455 (32)	9361 (46)	1546 (18)	89 (13)
H(12)	8882 (26)	8147 (40)	2425 (16)	76 (10)
H(11)	7961 (26)	5816 (38)	2265 (16)	69 (10)
H(17)	8956 (28)	3957 (40)	-546 (16)	68 (10)
H(18)	8695 (35)	1382 (57)	-821 (20)	117 (15)
H(19)	7642 (35)	-172 (52)	-182 (21)	117 (15)
H(20)	7050 (30)	1114 (46)	715 (17)	89 (12)
H(1)	5213 (29)	3882 (43)	389 (17)	79 (11)
H(2)	3508 (30)	3266 (45)	266 (17)	85 (12)
H(3)	2738 (36)	2449 (50)	1119 (19)	103 (14)
H(4)	3688 (28)	2032 (41)	2045 (17)	77 (11)
H(7)	6467 (25)	993 (36)	3082 (14)	54 (9)
H(8)	8123 (30)	632 (41)	3197 (18)	71 (12)
H(9)	9238 (28)	1487 (42)	2615 (15)	68 (10)
H(10)	8762 (27)	3038 (40)	1786 (15)	77 (10)
H(6A)	8617 (23)	6094 (36)	17 (15)	55 (9)
H(3A)	5225 (25)	2120 (36)	2512 (14)	55 (9)

2155 réflexions mesurées  
1988 réflexions  
indépendantes

Affinement

Affinement à partir des  $F^2$   
 $R[F^2 > 2\sigma(F^2)] = 0,042$   
 $wR(F^2) = 0,090$   
 $S = 1,093$

1988 réflexions  
173 paramètres

Tous les paramètres des  
atomes d'hydrogène  
affinés

1 réflexion de référence  
fréquence: 120 min  
variation d'intensité: 1,7%

$w = 1/[\sigma^2(F_o^2) + (0,0263P)^2 + 1,0719P]$   
où  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0,001$   
 $\Delta\rho_{\max} = 0,334 \text{ e } \text{Å}^{-3}$   
 $\Delta\rho_{\min} = -0,300 \text{ e } \text{Å}^{-3}$

Pas de correction  
d'extinction

Facteurs de diffusion des  
*International Tables for  
Crystallography* (Tome C)

## Références

- Enraf-Nonius (1994). *CAD-4 EXPRESS*. Version 5.1/1.2. Enraf-Nonius, Delft, Les Pays Bas.
- Fair, C. K. (1990). *MOLLEN. An Interactive Intelligent System for Crystal Structure Analysis*. Enraf-Nonius, Delft, Les Pays Bas.
- Johnson, C. K. (1976). *ORTEP II*. Rapport ORNL-5138. Oak Ridge National Laboratory, Tennessee, EU.
- Lumme, P. O., Knuutila, H. & Lindell, E. (1996). *Acta Cryst.* C52, 51-56.
- North, A. C. T., Phillips, D. C. & Mathews, E. S. (1968). *Acta Cryst.* A24, 351-359.
- Sheldrick, G. M. (1990). *Acta Cryst.* A46, 467-473.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. Université de Göttingen, Allemagne.
- Whitlow, S. H. (1973). *Inorg. Chem.* 12, 2286-2289.

Tableau 1. Paramètres géométriques (Å, °)

Cu1-O1	1,954(3)	N1-C5	1,474(4)
Cu1-O1	1,954(3)	N1-C5	1,486(4)
Cu1-N1	2,059(3)	N1-C1	1,489(4)
Cu1-N1	2,059(3)	C1-C2	1,513(4)
Cu1-O2	2,328(2)	C1-H1C1	0,97(4)
Cu1-O2	2,328(2)	C1-H2C1	0,93(4)
O1-C1	1,258(4)	C3-C4	1,512(4)
O2-C1	1,206(4)	C3-H1C3	0,90(4)
O3-C1	1,301(4)	C3-H2C3	0,95(3)
O3-H1O3	1,03(4)	C5-C6	1,523(4)
O4-C6	1,187(4)	C5-H1C5	1,00(3)
O5-C6	1,374(4)	C5-H2C5	0,88(3)
O5-H1O5	0,77(5)	OW1-H1W1	0,95(7)
O6-C2	1,213(4)	OW1-H2W1	0,79(7)
O1-Cu1-O1	180,0	C3-N1-C1	112,2(3)
O1-Cu1-N1	93,9(1)	C5-N1-C1	110,5(3)
O1-Cu1-N1	86,1(1)	C3-N1-Cu1	107,4(2)
O1-Cu1-N1	86,1(1)	C5-N1-Cu1	108,4(2)
O1-Cu1-N1	93,9(1)	C1-N1-Cu1	105,9(2)
N1-Cu1-N1	180,0	N1-C1-C2	113,5(3)
O1-Cu1-O2	87,2(1)	O6-C2-O1	131,6(3)
O1-Cu1-O2	92,7(1)	O6-C2-C1	116,3(3)
O1-Cu1-O2	102,23(9)	O1-C2-C1	119,1(3)
N1-Cu1-O2	77,77(9)	N1-C3-C4	112,2(2)
O1-Cu1-O2	92,7(1)	O2-C1-O3	124,5(3)
O1-Cu1-O2	87,2(1)	O2-C1-C3	123,3(3)
N1-Cu1-O2	77,77(9)	O3-C4-C3	112,2(3)
N1-Cu1-O2	102,23(9)	N1-C5-C6	113,8(3)
O2-Cu1-O2	180,0	O4-C6-O5	124,9(3)
C2-O1-Cu1	114,4(2)	O1-C6-C5	126,9(3)
C4-O2-Cu1	105,3(2)	O5-C6-C5	108,2(3)
C3-N1-C5	112,1(2)		

Code de symétrie: (i) -x, 1-y, -z.

La largeur de balayage est  $(1,00 + 0,70 \tan \theta)^\circ$ . Les intensités ont été corrigées des facteurs de Lorentz-polarisation. La structure a été résolue par les méthodes directes (*SHELXS86*; Sheldrick, 1990) puis affinée par la méthode des moindres carrés (*SHELXL93*; Sheldrick, 1993).

Collection des données: *CAD-4 EXPRESS* (Enraf-Nonius, 1994). Affinement des paramètres de la maille: *CAD-4 EXPRESS*. Réduction des données: *MOLLEN* (Fair, 1990). Graphisme moléculaire: *ORTEP II* (Johnson, 1976). Logiciel utilisé pour préparer le matériel pour publication: *SHELXL93*.

Des documents complémentaires concernant cette structure peuvent être obtenus à partir des archives électroniques de IUCr (Référence: MU1346). Les processus d'accès à ces archives est donné au dos de la couverture.

*Acta Cryst.* (1998), C54, 199-201

## Bis(di-2-pyridyl-N-amine)(propionato-O)-copper(II) Nitrate

SUJITRA YOUNGMIE,<sup>a</sup> KAMPHONG CHANDAVONG,<sup>a</sup>  
CHAVENG PAKAWATCHAI,<sup>b</sup> ZHONG-YUAN ZHOU<sup>c</sup> AND  
HOONG-KUN FUI<sup>d</sup>

<sup>a</sup>Department of Chemistry, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand, <sup>b</sup>Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat Yai 90112, Thailand, <sup>c</sup>Crystal Structure Analysis Laboratory, Chengdu Branch, Academia Sinica, Sichuan 610041, People's Republic of China, and <sup>d</sup>X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia. E-mail: hkfui@usm.my

(Received 27 June 1997; accepted 21 October 1997)


## Abstract

The crystal structure of the title compound comprises  $[\text{Cu}(\text{C}_3\text{H}_5\text{O}_2)(\text{C}_{10}\text{H}_9\text{N}_3)_2]^+$  cations and  $\text{NO}_3^-$  anions. The cation structure is intermediate between a distorted square-based pyramidal five-coordinate geometry with an additional long bond and an asymmetric *cis*-distorted octahedral geometry, both giving a  $(4+1+1^-)$ -type coordination.

## Comment

The crystal structures of the  $[\text{Cu}^{\text{II}}(\text{chelate})_2(\text{OXO})]^+$  cations, where chelate is di-2-bipyridyl or 1,10-*o*-phenanthroline and OXO is  $\text{ONO}^-$ ,  $\text{CH}_3\text{CO}_2^-$  or  $\text{HCO}_2^-$  are well characterized (Hathaway, 1984) as fluxional copper(II) stereochemistries (Hitzgerald *et al.*, 1981; Simmonds *et al.*, 1987). In order to extend this series to more flexible chelate nitrogen ligands, complexes with

the ligand di-2-pyridylamine (dpyam) have been examined. Only a few crystal structures of the dpyam complexes in this series have been reported (Aduldecha *et al.*, 1991).



The asymmetric unit comprises discrete [Cu(C<sub>3</sub>H<sub>5</sub>O<sub>2</sub>)(C<sub>10</sub>H<sub>9</sub>N<sub>3</sub>)<sub>2</sub>]<sup>+</sup> cations and NO<sub>3</sub><sup>-</sup> anions. The bond lengths and angles in the structure agree with those observed in bis(di-2-pyridylamine)(acetato)copper(II) nitrate (Aduldecha *et al.*, 1991). The Cu atom is basically five coordinate, but the second O atom of the propionate group occupies a sixth position at a distance of 2.669 (3) Å. Thus, the CuN<sub>4</sub>O<sub>2</sub> chromophore involves a *cis*-distorted octahedral stereochemistry with an asymmetric coordination of the OXO<sup>-</sup> anion. However, the structure can also be described as involving a square-pyramidal octahedral (4+1+1\*)-type stereochemistry by taking atoms N1, N2, N4 and O1 as the basal plane, and O1—Cu—N2 as the basal angle (Aduldecha *et al.*, 1991). The Cu atom is displaced from the basal plane by 0.2657 (3) Å. The basal plane has a slightly trigonal distortion with a  $\tau$  value of 0.22 (9) (Addison *et al.*, 1984). Hence, the structure can best be described as intermediate between an asymmetric *cis*-distorted octahedral geometry and a distorted square-based pyramidal

five-coordinate geometry with an additional long bond at the sixth coordinating position.

The planar pyridine rings involve dihedral angles of 27.46 (9) and 23.8 (1)° for the ligands defined by the N1/N5 and N2/N4 atoms, respectively. The crystal structure is stabilized by C—H...O and N—H...O-type hydrogen bonds, the details of which are given in Table 2.

### Experimental

The title complex was prepared by adding a boiling solution of [Cu(NO<sub>3</sub>)<sub>2</sub>].3H<sub>2</sub>O (0.48 g, 2.0 mmol) in 20 ml water to a warm solution of dpyam (0.68 g, 4.0 mmol) in 30 ml ethanol, after which CH<sub>3</sub>CH<sub>2</sub>COONa (0.38 g, 4.0 mmol) was added. After a few weeks, dark-green crystals formed.

#### Crystal data

[Cu(C<sub>3</sub>H<sub>5</sub>O<sub>2</sub>)(C<sub>10</sub>H<sub>9</sub>N<sub>3</sub>)<sub>2</sub>]-NO<sub>3</sub>

*M<sub>r</sub>* = 541.02

Monoclinic

*P*2<sub>1</sub>/*c*

*a* = 13.058 (3) Å

*b* = 8.6770 (10) Å

*c* = 21.842 (3) Å

$\beta$  = 92.790 (10)°

*V* = 2471.9 (7) Å<sup>3</sup>

*Z* = 4

*D<sub>r</sub>* = 1.454 Mg m<sup>-3</sup>

*D<sub>m</sub>* not measured

Mo K $\alpha$  radiation

$\lambda$  = 0.71073 Å

Cell parameters from 18 reflections

$\theta$  = 0.05–12.93°

$\mu$  = 0.932 mm<sup>-1</sup>

*T* = 293 (2) K

Slab

0.58 × 0.46 × 0.26 mm

Dark green

#### Data collection

Siemens P4 diffractometer

$\theta/2\theta$  scans

Absorption correction: none

5640 measured reflections

4333 independent reflections

3261 reflections with

$I > 2\sigma(I)$

*R<sub>int</sub>* = 0.021

$\theta_{\max}$  = 25.01°

*h* = -1 → 14

*k* = -1 → 10

*l* = -25 → 25

3 standard reflections

every 97 reflections

intensity decay: <3%

#### Refinement

Refinement on *F*<sup>2</sup>

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.105$

*S* = 0.984

4333 reflections

430 parameters

All H atoms refined

$w = 1/[\sigma^2(F_o^2) + (0.0685P)^2]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.497 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.346 \text{ e \AA}^{-3}$

Extinction correction: none

Scattering factors from

*International Tables for Crystallography* (Vol. C)




Fig. 1. The structure of the title compound showing 50% probability displacement ellipsoids.

Table 1. Selected geometric parameters (Å, °)

Cu—N4	1.989 (2)	Cu—N2	2.020 (2)
Cu—N1	2.005 (2)	Cu—N5	2.148 (2)
Cu—O1	2.020 (2)	Cu—O2	2.669 (3)
N4—Cu—N1	170.59 (10)	O1—Cu—N2	157.14 (9)
N4—Cu—O1	88.05 (10)	N4—Cu—N5	100.76 (10)
N1—Cu—O1	84.33 (9)	N1—Cu—N5	86.47 (9)
N4—Cu—N2	89.06 (9)	O1—Cu—N5	104.00 (9)
N1—Cu—N2	95.82 (9)	N2—Cu—N5	98.82 (9)

Table 2. Hydrogen-bonding geometry (Å, °)

D—H...A	D—H	H...A	D...A	D—H...A
C20—H20...O3	1.02 (4)	2.46 (4)	3.457 (6)	165 (3)
C2—H2...O5 <sup>i</sup>	0.90 (3)	2.52 (3)	3.291 (5)	144 (2)
N3—H1N3...O5 <sup>i</sup>	0.88 (3)	1.99 (3)	2.859 (5)	168 (3)
C3—H3...O4 <sup>ii</sup>	0.95 (4)	2.47 (4)	3.377 (9)	160 (3)
C13—H13...O5 <sup>iii</sup>	0.85 (5)	2.55 (4)	3.352 (6)	157 (4)
C14—H14...O2 <sup>iii</sup>	0.92 (4)	2.58 (4)	3.334 (4)	140 (3)
N6—H1N6...O2 <sup>iv</sup>	0.78 (3)	1.98 (3)	2.763 (4)	176 (3)

Symmetry codes: (i)  $2 - x, 1 - y, -z$ ; (ii)  $x, 1 + y, z$ ; (iii)  $x - 1, y, z$ ; (iv)  $1 - x, y - \frac{1}{2}, \frac{1}{2} - z$ .

The title structure was solved by direct methods and refined by full-matrix least-squares techniques. Atoms C22 and C23 of the propionate group showed very high disorder with unreliable C—C bond lengths. Hence, it was decided to consider the propionate group as two entities with different orientations for the bonds C21—C22 and C22—C23. The occupancies of C22 and C23 were initially refined and then fixed at 0.45 for C22A/C23A and 0.55 for C22B/C23B. The O atoms of the disordered nitrate group were divided into two sets with occupancies of 0.78 and 0.22. All the disordered atoms were refined anisotropically with C—C, N—O and O—O distances involving the disordered atoms restrained. The H atoms of C22A, C22B, C23A and C23B were fixed geometrically and allowed to ride on the atoms to which they are attached. All other H atoms were located from a difference Fourier map and refined isotropically.

Data collection, cell refinement and data reduction: XSCANS (Siemens, 1994); structure solution and molecular graphics: SHELXTL/PC (Sheldrick, 1990); structure refinement: SHELXL93 (Sheldrick, 1993); geometrical calculations: PARST (Nardelli, 1983, 1995).

The authors would like to thank the Malaysian Government and Universiti Sains Malaysia for research grant R&D No. 190-9609-2801.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: CF1199). Services for accessing these data are described at the back of the journal.

## References

- Addison, A. W., Nageswara Rao, T., Reedijk, J., van Rijn, J. & Verschoor, G. C. (1984). *J. Chem. Soc. Dalton Trans.* pp. 1340–1356.
- Aduldecha, S., Keniry, M. E., Akhter, P., Tyagi, S. & Hathaway, B. J. (1991). *Acta Cryst.* C47, 1378–1383.
- Fitzgerald, W., Murphy, B., Tyagi, S., Walsh, A., Walsh, B. & Hathaway, B. J. (1981). *J. Chem. Soc. Dalton Trans.* pp. 2271–2279.
- Hathaway, B. J. (1984). *Struct. Bonding (Berlin)*, 57, 55–118.
- Nardelli, M. (1983). *Comput. Chem.* 7, 95–98.
- Nardelli, M. (1995). *J. Appl. Cryst.* 28, 659.
- Sheldrick, G. M. (1990). *SHELXTL/PC Users Manual*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.
- Siemens (1994). *XSCANS. X-ray Single Crystal Analysis System*. Version 2.1. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Simmons, C. J., Hathaway, B. J., Amornjarusiri, K., Santar-Siero, B. D. & Clearfield, A. (1987). *J. Am. Chem. Soc.* 109, 1947–1958.

*Acta Cryst.* (1998). C54, 201–203

## $\mu$ -(2-Thioxo-1,3-dithiole-4,5-dithiolato)-bis{[tris(*p*-methoxyphenyl)phosphine]-gold(I)}

ELENA CERRADA, MARIANO LAGUNA AND M. DOLORES VILLACAMPA

*Departamento de Química Inorgánica, Universidad de Zaragoza-CSIC, Instituto de Ciencia de Materiales de Aragón, 50009 Zaragoza, Spain. E-mail: dvilla@posta.unizar.es*

(Received 12 June 1997; accepted 10 October 1997)

## Abstract

The molecule  $[\text{Au}_2(\mu\text{-C}_3\text{S}_5)\{\text{P}(\text{C}_6\text{H}_4\text{OCH}_3)_3\}_2]$  consists of two  $\text{AuPR}_3$  fragments bridged by one dmit group. Although the Au—S [2.327 (2) and 2.331 (2) Å] and Au—P [2.264 (2) and 2.265 (2) Å] bond lengths are similar to those observed in related dithiolate complexes, the intramolecular Au...S [3.013 (2) Å] and Au...Au [3.2217 (8) Å] contacts are longer.

## Comment

Dinuclear gold(I) complexes have been studied extensively during recent years (Melnik & Parish, 1986; Usón & Laguna, 1986; Mingos, 1984; Jones, 1981) because they provide a good basis for the study of weak metal–metal interactions (Schmidbaur *et al.*, 1989; Schmidbaur, 1990). Such interactions are often observed in polynuclear gold(I) derivatives and arise from relativistic effects (Pyykkö, 1988).

Interest in metal thiolate chemistry has increased because of the widespread applications of some of these compounds, *e.g.* for modelling protein sites, as anti-arthritis drugs, and in catalytic processes (Blower & Dilworth, 1987; Holm *et al.*, 1990; Wright *et al.*, 1990; Dance, 1986; Kuchn & Isied, 1980). In particular, several examples of complexes containing bridging dithiolate ligands coordinated to gold(I)— $\text{PR}_3$  units have been reported (Dávila *et al.*, 1993, 1994; Nakamoto *et al.*, 1993; Gimeno *et al.*, 1994; Cerrada *et al.*, 1996). We report here a dinuclear gold(I) derivative, (1), containing 2-thioxo-1,3-dithiole-4,5-dithiolate (dmit) as the bidentate ligand, which is an example of a so-called ‘open-ring’ complex (Dávila *et al.*, 1994).

