

**IMMUNOPHENOTYPES, CYTOKINE PRODUCTION, AND
CYTOTOXIC ACTIVITY OF NATURAL KILLER CELLS
IN HIV-1 INFECTED CASES**

SUJIN ASSAWAWITOONTIP

**A THESIS SUBMITTED IN PARTIAL FULFILLMENT
OF THE REQUIREMENTS FOR
THE DEGREE OF DOCTOR OF PHILOSOPHY
(MEDICAL MICROBIOLOGY)
FACULTY OF GRADUATE STUDIES
MAHIDOL UNIVERSITY
2005**

**ISBN 974-04-6071-2
COPYRIGHT OF MAHIDOL UNIVERSITY**

Thesis
Entitled

**IMMUNOPHENOTYPES, CYTOKINE PRODUCTION, AND
CYTOTOXIC ACTIVITY OF NATURAL KILLER CELLS
IN HIV-1 INFECTED CASES**

.....
Mr. Sujin Assawawitoontip
Candidate

.....
Prof. Pilaipan Puthavathana,
Ph.D.
Major-Advisor

.....
Prof. Kovit Pattanapanyasat,
Ph.D.
Co-Advisor

.....
Dr. Chitraporn Karnasuta
Ph.D.
Co-Advisor

.....
Associated. Prof. Prasert Auewarakul,
M.D.,Dr. med.
Co-Advisor

.....
Assoc. Prof. Rassmidara Hoonsawat,
Ph.D.
Dean
Faculty of Graduate Studies
Mahidol University

.....
Assoc. Prof. Uraivan Kositanont,
Ph.D.
Chair
Doctor of Philosophy Program
in Medical Microbiology
Faculty of Medicine Siriraj Hospital

Thesis
Entitled

**IMMUNOPHENOTYPES, CYTOKINE PRODUCTION, AND
CYTOTOXIC ACTIVITY OF NATURAL KILLER CELLS
IN HIV-1 INFECTED CASES**

was submitted to the Faculty of Graduate Studies, Mahidol University
For the degree of Doctor of Philosophy (Medical Microbiology)

on
28 April, 2005

.....
Mr. Sujin Assawawitoontip
Candidate

.....
Prof. Pilaipan Puthavathana,
Ph.D.
Chair

.....
Col. Dr. Sorachai Nitayaphan,
M.D., Ph.D.
Member

.....
Prof. Kovit Pattanapanyasat,
Ph.D.
Member

.....
Dr. Chitraporn Karnasuta
Ph.D.
Member

.....
Associated. Prof. Prasert Auewarakul,
M.D., Dr. med.
Member

.....
Assoc. Prof. Rassmidara Hoonsawat,
Ph.D.
Dean
Faculty of Graduate Studies
Mahidol University

.....
Clin. Prof. Piyasakol Sakolsatayadorn,
M.D., Grad. Dip. in Surgery, Dip. Thai
Board of General Surgery, F.R.C.S.T.
Dean
Faculty of Medicine Siriraj Hospital
Mahidol University

ACKNOWLEDGEMENTS

I would like to express my deepest appreciation and sincere gratitude to my advisor, Professor Dr. Pilaipan Puthavathana for her competent supervision, excellent guidance and generous encouragement throughout my study. I am very grateful to my co-advisors, Professor Dr. Kovit Pattanapanyasat, Associate Professor Dr. Prasert Auewarakul, Dr. Chitraporn Karnasuta, and the external committee, Col. Dr. Sorachai Nitayaphan, for their kind suggestion and critical comments and training on laboratory techniques.

I acknowledge the Royal Golden Jubilee Scholarships for Ph.D. Program, the Thailand Research Fund; and the Department of Medical Science, Ministry of Public Health, Thailand, for financial support.

I wish to especially thank Professor Dr. Surapol Suwanagool and members of the Department of Preventive and Social Medicine, Faculty of Medicine Siriraj Hospital for their kindness and providing clinical specimens. Particular note of thanks is undoubtedly expressed to all of our subjects, especially to all HIV infected persons who donated the precious blood for this study.

Special thanks to Associate Professor Dr. Uraiwan Kositanont and Associate Professor Dr. Suda Louisirochanakul for their great advice and valuable suggestion. Special appreciation is also expressed to Mrs. Rawewan Kanyok, Miss Rumporn Kulab, Miss Kannika Nateerom, Miss Phisnu Pooruk, all staff members and graduate students at the Department of Microbiology, and the Office for Research and Development, Faculty of Medicine Siriraj Hospital, Mahidol University for their kindness, help and cheerfulness throughout the course of this thesis.

I also mostly appreciate to Miss Sauwalak Thammataksin for their kindly advice on statistical analysis.

Finally, I would like to dedicate this thesis to the persons whom I can never leave out-my family, especially my dearest father and mother. I am most grateful for their great advice, unconditional love and encouragement of all my strives to reach the goal. Though my mother is no longer with us today to witness this success in person, I am certain she is now rejoining with us from the heaven above.

Sujin Assawawitoontip

IMMUNOPHENOTYPES, CYTOKINE PRODUCTION, AND CYTOTOXIC ACTIVITY OF NATURAL KILLER CELLS IN HIV-1 INFECTED CASES

SUJIN ASSAWAWITOONTIP 4037136 SIMM/D

Ph.D. (MEDICAL MICROBIOLOGY)

THESIS ADVISORS: PILAIPAN PUTHAVATHANA, Ph.D., KOVIT PATTANAPANYASAT, Ph.D., PRASERT AUEWARAKUL, M.D.,Dr.med., CHITRAPORN KARNASUTA, Ph.D.

ABSTRACT

This research emphasized the investigation of NK cell numbers and their function in HIV-1 infected cases. In addition, the lymphocyte immunophenotype reference ranges of CD3⁺, CD4⁺ and CD8⁺T cells, CD19⁺B cells, CD16⁺, CD56⁺ and CD16⁺/CD56⁺NK cells and CD4/CD8 ratios were established in a total of 125 HIV seronegative healthy Thai adults. Flow cytometric analyses demonstrated relatively lower CD4⁺T counts, but a higher NK cell count in Thais as compared to Caucasians. Our study was supported by other groups of investigators on Thai and other Mongoloid people. Statistical analyses showed that females had a significantly higher total CD3⁺T cells, but lower NK cell counts as compared to males ($p < 0.05$); and approximately 86% of NK cells carried both CD16 and CD56 molecules on the same cell. Regarding age variation, an increase of 1.1% of CD4⁺T cells per decade was seen. We also studied the changes in lymphocyte subsets in correlation with the level of CD4⁺T cell count in HIV seropositive subjects with CD4⁺T cells <14 or $\geq 14\%$. The results showed that a decrease in CD4⁺T cell count had resulted in an increase in number of CD8⁺T cells and NK cells. Meanwhile, the change on CD4⁺T cell count had no effect on level of CD19⁺B cell count.

A three color flow cytometric analysis was used to detect intracellular cytokine/chemokine production in 10 HIV seronegative and 28 HIV seropositive subjects. It was revealed upon stimulation with PMA and ionomycin that both NK and CD8⁺T cells were important sources of IFN- γ and MIP-I β ; whereas RANTES and MIP-I α were mainly produced by CD8⁺T cells. Moreover, the study demonstrated that HIV seronegative and seropositive subjects are not different in the ability to produce chemokine/cytokine.

NK cell cytotoxic activity assay was performed in 69 HIV seronegative and 107 HIV seropositive subjects by using a standard ⁵¹chromium release assay against K-562 target cells and its cytotoxic activity was expressed as lytic unit (LU20) per 10⁶PBMCs. The results showed that NK cell cytotoxic activity in HIV seronegative subjects had no difference by sex with an average LU20 of 10.5 \pm 8.4. A decrease in NK cytotoxic activity was observed in the late stage of HIV infection i.e., in subjects with CD4⁺T cells <14% as compared to HIV seronegative subjects. However, level of CD4⁺T cell count in HIV infected cases had no correlation with level of NK cell cytotoxic activity. Collective data from this study suggested that NK cells may be an alternative to CD8⁺T cells and neutralizing antibodies in the prevention against HIV disease progression. A longitudinal study is needed in order to explore this point of view.

KEY WORDS: IMMUNOPHENOTYPES/ CYTOKINE/ CYTOTOXIC ACTIVITY/ NK CELLS/ HIV
128 P. ISBN 974-04-6071-2

IMMUNOPHENOTYPES, การผลิตสารไซโตไคน์, และการทำลายเซลล์แปลกปลอม ของเซลล์ NATURAL KILLER ในผู้ติดเชื้อเอชไอวี ไทยปี 1 (IMMUNOPHENOTYPES, CYTOKINE PRODUCTION, AND CYTOTOXIC ACTIVITY OF NATURAL KILLER CELLS IN HIV-1 INFETED CASES)

สุจินต์ อัสวีทฤทธิ์ 4037136 SIMM/ D

ปร.ด. (จุลชีววิทยาการแพทย์)

คณะกรรมการควบคุมวิทยานิพนธ์ : พิไลพันธ์ พุทธิวัฒนะ, ปร.ด., โกวิท พัฒนาปัญญาสัตย์, Ph.D., ประเสริฐ เอื้อวรากุล, M.D., Dr. med., จิตรพร วรรณสุด, Ph.D.

บทคัดย่อ

โครงการวิจัยนี้ เน้นการศึกษา ชนิด จำนวนและการทำงานของเซลล์เอ็นเคในผู้ติดเชื้อเอชไอวี และการศึกษานี้ยังได้สร้างค่าอ้างอิงมาตรฐานของเซลล์ลิมโฟไซต์ชนิดอื่น ๆ อีกด้วย ได้แก่ เซลล์ $CD3^+$, $CD4^+$ และ $CD8^+$ T cells, $CD19^+$ B cells, $CD16^+$, $CD56^+$ และ $CD16^+/CD56^+$ NK cells และอัตราส่วน $CD4/CD8$ ในคนไทยที่มีสุขภาพแข็งแรง ไม่มีการติดเชื้อเอชไอวี จำนวน 125 คน การศึกษา lymphocyte immunophenotypes โดยวิธี flow cytometry พบว่าคนไทยมีปริมาณ $CD4^+$ T cells น้อยกว่าชาวคอเคเซียน ซึ่งคล้ายกับที่มีผู้เคยรายงานไว้ในคนไทยหรือชาวมองโกลอยด์ในประเทศอื่น ในแง่การศึกษาเซลล์เอ็นเค ได้พบว่าชาวไทยมีจำนวนเซลล์นี้มากกว่าชาวคอเคเซียน และเพศหญิงมีปริมาณ $CD3^+$ T cells สูงกว่าในเพศชาย แต่ในเพศชายมีปริมาณ เซลล์เอ็นเค สูงกว่าในเพศหญิง และเซลล์เอ็นเคประมาณร้อยละ 86 มีโมเลกุลของ $CD16$ และ $CD56$ อยู่บนเซลล์เดียวกัน และการศึกษาความผันแปรทางอายุ พบว่าจำนวนของเซลล์ $CD4$ จะเพิ่มขึ้นร้อยละ 1.1 เมื่ออายุมากขึ้นทุก10 ปี

งานวิจัยได้ทำการวิเคราะห์หาการเปลี่ยนแปลงจำนวนของ immune cells ชนิดต่างๆ เมื่อจำนวน $CD4^+$ T cells เปลี่ยนแปลงไป โดยทำการศึกษาในผู้ติดเชื้อเอชไอวี 2 กลุ่ม คือกลุ่มที่มี $CD4^+$ T cells <14% และ $\geq 14%$ และพบว่าเมื่อจำนวน $CD4^+$ T cells ลดลง จำนวน $CD8^+$ T cells และเซลล์เอ็นเค จะเพิ่มขึ้น แต่จำนวน B cells จะไม่มีการเปลี่ยนแปลง

การศึกษาความสามารถของเซลล์ในการสร้างสารไซโตไคน์ และคีโมไคน์ โดย flow cytometry ในเม็ดเลือดขาวของผู้ไม่ติดเชื้อเอชไอวี จำนวน 10 คน และผู้ติดเชื้อจำนวน 28 คน ด้วยการกระตุ้นด้วย PMA และ ionomycin แสดงให้เห็นว่า ทั้งเซลล์เอ็นเค และ $CD8^+$ T cells เป็นแหล่งสำคัญในการผลิตสารอินเตอร์เฟอรอนแกมมา และ MIP- β ในขณะที่เซลล์ซึ่งสร้างสารRANTES และ MIP- α ส่วนใหญ่เป็น $CD8^+$ T cells และเซลล์ของคอนปกติ และผู้ติดเชื้อเอชไอวี ไม่มีความแตกต่างกันในด้านความสามารถในการผลิตสารไซโตไคน์ และคีโมไคน์

จากการศึกษาการทำลายเซลล์แปลกปลอม โดยเซลล์เอ็นเค (NK cell cytotoxicity assay) ในคนปกติจำนวน 69 ราย และผู้ติดเชื้อเอชไอวี 107 ราย ด้วยวิธีการปลดปล่อยสารโครเมียม-51 จากเซลล์เป้าหมาย K562 โดยวัดเป็นค่า lytic unit (LU20)/ 10^6 PBMCs พบว่าค่าเฉลี่ย $LU20 \pm$ ค่าเบี่ยงเบนมาตรฐานในคนปกติคือ 10.5 ± 8.4 และค่าที่ได้ไม่มีความแตกต่างทางด้านเพศ และการศึกษาี้แสดงว่าการทำงานของเซลล์เอ็นเค ในผู้ติดเชื้อเอชไอวีลดลงเมื่อเปรียบเทียบกับคนปกติ อย่างไรก็ตาม การศึกษาในผู้ป่วยเอชไอวี ไม่พบความสัมพันธ์ระหว่างปริมาณ $CD4^+$ T cells และการทำงานของเซลล์เอ็นเค ในการทำลายเซลล์แปลกปลอม

การศึกษานี้ชี้แนะว่าเซลล์เอ็นเค น่าจะเป็นเซลล์ที่ตีในการป้องกันและควบคุมการดำเนินโรคในผู้ติดเชื้อเอชไอวี นอกเหนือไปจากเซลล์ $CD8^+$ และนิวทรอไฟลิ่ง แอนติบอดี การศึกษาคิดตามผู้ป่วยอย่างต่อเนื่อง จะให้ข้อมูลที่ชัดเจนขึ้น

CONTENTS

	PAGE
ACKNOWLEDGEMENT	iii
ABSTRACT	iv
LIST OF TABLES	ix
LIST OF FIGURES	x
LIST OF ABBREVIATIONS	xii
CHAPTER	
I INTRODUCTION	1
II OBJECTIVES	3
III LITERATURE REVIEW	4
IV MATERIALS AND METHODS	38
1. Subjects.....	38
2. Blood specimen collection and processing.....	39
3. Immunophenotype staining.....	40
3.1 Staining procedure.....	41
4. Intracellular cytokine staining.....	43
4.1 Cell stimulation for cytokine or chemokine production..	44
4.2 Intracellular cytokine or chemokine staining.....	45
5. Flow cytometric analysis.....	46
6. Evaluation of natural killer cell cytotoxic activity.....	47
6.1 Preparation of effector cells.....	47
6.2 Preparation of target cells.....	48
6.3 Cryopreservation.....	49
6.4 Thawing frozen cells from liquid nitrogen tank.....	49
6.5 Labelling of K562 target cells.....	50
6.6 Plating of effector and target cells.....	51

CONTENTS (continued)

CHAPTER	PAGE
6.7 Running NK cytotoxic activity assay.....	51
6.8 Harvest the assay plate.....	53
6.9 Data interpretation.....	53
7. Anti-HIV testing.....	54
8. Statistical analysis.....	54
V RESULTS	55
1. Leucocyte differential count in HIV seronegative, healthy subjects.....	55
2. Reference ranges of lymphocyte immunophenotype subsets with respect to sex and age.....	57
3. Determination for the influence of sex on the number of NK cells and their subpopulations.....	60
4. Reference ranges of CD4 ⁺ T cell in healthy Thai adults based on absolute number of CD4 ⁺ T cell count.....	62
5. Correlation between level of CD4 ⁺ T cell count and changes in level of CD19 ⁺ B, CD8 ⁺ T or NK lymphocytes in HIV seropositive subjects.....	64
6. Synthesis of IFN γ , RANTES, MIP-I α and MIP-I β in different lymphocyte subsets of HIV seronegative and HIV seropositive subjects.....	69
6.1 Spontaneous production of cytokine/chemokine without antigenic stimulation.....	74
6.2. Production of cytokine/chemokine upon stimulation with PMA and ionomycin.....	78
7. NK cell cytotoxicity assay.....	81

CONTENTS (continued)

CHAPTER	PAGE
VI DISCUSSION.....	88
VII CONCLUSION.....	97
REFERENCES....	99
APPENDIX.....	121
BIOGRAPHY.....	128

LIST OF TABLES

TABLE	PAGE
1. Published data on lymphocyte immunophenotypes in healthy adults of Caucasean and Mongoloid origins	13
2. Summary of CDC 1993 revised classification system for HIV infection for adolescents and adults.....	18
3. Equivalence between absolute number and percentages.....	19
4. Biological properties of chemokines.....	26
5. Phenotypic characteristics of NK cells.....	31
6. Characterization of lymphocyte immunophenotypes.....	42
7. Monoclonal antibodies to characterize immune cells expressing cytokine and chemokine.....	46
8. Leukocyte differential count of the HIV seronegative, healthy subjects.....	56
9. Lymphocyte immunophenotype reference ranges with respect to sex and age by percentages.....	58
10. Lymphocyte immunophenotype reference ranges with respect to sex and age by absolute number.....	59
11. NK cell subpopulations by percentages and absolute numbers.....	61
12. Distribution of subjects with different ranges of CD4 ⁺ Tcells.....	63
13. Correlation between level of CD4 ⁺ T cell and CD19 ⁺ B cell, CD8 ⁺ T cells or NK lymphocytes in HIV seropositive subjects.....	68
14. The background value of cytokine expressing cell without antigenic stimulation.....	74
15. Flow cytometric analyses on cytokine and chemokines producing cell upon stimulation by PMA and ionomycin.....	80
16. Determination for the amount of NK cells and their function in HIV seronegative, healthy Thais which classified by sex.....	82
17. Correlation between NK cell number and NK cytotoxic activity in HIV seronegative and HIV seropositive subjects with different ranges of CD4 ⁺ T lymphocyte count.....	83

LIST OF FIGURES

FIGURE	PAGE
1.	The basic structure of HIV particle..... 5
2.	Genomic organization and feature of the HIV genome 6
3.	Typical course of HIV infection..... 10
4.	Recognition of target cells by NK cells..... 33
5.	Mechanism of NK cells can distinguish infected from non-infected cells..... 35
6.	Correlation between level of CD4 ⁺ T lymphocyte and CD19 ⁺ B lymphocyte count in HIV seropositive subjects with; a) CD4<14% b) CD4≥14% c)total subjects..... 65
7.	Correlation between level of CD4 ⁺ T lymphocyte and CD8 ⁺ T lymphocyte count in HIV seropositive subjects with; a) CD4<14% b) CD4≥14% c)total subjects..... 66
8.	Correlation between level of CD4 ⁺ T lymphocyte and NK lymphocyte count in HIV seropositive subjects with; a) CD4<14% b) CD4≥14% c) total subjects..... 67
9.	Flow cytometric three-color analysis of IFN γ expressing CD3, CD8 and NK cells from A : unstimulated control; and B: PMA and ionomycin stimulation..... 70
10.	Flow cytometric three-color analysis of RANTES expressing CD3,CD8 and NK cells from A : unstimulated control; and B: PMA and ionomycin stimulation..... 71
11.	Flow cytometric three-color analysis of MIP-I α expressing CD3, CD8 and NK cells from A: unstimulated control; and B: PMA and ionomycin stimulation..... 72
12.	Flow cytometric three-color analysis of MIP-I β expressing CD3, CD8 and NK cells from A : unstimulated control; and B: PMA and ionomycin stimulation..... 73

LIST OF FIGURES (Continued)

FIGURE	PAGE
13. Production of intracellular cytokines, e.g., IFN γ , RANTES, MIP-I α and MIP-I β in non-activated CD3 ⁺ T cells from HIV seronegative (n=10) and seropositive subjects (n=28).....	75
14. Production of intracellular cytokines, e.g., IFN γ , RANTES, MIP-I α and MIP-I β in non-activated CD8 ⁺ T cells from HIV seronegative (n=10) and seropositive subjects (n=28).....	76
15. Production of intracellular cytokines, e.g., IFN γ , RANTES, MIP-I α and MIP-I β in non-activated NK cells from HIV seronegative (n=10) and seropositive subjects (n=28).....	77
16. Corrected percentages of IFN γ , RANTES, MIP-I α and MIP-I β expressing CD3 ⁺ T, CD8 ⁺ T and NK lymphocytes in HIV seronegative (n=10) and seropositive subjects(n=28).....	79
17. Correlation between LU20/ 10 ⁶ PBMC and NK cell numbers in a) HIV seronegative subjects b) HIV seropositive subjects with CD4<14% and c) HIV seropositive subjects with CD4 \geq 14%.....	84
18. Correlation between LU20 / 10 ⁶ PBMC and %CD16 ⁺ NK cell subpopulation in a) HIV seronegative subjects b) HIV seropositive subjects with CD4 \geq 14% and c) HIV seropositive subjects with CD4<14%	85
19. Correlation between LU20 / 10 ⁶ PBMC and %CD56 ⁺ NK cell subpopulation in a) HIV seronegative subjects b) HIV seropositive subjects with CD4 \geq 14% and c) HIV seropositive subjects with CD4<14%.....	86
20. Correlation between LU20 / 10 ⁶ PBMC and %CD4 ⁺ T cell count in a) HIV seronegative subjects b) HIV seropositive subjects with CD4<14% and c) HIV seropositive subjects with CD4 \geq 14%.....	87

LIST OF ABBREVIATIONS

%	Percentage
⁵¹ Cr	Chromium-51 radioisotope
ADCC	Antibody-dependent cell-mediated cytotoxicity
Ag	Antigen
AIDS	Acquired immunodeficiency virus
ARC	AIDS related complex
BFA	Brefeldin A
bp	Base pair
CA	California
CD	Cluster of differentiation
CDC	Center for Disease Control
Ci	Curie
cm	Centimeter
CO ₂	Carbon dioxide
cpm	Count per minute
CRF	Circulating recombinant form
CT	Connecticut
CTL	Cytotoxic T lymphocyte
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleic acid
E	Effector
e.g.	exempli gratia (for example)
EDTA	Ethylenediaminetetraacetic acid
ELISA	Enzyme-linked immunosorbent assay
ELISpot	Enzyme-linked immunospot
Env	Envelope glycoprotein
<i>env</i>	Envelope gene
<i>et al.</i>	et alii (and others)

LIST OF ABBREVIATIONS (Continued)

F	Female
FACS	Fluorescence activated cell sorter
Fc	Fragment crystallizable
Fc γ R	Fc gamma receptor
FCS	Fetal calf serum
Fig	Figure
FITC	Fluorescein isothiocyanate
g	Gram
GMCSF	Granulocyte-macrophage colony-stimulating factor
gp	Glycoprotein
HEPES	N-2 hydroxyethylpiperazine-2-ethane sulfonic acid
HIV	Human immunodeficiency virus
I	Ionomycin
i.e.	id est (that is)
IFN	Interferon
IL	Interleukin
ITAM	Immunoreceptor tyrosine-based activation motif
Kb	Kilobase
KCl	Potassium chloride
KD	Kilo Dalton
lbs	Pound (load balance system)
LGL	Large granular lymphocyte
LTR	Long terminal repeat
LU	Lytic unit
M	Male
MA	Matrix
mAb	Monoclonal antibody

LIST OF ABBREVIATIONS (Continued)

MCP	Macrophage chemoattractant protein
MD	Maryland
MHC	Major histocompatibility complex
Min	Minute
MIP-I α	Macrophage inflammatory protein-I alpha
MIP-I β	Macrophage inflammatory protein-I beta
ml	Milliliter
mM	Millimolar
MO	Missouri
mRNA	Messenger ribonucleic acid
Na ₂ ⁵¹ CrO ₄	Sodium chromate (Radioactivated isotope)
NaCl	Sodium chloride
NaOH	Sodium hydroxide
N-CAM	Neural cell adhesion molecule
Nef	Negative factor
NJ	New Jersey
NK	Natural killer cell
NY	New York
nm	Nanometer
No	Number
NS	Not significant
O.D.	Optical density
°C	Degree celsius
PBMC	Peripheral blood mononuclear cell
PBS	Phosphate buffer saline
PCR	Polymerase chain reaction
PE	Phycoerythrin
PerCP	Peridin chlorophyl protein

LIST OF ABBREVIATIONS (Continued)

PFA	Paraformaldehyde
PGL	Persistent generalized lymphadenopathy
pH	Potential of hydrogen
PMA	Phorbol 12-myristate 13-acetate
RANTES	Regulation-upon activation, normal T-cell expressed and secreted
RNA	Ribonucleic acid
rpm	Revolutions per minute
RPMI	Roswell Park Memorial Institute
RT	Reverse transcriptase
RT	Room temperature
SD	Standard deviation
SDF-1	Stromal-derived factor 1
SDS	Sodium dodecyl sulfate
sIg	Surface immunoglobulin
T	Target
T _H	T helper
U	Unit
μg	Microgram
μl	Microliter
USA	United States of America
UV	Ultraviolet
Vpr	Viral accessory gene product
WBC	White blood cell

CHAPTER I

INTRODUCTION

Determination of the distribution of immunoregulatory cells in the peripheral blood is important in the management of diseases that involve alterations in lymphocyte subpopulations. Changes in the immunoregulatory lymphocyte subsets have been found in many pathological conditions such as immunodeficiency including AIDS (1, 2), sarcoidosis (3), measles (4) and leukemia (5). The process of measuring the number of lymphocyte subsets in the whole blood sample is referred to as immunophenotyping and is usually accomplished by flow cytometric analysis. Profound changes of immune cell number occur throughout the HIV disease course. Therefore, lymphocyte immunophenotypes can be used as markers of immune status which is important to aid in monitoring of HIV disease progression and the response to therapy (6).

The high rate of antigenic variation renders HIV the ability to escape from both humoral and cell-mediated immune response. There are clear evidences that rapid mutation in the cytotoxic T lymphocyte (CTL) epitopes can lead to an escape from CTL recognition (7, 8). Moreover, during the course of HIV infection, the Nef protein downregulates MHC class I expression and may thus reduce the role of CTLs which results in viral persistence. HIV antigenic variation has no affect on NK cell recognition of the infected cells. Moreover, the downregulation of MHC class I molecules on the infected cells may render themselves more sensitive to NK cell-mediated cytotoxicity (9). Previous studies had shown that Thai people had higher frequency of NK cells as compared to the Caucasians (10). So in this study, we focused on activities of NK cells, the cell which is an important component of innate and adaptive immune system.

NK cells were defined by the capacity to bind and kill certain tumor, parasitic and virus-infected cells without prior sensitization. They can function nonspecifically in mediating lytic activity in term of antigen recognition; they also function specifically through the mechanism of antibody dependent cell cytotoxicity (11-14). After NK cells are activated by the infection, they release cytokines and chemokines that induce inflammatory responses, modulate hematopoiesis, control monocyte and granulocyte cell growth and function, and influence the type of adaptive immune responses that follow. Peripheral blood NK cell number and functional activity are in general, stable for a given individual overtime (15). However, NK cell number and function are altered in many diseases, including HIV, cancers, viral infections and autoimmune disease. In particular, NK cells are thought to be important in early host defense against viral infection, including HIV.

Variable levels of deficiency in both NK cell number and function has been shown in adult patients with AIDS (16, 17). Rapid disease progression after HIV infection also has been correlated with a decrease in NK cell activity (18). Furthermore, NK cells are important sources of cytokines and CC-chemokines which regulate immune response and suppress HIV entry or replication *in vitro* (19, 20). Vitale *et al*, (20) found a higher percentages of IFN- γ producing cells in NK cell population than in T cells in both healthy and HIV infected subjects. Moreover, Kottilit *et al*, demonstrated that NK mediated suppression of HIV replication is as potent as that of CD8⁺T cells; The suppression is mediated predominantly by secretion of CC-chemokines (19, 21).

This current research studied the lymphocyte immunophenotype subpopulations in HIV seronegative subjects and the HIV seropositive cases with different disease stages. In addition, NK cell functions: both cytolytic activity and the cytokine/ chemokine production were compared between the HIV seronegative and the HIV seropositive cases.

CHAPTER II

OBJECTIVES

This study was conducted to investigate the various properties of NK cells including lymphocyte immunophenotypes, cytokine/ chemokine production and cytotoxic activity in HIV seronegative and HIV seropositive subjects. The aims of this studies were:

1. To determine lymphocyte immunophenotype reference ranges of CD3⁺T, CD4⁺T, CD8⁺T, CD19⁺B as well as NK cells in HIV-1 seronegative, healthy Thais
2. To determine the changes in lymphocyte immunophenotype subsets in HIV seropositive subjects in accordance to level of CD4⁺ T cell count
3. To determine cytokine and chemokine production in NK cells and CD8⁺T lymphocytes
4. To measure NK cell cytotoxic activity in HIV seronegative, healthy subjects and HIV-1 seropositive subjects by using ⁵¹Cr–release assay

CHAPTER III

LITERATURE REVIEW

1. Human immunodeficiency virus (HIV)

HIV is divided into two groups denoted type 1 (HIV-1) and type 2 (HIV-2). As viewed by electron microscopy, HIV has the characteristics of a lentivirus with a cone-shaped capsid core of 80-100 nm in diameter. The surface of the HIV is characteristically made up of 72 knobs containing trimers of the envelope glycoproteins. The envelope structures are derived from a 160 kd precursor, gp160, which is cleaved into gp120 external surface envelope protein and a gp41 transmembrane protein (22, 23). The virion gp120, located on the virus surface, contain the binding site(s) for the cellular receptor(s) and the major neutralizing domains (24). The capsid core is composed of the viral p24 Gag protein. Inside this capsid are two identical RNA strands of approximately 9.2 kb in length, with which the viral RNA-dependent DNA polymerase or reverse transcriptase (RT) and the nucleocapsid proteins are closely associated (Fig. 1) (25). Diagram of HIV virion is shown in Figure 1. The HIV genome includes the structural genes *gag*, *pol*, and *env* genes and six regulatory genes, namely *tat*, *rev*, *vif*, *nef*, *vpr* and *vpu* (found in only in HIV-1) or *vpx* (found only in HIV-2) of which their products regulate the replication cycle. The *gag* sequence encodes the viral core; *pol* sequence encodes RT, integrase, and viral protease enzymes required for viral replication; and the *env* sequence encodes the envelope glycoprotein gp120 and gp41, which are required for infection of cells (26, 27). HIV genomic organization and the encoded proteins are shown in Figure 2.




Figure 1. The basic structure of HIV particle
(<http://www.critpath.org/aric/library/img002.htm>)




Figure 2. Genomic organization and feature of the HIV genome
 (<http://www.hivmedicine.com>)

Phylogenetic analysis of *env* gene sequences of HIV-1 isolates obtained from various countries led to the classification of HIV-1 into three distinct groups: M, N and O. The predominant M or major group contains 9 subtypes or clades: A-D, F-H, J and K. Similarly, six subtypes of HIV-2 (A-F) have been identified. Moreover, there is extensive variability within each subtype (28). Subtype C is found in the highest frequency among all subtypes, nearly all the identified HIV-1 subtypes (A-E, G, H, J and O) can be found in sub-Saharan Africa (25, 29, 30). Subtype B was responsible for the first homosexual epidemic, but also shown to be capable for spreading parenterally and heterosexually. Subtype B is the most common subtype found in the United States and Western Europe. While subtype E predominates in Thailand and Southeast Asia where its primary mode of transmission is heterosexual. The subtype predominated in Thailand are B' and E. Subtype E takes part of more than 90% of the cases. The Thailand subtype B' is different from the subtype B viruses in North America by the amino acids in its V3 loop (GPGQ or GPGH *versus* GPGR) (25). The subtype E in Thailand is currently renamed as circulating recombinant form (CRF) 01_AE. CRF 01_A_E is the mosaic virus of which most of the genome is derived from subtype A, but *env* gene and assignment of the LTR are derived from subtype E (31, 32).

2. Pathogenesis of HIV infection

Transmission of HIV occurs through direct contact with infected body fluids including blood products, semen, vagina and cervical secretion, amniotic fluid and breast milk via three major routes : sexual transmission, parenteral transmission and vertical transmission from mother to newborn. Sexual transmission is the great majority of HIV infections worldwide. HIV has been reported to infect a wide range of cells *in vitro*, including CD4⁺T bone marrow cells, precursor CD4⁺ bone marrow cells, CD8⁺ T cells, B cells, NK cells, eosinophils, peripheral blood dendritic cells, follicular dendritic cells, immature thymic precursor cells, Langerhans cells, megakaryocytes, astrocytes, oligodendroglia, renal epithelial cells, cervical cells, rectal and bowel mucosal cells, goblet, and columnar epithelial cells, trophoblastics cells, as well as cells and tissues from organs such as liver, lungs, salivary glands, eyes, prostate gland, testes, and adrenal gland (33, 34). Because the only cell types

that are consistently found to be infected with HIV *in vivo* are CD4⁺T lymphocytes and macrophage-lineage cells, the relevance of *in vitro* viral replication in other cell types to HIV disease is unclear at present. It is now accepted that the major viral reservoir during asymptomatic phase of HIV infection is lymph node tissue. There are two sources of HIV in lymphoid tissue, the productively infected cells (mononuclear cells, including both CD4⁺T lymphocytes and monocyte/macrophages) and the virions trapped as immune complexes on the follicular dendritic cell network of germinal center. It has been postulated that these trapped virions could act as a source of infectious virus (35-37).

The life cycle of HIV-1 can be divided into early and late phases. The early phase of HIV replication begins with the attachment of mature virion to the target cell. The entry of HIV-1 into the target cell is mediated by interaction between the viral envelope glycoprotein, the CD4 receptor, and the HIV-1 coreceptor. The primary HIV-1 coreceptors are the chemokine receptor CCR5, used by R5 HIV-1 (M-tropic, nonsyncytial inducing HIV-1), and CXCR4 used by X4 HIV-1 (T-tropic, syncytial inducing HIV-1) (38, 39). The entry process is a two-step event where an initial interaction between gp120 on envelope and CD4 molecules on cell surface elicits conformational changes in gp120 to bind the chemokine receptor binding site. Subsequent interaction of the gp120-CD4 complex with the relevant chemokine receptor molecule results in further conformational changes in the gp41 subunit which in turn facilitates the fusion event between the viral and host cell membranes, respectively, and then entry and uncoating in cytoplasm (40). Once inside the cytoplasm, the conversion of the viral RNA into double-stranded DNA provirus commences as the viral reverse transcriptase becomes active. Reverse transcriptase synthesizes a double-stranded DNA copy from the single-stranded viral RNA and generating a DNA provirus. The viral DNA migrates to and enters the host cell nucleus (a process facilitated by the HIV proteins Vpr and MA) and integrates into the host genome by the catalytic activity of integrase enzyme. Provirus can then remain latent or be active. When the host cell divides, the provirus is transmitted to daughter cells as part of host cell chromosomes. During the late phase of replication cycle, proviral DNA will be transcribed into species of viral mRNAs and later into

the viral RNA genomes . The viral components assembly into nucleocapsid which will bud from cytoplasmic membrane and generate hundred of progeny virions (41, 42).

3. Natural history of HIV infection

The immunopathogenesis of HIV infection is extremely complex. A variety of virologic and immunologic mechanisms contribute to the progressive deterioration of immune function and to progression of HIV disease to AIDS. It is now recognized that the typical case may take 8-10 years or more for AIDS to develop after seroconversion (43). A dramatic increase in the knowledge of the natural history of HIV infection has yielded insights into mechanisms of HIV disease pathogenesis. Although the course of HIV infection may vary somewhat among individual patients, a common pattern of development has been recognized (Fig. 3). The typical course of HIV infection is divided into 3 phases as follows;

Primary infection : This period is followed by the development of detectable humoral and cellular immune responses. After exposure to HIV, an infected individual becomes acutely viremic and HIV is widely disseminated during this early stage of infection, particularly in lymphoid organs (44). Moreover, within one week to three months, there is an immune response to the HIV infection and the HIV-specific immunity initiates during this stage which is associated with a dramatic decline in viremia (43, 45). However, this immunity is apparently inadequate to suppress viral replication completely, since the viruses can persist in lymph node (46).

Clinical latency : The period of inapparent infection may last for years. All patients have a gradual deterioration of the immune system, and manifest by the depletion of CD4⁺T lymphocytes (47).

Clinically apparent disease : The inevitable outcome of the progressive deterioration of the immune system that occur in most patients with HIV infection is clinically apparent disease or an acquired immunodeficiency syndrome (AIDS)-defining illness, either severe with persistent constitutional sign and symptoms or an opportunistic infection or neoplasm (48).




Figure 3. Typical course of HIV infection
 (Pantaleo G *et al.* N Eng J Med 1993; 328: 327-35)

On the basis of the duration of HIV infection and the kinetics of virologic and immunologic events, three dominant patterns of evolution of HIV disease have been described as follows (45):

a) Typical progressors : 80 to 90 % of HIV infected persons are “typical progressors”. Progression from clinically apparent disease to AIDS occurs within 8 to 10 years as the result from the continuous replication of virus in the lymphoid tissue and severe impairment of immune function (45).

b) Rapid progressors : This category occurs in 5 to 10% of HIV-infected persons. Rapid progression to AIDS occurs within 2 to 3 years after seroconversion (49). Immune response is usually defective, level of antibodies against HIV proteins and neutralizing antibody are low to absent in this group of patients (50).

c) Long-term nonprogressors : This group comprises a small percentage (5%) of infected persons who do not experience clinical progression of HIV infection, and have stable CD4⁺T cell count, absence of symptoms, and both HIV specific humoral and cell-mediated immune responses are very strong for many years (7 or more years) despite lack of therapy (51).

4. Lymphocyte immunophenotypes

Lymphocyte subpopulations comprising T, B and natural killer (NK) cells have been successfully enumerated by flow cytometry employing a panel of monoclonal antibodies to immunophenotypic surface markers. Determination of lymphocyte immunophenotypes may be useful in the disease diagnosis, prognosis, assessing the efficacy of treatment and/or searching for clues to pathogenesis. However, normal values of certain lymphocyte subsets can vary according to age (1, 10, 52-61), sex (1, 10, 53, 56, 57, 61), ethnic group (1, 10, 53, 56, 58, 60, 61), circadian rhythm (62) and methodologies (2, 63-65). Changes of lymphocyte immunophenotypes from the reference ranges were reported in several diseases of infectious and noninfectious origins, e.g. AIDS (2, 66), measles (4), cancer (67) and diabetes mellitus (68). In persons infected with HIV, level of CD4 cell count is established as a guideline for AIDS case definition or disease staging, starting anti-retroviral drugs and prophylaxis for opportunistic infections as well as a marker of treatment success (69-71).

Collective data from Asian countries including Malaysia, Hong Kong and Thailand suggested that Mongoloid people have less CD4⁺ T cells, but higher NK cells as compared to the Caucasians (Table 1)(1, 10, 56, 58). High NK cell number, therefore, may be a compensation for low CD4 cell count. Thai clinicians often come across with AIDS cases who have low CD4 cell count, but are still capable to sustain normal life activity. The published data on lymphocyte immunophenotypes in healthy adults of Caucasian and Mongoloid origins was shown in Table 1 (1, 10, 53-56, 72, 73).

Table 1. Published data on lymphocyte immunophenotypes in healthy adults of Caucasian and Mongoloid origins

Lymphocyte subset	Number of cells by % and ranges; and absolute number and ranges									
	Reichert <i>et al.</i> ^a (n=261)	Erkeler Yuksel <i>et al.</i> ^a (n=101)	Hulstaert <i>et al.</i> ^a (n=85)	Choong <i>et al.</i> ^b (n=212)	Webster <i>et al.</i> ^b (n=217)	Kam <i>et al.</i> ^b (n=208)	Vithayasai <i>et al.</i> ^b (n=150)	Webster <i>et al.</i> ^b (n=143)	Lee <i>et al.</i> ^b (M=33,F=38)	
CD3⁺	73±6.2%	72 (67-76)% 1.4 (1.1-1.7) ^c	75 (71-79)% 1.3 (1.0-1.5) ^c	67.5±8.5%	M (n=131); 65.3±8.4% F (n=86); 70.2±6.4%	69±7.7% (1.37±0.4) ^c	64.0±8.8% (1.63±0.6) ^c	69±8.5% (1.36±0.43) ^c	M; 67.5(61-73)% F; 71(67-76)% 0.8(0.7-1) ^c 1.1(0.9-1.3) ^c	
CD4⁺	43±7.5%	42 (38-46) % 0.8 (0.7-1.1) ^c	48 (43-54)% 0.7 (0.6-.98) ^c	35.5±7.8%	Both (n=154); 37.4±6.8%(0.84±0.240) ^c M (n=90); 36.1±6.7% (0.82±0.25) ^c F (n=64); 40.4±6.5% (0.93±0.22) ^c	36.4±7.5% (0.73±0.23) ^c	36.1±6.4% (0.91±0.3) ^c	40±8.0% (0.78±0.28) ^c	M;33.5(27-39)% F;37(30-43)% 0.4(0.4-0.5) ^c 0.5(0.4-0.7) ^c	
CD8⁺	33±7.5%	35 (31-40) % 0.7 (0.5-0.9) ^c	32 (28-37) %	36.8±8.5%	M (n=131); 27.6±6.6% F (n=86); 30.2±6.4%	29.7±7.2% (0.59±0.21) ^c	25.7±7.3% (0.67±0.35) ^c	26±6.9% (0.51±0.20) ^c	M; 28(24-32)% F;30(26-35)% 0.4(0.3-0.4) ^c 0.4(0.4-0.8) ^c	
CD4/CD8	1.4±0.6	1.2 (1.0-1.5)	1.8 (1.4-2.4)	1.1±0.6	1.35±0.48	1.0±0.4	1.5±0.6	1.7±0.7	M; 1.2 (0.9-1.5) F; 1.2(1.0-1.6)	
CD19⁺	14±4.2%	13 (11-16) % 0.3 (0.2-0.4) ^c	13 (11-16) % 0.21 (0.16-0.27) ^c	12.4±4.5%	14.7±4.5%	11.1±3.9% (0.22±0.11) ^c	-	14±4.8% (0.28±0.14) ^c	M; 10(9-14)% F;12(10-15)% 0.1(0.1-0.2) ^c 0.2(0.1-0.2) ^c	
NK	14±6.1%	14(10-19) % 0.3 (0.2-0.4) ^c	11 (8-15) %	17.9±8.1%	M (n=131); 20.6±8.5% F (n=85); 15.3±6.4%	19.8±8.1% (0.39±0.19) ^c	-	17±8.7% (0.34±0.21) ^c	M; 20(17-28)% F;14(10-19)% 0.3(0.2-0.4) ^c 0.2(0.1-0.3) ^c	
WBC	5.7-5.9 ^c	5.9 (4.6-7.1) ^c	5.3 (4.4-6.3) ^c	-	-	-	7.8±2.3 ^c	5.8±1.5 ^c	M; 4.8 (4.1-6.7) ^c F; 5.1(4.4-6.7) ^c	
Lymphocyte	30-33%	32 (28-39) %	31 (27-34) % 1.6 (1.3-1.9) ^c	-	2.4±0.5%	30.6±7.7% (1.98±0.53) ^c	33.8±8.1%	34±7.7% (1.97±0.59) ^c	M; 30(23-32)% F; 27(24-33)% 1.3 (1.1-1.5) ^c 1.5(1.2-1.9) ^c	

a = Caucasians ; b = Mongoloids ; c = absolute number (x 10³ cell/ µl)

5. Lymphocyte immunophenotypes in HIV infected subjects

Profound changes of immune cellular activation occur after infection with HIV and persist throughout the disease course. Prospective studies of the natural history of HIV infection have demonstrated that immunological deterioration often begins soon after HIV-1 seroconversion (74). Subsequently, most infected individuals experience a progressive loss of immunologic capacity and a decline in the number of circulating CD4⁺lymphocytes with a corresponding increase of CD8⁺T lymphocytes, leading to a drastically reduced CD4/CD8 ratio (74, 75-77). Sampalo *et al.* reported that HIV-infected patients had a striking increase in amount of B lymphocytes (78, 79). Regarding NK cell number, there are several studies with conflict results. One study had reported on the reduction of CD4⁺T cell count in associated with a decrease in number of circulating NK cells (80, 81); however, the other studies had demonstrated that numbers of circulating NK cells did not differ significantly between HIV seropositive and HIV seronegative subjects (82, 83). De Souza *et al.* (84) reported that there was no significantly different in NK cell number as compared between the HIV-seronegative and seropositive Thais, while the significant difference was demonstrated as compared between HIV-seronegative and seropositive North Americans. Moreover, Jason *et al.* (85) demonstrated an increase in the percentage of NK cells in HIV seropositive Thais as compared with HIV-seronegative subjects. Additionally, extensive prospective study by Ullum *et al.* in Europe demonstrated that the amount of NK cells was of no prognostic value for HIV disease progression (86).

6. HIV classification system for adolescents and adults

CDC has revised the classification system for HIV infection to emphasize the clinical importance of the CD4⁺T lymphocyte count in the categorization of HIV related clinical conditions. The classification system establishes mutually exclusive subgroups for which the spectrum of clinical conditions is integrated with the CD4⁺T lymphocyte count. The 1993 revised classification system for HIV infection is based on three ranges of CD4⁺T lymphocyte counts and three clinical categories and is represented by a matrix of nine mutually exclusive categories (Table 2) (87, 88).

The three CD4⁺ T lymphocyte categories are defined as follow :

Category 1 : ≥ 500 cells/ μ l

Category 2 : 200-499 cells/ μ l

Category 3 : < 200 cells/ μ l

The clinical categories of HIV infection are defined as follow:

Category A

Category A consists of more of the conditions listed below in an adolescent or adult (≥ 13 years) with documented HIV infection. Conditions listed in categories B and C must not have occurred.

- Asymptomatic HIV infection
- Persistent generalized lymphadenopathy
- Acute (primary) HIV infection with accompanying illness or history of acute HIV infection

Category B

Category B consists of symptomatic conditions in HIV infected adolescent or adult that are not included among condition listed in clinical category C and that meet at least one of the following criteria: a) the conditions are attributed to HIV infection or are indicative of a defect in cell-mediated immunity; or b) the conditions are considered by physicians to have a clinical course or to require management that is complicated by HIV infection. Examples of conditions in clinical category B include, but are not limited to:

- Bacillary angiomatosis
- Candidiasis oropharyngeal (thrush)
- Candidiasis vulvovaginal
- Cervical dysplasia/ cervical carcinoma *in situ*
- Constitutional symptoms such as fever (38.5° C) or diarrhea (lasts >1 month)
- Hairy leucoplakia, oral
- Herpes zoster (shingles), involving at least two distinct episodes
- Idiopathic thrombocytopenic purpura
- Listeriosis
- Pelvic inflammatory disease, peripheral neuropathy.

Category C

Category C includes the clinical conditions listed in the AIDS surveillance case definition. For classification purpose, once a category C condition has occurred, the person will remain in category C. The conditions included in AIDS surveillance case definition as follow;

- Candidiasis of bronchi, trachea, esophageal, or lungs
- Cervical cancer, invasive
- Coccidioidomycosis, disseminated or extrapulmonary
- Cryptococcosis, extrapulmonary
- Cryptosporidiosis, chronic intestine (>1 month's duration)
- Cytomegalovirus disease (other than liver, spleen, or nodes)
- Cytomegalovirus retinitis (with loss of vision)
- Encephalopathy, HIV-related
- Herpes simplex: chronic ulcer (>1 month's duration); or bronchitis, pneumonitis, or esophagitis
- Histoplasmosis disseminated or extrapulmonary
- Isosporiasis, chronic intestine (1 month's duration)
- Kaposi's sarcoma
- Lymphoma, Burkitt's (or equivalent term)
- Lymphoma, primary of brain

- *Mycobacterium avium* complex or *M. kansasii*, disseminated or extrapulmonary
- *Mycobacterium tuberculosis*, any site (pulmonary or extrapulmonary)
- *Mycobacterium*, other species or unidentified species, disseminated or extrapulmonary
- *Pneumocystis carinii*
- Pneumonia, recurrent
- Progressive multifocal leukoencephalopathy
- *Salmonella* septicemia, recurrent
- Toxoplasmosis of brain,
- Wasting syndrome due to HIV.

Table 2. Summary of CDC 1993 revised classification system for HIV infection for adolescents and adults

CD4⁺T cell categories	Clinical categories		
	(A) Asymptomatic, acute HIV or PGL	(B) Symptomatic, not (A) or (C) conditions	(C) AIDS-indicator conditions
1) >500 cells / μ l	A1	B1	C1
2) 200-499 cells / μ l	A2	B2	C2
3) < 200 cells / μ l	A3	B3	C3

7. Equivalences for CD4⁺T lymphocyte count by absolute numbers and percentages

Enumeration of CD4⁺T lymphocytes in term of absolute number is more accurate than that presented in term of percentage upon repeated measurements. However, data correlating natural history of HIV infection with the CD4⁺T cell percentages have not been as consistently available as the data on absolute CD4⁺T lymphocyte count. Therefore, the CDC classification system emphasizes the use of CD4⁺T lymphocyte counts by absolute number, but the use of CD4⁺T lymphocyte by percentages are also accepted. Estimation for an equivalence between absolute number and percentages of CD4⁺T lymphocyte count is shown in Table 3 (87).

Table 3. Equivalence between absolute numbers and percentages

CD4 ⁺ T cell category	CD4 ⁺ T lymphocytes / μ l	% CD4 ⁺ T lymphocyte
(1)	≥ 500	≥ 29
(2)	200-499	14-28
(3)	< 200	<14

8. Immune response to HIV infection

The immune system in HIV infection has been divided into innate and adaptive component, each with a different function and role (89, 90). Innate immunity is the first line of defense against invading infection. The mechanism of innate immunity exist before an encounter with the viruses and are rapidly activated by the viruses before the development of adaptive immune response. The innate immune system consist of epithelial barriers, circulating effector cells (granulocytes, macrophages, NK and dendritic cells) and circulating effector proteins (e.g. complements) (26). The NK cells display at least two effector functions that can contribute to control of infection : first, they can kill infected cells ; second, they are a rich source of inflammatory cytokines, especially $IFN\gamma$ and chemokines such as RANTES, $MIP-1\alpha$ and $MIP-1\beta$ with antiviral activity. Macrophages and dendritic cells also play a key role in the early control of infection by secreting IL-2 and IL-18 that can induce $IFN\gamma$, to elicit direct and indirect antiviral effects (91).

The adaptive immune response consists of cellular and humoral immunity which are important in controlling the HIV infection. Humoral immunity directs antibody response within 6 to 9 weeks after infection to a variety of HIV antigens. However, little evidences show that the antibodies have any beneficial effect in limiting the disease (26). The cellular immune response play a vital role in HIV pathogenesis. The studies of the long-term nonprogressors, and the exposed yet uninfected persons have suggested that certain aspects of cellular immunity may protect against HIV infection or slow disease progression (41). Two aspects of cellular immune response are the cytotoxic T-cells and the response that release cytokines (92-94) and lyse target cell expressing foreign antigens; and T-helper cells which is responsible directly or indirectly for induction of a wide array of lymphoid and lymphoid cell functions. These effects include activation of macrophages; induction of cytotoxic T cells; NK cells and B cells; and secretion of a variety of soluble factors that induce growth and differentiation of lymphoid cells and affect hematopoietic cells (95). Cellular immunity generally occurs in parallel with the humoral (antibody) response, although the two can be separated in certain circumstances. The infection usually evoke both arms of immune, which broadly differ in their function: the cellular immune response

controls the infection and the humoral response prevents further infection with the same agent (96). There is clear evidence that both during acute and chronic phases of HIV infection, rapid mutation in epitopes which are recognized by HIV-specific CTL may occur. In addition, the Nef protein can down regulate the molecules of MHC class I expressing on the surface of infected cell, thus lead to a reduction in the role of CTLs (97). These events result in HIV escape from the immune response. However, HIV antigenic variation has no affect on NK cell activity and MHC Class I down regulation may render themselves more sensitive to NK cell-mediated cytotoxicity.

9. Cytokines and chemokines

Cytokines are important mediators generated during immune response and are produced by a variety of activated cells, such as CD8⁺T cells, T_H0 and T_H1 CD4⁺T cells and NK cells (98, 99). They play an important role in regulating the immune response and determining resistance or susceptibility to disease. Chemokines constitute a large family of small cytokines with four conserved cysteins linked by disulfide bonds. When chemokines bind to HIV coreceptors, they are potent and selective inhibitors of HIV infection. Cytokines are a family of proteins that mediated many of responses of innate and adaptive immunity (26). The same cytokines may be produced by many cell types and individual cytokine often acts on diverse cell types. They are synthesized in response to inflammation or antigenic stimuli and usually act locally, in an autocrine or paracrine fashion, by binding to high affinity receptors on target cells (27). Thus, cytokines serve many functions that are critical to host defense against the pathogens and provide link between innate and adaptive immunity (26). Depending on the nature of the infecting virus, the infected cells may be triggered to produce anti-viral cytokines and chemokines that inhibit one or more steps in the viral life cycle (90), thereby limiting the extent of the infection. The complex network of cytokines involved in inflammatory and immunoregulatory responses has an important role in several components of the pathogenesis of HIV infection and AIDS (100).

Recently, several cytokines, e.g., IFN- γ , IFN- α , IFN- β , IL-2, IL-12 and IL-16 have been demonstrated to profoundly affect the state of HIV expression, overall spreading, and suppress HIV infection. In addition, some chemokines e.g., RANTES, MIP-I α , MIP-I β , SDF-1, MCP-1, MCP-2, MCP-3 and IP-10 function as the HIV-suppressive factors by interacting with chemokine receptors. When viruses infect cells, production of IFN is induced, and consequently, viral replication is inhibited. Interferon is believed to be part of the body's first line of defense against viral infection and to be the first cytokine to be recognized. Interferon regulate humoral and cellular immunity and has broad cell growth regulatory activities (27, 98, 100-102).

IFNs are cytokines that play a complex and central roles in the resistance of mammalian host to pathogens. IFNs, although best known for their anti-viral properties, are potent regulators of cell growth and have immunomodulatory activity (103-105). IFN is a family of related proteins that may be grouped into two types, I and II. Type I IFNs are composed of two distinct groups of proteins called IFN- α and IFN- β which are synthesized by many cell types after viral infection. Double-stranded RNA is a potent inducer of IFN synthesis (26, 104). There are two distinct families (I and II) of IFN- α . IFN- α (I) is 166 amino acid long whereas IFN- α (II) has 172 amino acids. Major sources of IFN- α are lymphocytes, monocytes, macrophages. IFN- β is related to IFN- α with 30% amino acid sequence homology and source of IFN- β are fibroblast and some epithelial cells. IFN- α and IFN- β seem to share the same receptor and have very similar biological activities (26, 106).

The action of type I IFNs are to protect cells against viral infection and promote cell-mediated immunity against intracellular microbes by inhibiting viral replication, increase expression of class I MHC molecule, stimulates the development of T_H1 cells and inhibit the proliferation of many cell types (26, 106). IFN- α is the first cytokine for which a role as an endogenous suppressor factor of HIV replication has been reported. Addition of the neutralizing antibody to IFN- α into the PBMC culture derived from HIV infected cases has been shown to increase both the levels of HIV replication and the frequency of positive viral (107, 108). Defective production of

IFN- α *in ex vivo* has been described in HIV infected individuals (109). While some reports demonstrated high level of IFN- α during HIV infection (110, 111). Regarding IFN- β , significant changes in the number of HIV proviral copies PBMCs have not been found after treatment of HIV infected patients with this agent (112). It implies that IFN- β may not play significant role in HIV infection.

Type II IFN or IFN- γ is the principle macrophage-activating cytokine and serves critical functions in innate immunity and in specific cell-mediated immunity. It has some anti-viral activity, but it is not a potent anti-viral IFN, and it functions mainly as an effector cytokine of immune response. IFN- γ is a homodimeric protein produced by NK, CD4⁺T_{H0}, and T_{H1}, and CD8⁺T cells. In innate immunity, NK cells secrete IFN- γ in response to recognition of unknown components of microbes or in response to IL-12; in this setting IFN- γ functions as a mediator. In adaptive immunity, T cells produce IFN- γ in response to antigen recognition, and this is enhanced by IL-12. The sequence of reactions involving IL-12 and IFN- γ is central to cell-mediated immunity against intracellular microbes. IFN- γ functions by ;1) providing the T lymphocytes and NK cells to activate macrophages to kill phagocytosed microbes (104, 113) stimulation of expression of class I and class II MHC molecules (114); 3) acting as costimulator on antigen presenting cells (26); 4) promoting the differentiation of naïve CD4⁺ T cells to the T_{H1} subset and inhibiting the proliferation of T_{H2} (114, 115); 5) acting on B cells to promote switching to certain IgG subclasses; 6) activating of neutrophils (104); and 7) stimulating to the cytolytic activity of NK cells (116).

Several studies have described an elevation of IFN- γ production in HIV-infected individual; however, there are conflicting reports on the status of IFN- γ production by cells isolated from HIV infected patients. Highly increased constitutive production of IFN- γ has been observed in PBMCs of asymptomatic HIV-infected patients with CDC group II/III as well as patients who have progressed to CDC group IV, as compared to healthy subjects (117, 118). However some studies demonstrated that PBMCs from HIV-infected individuals produce less IFN- γ in response to the stimuli (119, 120). Vitale *et al.* demonstrated that both in HIV uninfected and HIV-infected subjects, IFN- γ was produced from NK cells in a higher percentages than T cells (20).

Chemokines are 8-10 kd peptides that are the product of distinct genes clustered to human chromosomes 4 and 17. Chemokines regulate to selectively promote the rapid adhesion, chemotaxis and activation of leukocyte effector subpopulations (121, 122). In addition, some are able to regulate the proliferative potential of hematopoietic progenitor cells, endothelial cells, and certain types of transformed cells (123, 124). Chemokines act via a family of G-protein-coupled receptors and these types of receptor are usually large ligand (60 amino acids). Chemokine can be divided into two groups, α and β by the arrangement of the first two of four conserved cysteines. Members of the β chemokines possess the first two conserved cysteines which are adjacent. The gene for the β chemokines is on human chromosome 17, and they generally activate monocytes, lymphocytes, basophils and eosinophils. The major human chemokine β family includes MIP-1 α , MIP-1 β and RANTES which have been identified as the major M-tropic HIV suppressive factors (125). The biological properties of MIP-1 α , MIP-1 β and RANTES have been characterized in Table 4 (104, 106, 123).

β Chemokines act by competitive binding to a coreceptor, CC chemokine receptors (CCR5), that is also required by macrophage tropic HIV-1 strains as a coreceptor for entry into the host cell. This receptor-ligand binding renders CCR5 unavailable to the virus either by competitively blocking virus-cell interactions or by causing down-regulation from host cell surface ((126-130).

The α chemokine, also known as the CXC chemokines, contain a single amino acid between the first and second cystein residues. The gene that instructs synthesis of the α chemokine is on human chromosome 4. The major α chemokine is stroma cell-derived cofactor (SDF-1) which is an extremely efficacious chemoattractant for T lymphocytes and play a prominent role in B cell development. SDF-1 is a 352 amino acid proteins, mostly closely related to the receptor for IL-8 (131, 132). In addition, α chemokine (SDF-1) was shown to specifically inhibit CXCR4 mediated infection by T cell line tropic HIV-1. SDF-1 has been found to be responsible for competitive binding to a coreceptor CXCR4 which was identified as a specific coreceptor essential for entry of T-tropic HIV-1 into target cells(132).

The accumulating data have suggested that HIV infection that augmented production of RANTES, MIP-1 α and MIP-1 β may limit viral replication *in vivo* (133) and provided evidence for a significant inverse correlation between level of these chemokines and a reduction of virus burden *in vivo*. A specific immune response involving a high production of β chemokine by T cells seems to play role of protection in exposed uninfected individuals (134) and may contribute to the control of viral replication in long-term nonprogressors (135). Several studies demonstrated that PBMCs from HIV-1 infected, long term nonprogressors often produce higher levels of β -chemokines upon antigen stimulation *in vitro* than do PBMCs from rapid progressors (136, 137); and higher production of MIP-1 β by PBMCs has been associated with an asymptomatic status and a decreased risk of disease progression (138). Infection without disease progression for longer than 10 years, even though occurs in a minority of HIV-infected people, may be attributable from chemokine production, chemokine receptor polymorphisms including host factors (139).

The previous studies have demonstrated that CD8⁺ T cells, CD4⁺ T cells and macrophages, as well as several other cell types, secrete cytokines and CC-chemokines productions (140, 141). In addition, a few studies also showed the roles of NK cells for cytokine and chemokine production.

Table 4. Biological properties of chemokines

Chemokine	Percent identity to MIP-1α	Receptor	Source	Targets/ actions	Major effect
β Chemokine; MIP-1α	100	CCR1, 3, 5	B, T and NK lymphocytes, mast cells, fibroblasts, macrophages, monocytes	Neutrophil and monocyte chemotaxis, neutrophil and macrophage activation, stem cell suppression, potentiation of GM-CFU stimulation by GM-CSF, PGE-independent endogenous pyrogens Monocyte, NK, T, basophil and dendritic cell chemoattraction, monocytes, NK, T, and dendritic cell chemoattraction	Competes with HIV-1 for binding to coreceptors, anti-viral defense, promotes T _{H1} immunity
MIP-1β	67	CCR1, 3, 5	B, T and NK lymphocytes, macrophages, monocytes, neutrophils	Inhibits MIP-1 α action on stem cells, potentiates GM-CFU stimulation by GM-CSF, inhibits macrophage activation by MIP-1 α , monocytes, NK, T, and dendritic cell chemoattraction	Competes with HIV-1 for binding to coreceptors
RANTES	46	CCR1, 3, 5	T and NK lymphocytes, platelets, endothelial cells	Monocyte, granulocytes, memory T lymphocytes, NK, basophil, eosinophil, and dendritic chemoattraction	Competes with HIV-1 for binding to coreceptors, degranulates basophils, activate T cells, chronic inflammation
α Chemokine; SDF-1		CXCR4	T and NK lymphocytes, dendritic cells, bone marrow stroma cells	B cell maturation and lymphopoiesis, hematopoiesis, transendothelial chemotaxis on lymphocyte, monocyte but not monocyte, potent mononuclear cell attractant	Competes with HIV-1 for binding to coreceptors

10. Cytokine and chemokine assays

Several novel methods have been developed which facilitate the detection of cytokine and chemokine production by immune cells. Cytokine and chemokines can be assayed by: 1) enzyme-linked immunosorbent assay (ELISA) for quantifying the amount of secreted cytokine and chemokine in serum or cell culture supernatant; 2) determination of specific mRNA in chemokine and cytokine producing cells using polymerase chain reaction (PCR) (142); 3) enumeration of activated cells secreting cytokine by using enzyme-linked immunospot (ELISpot) (143, 144). However, these methods are time consuming by requiring long activation times and they are labour intensive (93). In addition, these methods can not specify cell type which produce cytokine/ chemokine because the same cytokine/ chemokine may be produced from several cell types. For instance, IFN γ may be produced from CD8 $^+$, CD4 $^+$ and NK cells.

A highly sensitive flow cytometric technique, that is based on the ability to quantitate and phenotypically characterize cytokine expressing cells simultaneously without prior sorting has been developed (145). This technique has the advantage of rapid determining intracellular cytokine production by activating the cells with mitogen in a short incubation time. Subsequently, addition of brefeldin A (BFA) to block the secretory pathway, a process that disrupts intracellular Golgi-mediated transport and allows the cytokines to accumulate and yield an enhanced signal for the detection by flow cytometry. This method has been applied to detect several types of cytokines in every discreted cellular population, and also phenotypically characterize the cytokine producing cells in various diseases such as HIV disease, autoimmunity and allergy. It is also possible to detect intracellular cytokine production for evaluation of human immune response, vaccine efficacy, drug and therapeutic modalities.

11. Characteristics of NK cells

NK cells are one component of the innate immune system and have the ability to both lyse target cells and provide an early source of immunoregulatory cytokines. Human NK cells are defined phenotypically by their expression of CD16, CD56 and lack of expression of CD3; and comprise about 15% of all lymphocytes (146). However Thai people have higher number of NK cells as compared to the Caucasians (10). NK cells are central in the innate immune response against tumor, parasites and cells infected with virus and intracellular bacteria (147, 148).

Previous studies have provided strong evidence that NK cells belong to the lymphocyte lineage. NK cells originated and differentiated in bone marrow (149, 150). Most of the characteristic of NK cells are summarized in Table 5 (86). NK cells are more closely related to T cells than B cells, because neither T nor NK cells express surface immunoglobulin (sIg), and they share expression of many cell surface molecules and functional attributes, including effector mechanism (151). NK cells can develop independently from mature antigen specific T or B lymphocytes and do not productively rearrange or express $\alpha\beta$ or $\gamma\delta$ T-cell receptor on their cell surface (147). NK cells were originally identified by their ability to spontaneously mediate lysis of certain susceptible tumor cells (148).

Morphologically, NK cells are characterized as large granular lymphocytes (LGLs) with a kidney-shaped nucleus and prominent azurophilic cytoplasmic granules. NK cells are members of lymphoid lineages. They are larger than most B or T cells with ranging in diameter from 12 to 15 μm . The majority of human NK cell activity is mediated by CD3^- , CD16^+ , CD56^+ lymphocytes. Due to their expression of CD16 (immunoglobulin Fc receptor, $\text{Fc}\gamma\text{RIII}$) these cells ($\text{CD16}^{\text{bright}}$) exhibit high level of antibody-dependent cellular cytotoxicity (ADCC) activity which is an arm of adaptive immunity. However CD16^- NK cells also exist (152).

Approximately, 90% of human NK cells have low-density expression of CD56 (CD56^{dim}) and express high level of $\text{Fc}\gamma$ receptor III ($\text{Fc}\gamma\text{RIII}$, $\text{CD16}^{\text{bright}}$), whereas about 10 % of NK cells are $\text{CD56}^{\text{bright}} \text{CD16}^{\text{dim}}$ or $\text{CD56}^{\text{bright}} \text{CD16}^-$ (151). CD56 is a 140 kd molecule, an isoform of the neural cell adhesion molecule (N-CAM),

found on neural tissue and some tumor cell. The CD56 is not expressed by other hematopoietic cells or lymphocytes (151, 153). The CD16 is a 70-kd glycoprotein of Ig superfamily and is expressed as a transmembrane protein on the majority of human peripheral blood NK cells (154). CD16 is the low affinity Fc γ RIII on the surface of NK cells, which binds to antibody-coated (opsonized) targets and signals through associated subunits containing an immunoreceptor tyrosine-based activation motif (ITAM) on target cell to direct ADCC (155). Now, there are ample evidence to suggest that these NK-cell subsets have unique functional attributes and therefore, distinct roles in the human immune response. The CD56^{dim} CD16^{bright} NK cell subset is more naturally cytotoxic and express higher levels of Ig-like NK receptors and Fc γ RIII than the CD56^{bright} NK cell subset. By contrast, the CD56^{bright} subset has the capacity to produce abundant cytokines following activation of monocytes, but has low natural cytotoxicity and is CD16^{dim} or CD16⁻ (146).

NK cells are derived from pluripotent hematopoietic stem cells. A committed precursor for T and NK cells expressing Fc γ RIII has been demonstrated both in human and mice (156). NKT cells are a subset of lymphocytes that are characterized by the co-expressing of the NK cell receptor, CD161 and a single large invariant TCR α chain (V α 24J α Q in the human and V α 14J α 28 in the mouse). In contrast to conventional T lymphocytes, the NKT TCR does not interact with peptide antigen presented by classical MHC class I or II molecule, but instead it recognizes glycolipids presented by CD1d (non classical-MHC-class-I like molecule). They have the unusual property of being readily stimulated through their TCR as a result of presentation of the nonpeptide antigen, α -galactosylceramide (α -GalCer), by CD1d molecules (157-160). Phenotypically, NKT cell are either CD4⁺CD8⁻ or CD4⁻CD8⁺ and this T cell population represents a major fraction of the mature T cell in thymus, nearly 50 % of α / β TCR⁺ T cells in liver and up to 5% of splenic T cells, but are rare in lymph node (161). As yet, little information is available on the tissue distribution of human invariant NKT cells. They constitute approximately 0.02-0.2 % of peripheral blood T cell compartment in both males and females (162).

NKT cells are capable of rapidly producing both T_H1 (e.g. IFN- γ) and T_H2 type cytokine (e.g. IL-4, IL-5, IL-10) upon triggering suggesting a potential role of NKT cells in immunoregulation (163). Nevertheless, NKT cells have now been shown to control various immune responses, including autoimmune, allergic, antitumor, and antimicrobial immune response. In HIV infection, depletion of NKT cells upon infection with HIV-1 results from the high susceptibility to infection of CD4 and CCR5 coexpressing NKT cells, potentially in combination with activation-induced cell death during persistent immune activation (162-164). Since a subset of NKT cells is CD4⁺T and also highly express level of the chemokine receptor CCR5 and CXCR6 but lower level of CXCR4 and CCR7. Unutmas *et al.* found that CD4⁺NKT cells are indeed higher susceptible to R-tropic but less so to X-4 tropic virus and caused rapid destruction of this subset *in vitro* (165, 166). Recent findings have shown that NKT cells in PBMCs of HIV-1 infected individuals are dramatically reduced, compared with healthy donors and they established a positive correlation with high viral load and lower CD4⁺ NKT numbers (164, 166).

Innate immunity effector cells, including NK cells recognize the presence of different pathogens mainly through a recently identified family of 10 genetically invariant receptors, the Toll-like receptors (TLR 1-10), capable of recognizing distinct molecular components of microbes (167). In particular, TLR3 is the specific receptor for dsRNA, a common intermediate in the reproductive cycle of many viruses. TLR3 expression on human NK cells and the ability of dsRNA to augment NK cell functions has been reported (168), but the mechanisms enabling NK cells to functionally respond to this product of viral infections are mostly unknown. Sivori *et al.* demonstrated that NK cells are able to respond to stimuli acting on different TLRs expressed by myeloid dendritic cells. Thus, after culture in the presence of the microbial products such as dsRNA (virus-derived) as well as CpG (of bacterial origin) can induce peripheral blood NK cells activation characterized by; 1) *de novo* expression of activation markers such as CD69 and CD25; 2) release of various cytokines including IFN- γ and TNF- α ; 3) up-regulation of antitumor cytotoxicity, and; 4) acquisition of cytotoxicity against immature dendritic cells (169).

Table 5. Phenotypic characteristics of human NK cells (170)

Characteristics	
1. General morphology	
Size	Large, 12-15 μm (mouse, 8-10 μm)
Cytoplasmic granules	+
Kidney-shaped nucleus	+
Adherence	-
Phagocytosis	-
2. Histochemical properties	
Acid phosphate	+
Non-specific esterase	+
β -Glucuronidase	+
Peroxidase	-
TdT	-
3. Cell surface antigens	
CD2	$\geq 75\%$ (rat 30%)
CD3	$\leq 5\%$
CD4	$\leq 1\%$
CD5	$\leq 5\%$
CD7	80% (Not tested in rodents)
CD8	25% (1% in mouse, >90% in rat)
CD11	>90% (Not tested in rodents)
CD16 (Fc γ RIII)	50-90% (ADCC is functionally weak in rodents)
CD56	10 % of human NK (not reliable marker for mouse)
Fc μ R	<1%
sIg	<1%
Ia	10% (<1% in mouse)
Asialo GM1	>90%

Discordance between species indicated by parenthesis +, >90% of the cells are positive for this characteristic; -, negative for the indicated characteristics

12. NK cell activity

The principal physiologic role of NK cells is to get rid of cells harbouring foreign antigens. NK cells are activated by recognition of three types of targets: 1) antibody coated cells (ADCC); 2) cells infected by viruses and some intracellular bacteria; and 3) cell lacking class I MHC molecules (Fig. 4) (26).

ADCC process requires three components: 1) target cell expressing pathogen's antigens on the surface; 2) antibodies of the IgG isotype to the target antigen, and 3) effector cells bearing Fc gamma receptor (FcγR) (13). Therefore, ADCC occurs only when the target cell is coated with antibody, and free IgG in plasma neither activates NK cells nor competes effectively with cell-bound IgG for binding to FcγRIII (13, 26, 152, 156, 171, 172). Engagement of FcγRIII by antibody-coated target cells activates the NK cells to synthesize and secrete cytokines such as IFN-γ, as well as discharge the content of their granules, which contain several proteases and nuclease as well as perforin, a molecule which polymerizes to form pores in the membrane of the target cell (171, 173). If NK cells are to mediate direct cytotoxicity against cells infected with viruses or other pathogens, they must have some mechanism for distinguishing the infected from the uninfected cells (27). Since all nucleated cells normally express MHC class I molecules, many viruses have evolved means to inhibit the expression of these molecules in infected cells (26, 27). The downregulation of MHC class I on surfaces of the infected cells can protect themselves from lysis by virus specific CD8⁺ cytotoxic T lymphocytes, but these viral infected cells may render themselves more sensitive to NK cell mediated cytotoxicity (Fig. 4) (148, 172, 174).




Figure 4. Recognition of target cells by NK cells. (A) NK cell recognizes antibody-coated targets by the FcγRIII receptor; (B) NK cell binds to target by adhesion molecules and other unknown ligands; (C) NK cell is activated by target cell lacking class I major histocompatibility complex (MHC) molecules; the activating receptors involved in the response to class I-deficient target are not known (26)

The possible mechanisms whereby NK cells distinguish infected from non-infected cells was shown in Figure 5. A proposed mechanism of NK cell recognition is shown. NK cells can use several different receptors that signal them to kill infected cells, including lectin-like receptors that recognize carbohydrate on self cells. However, another set of receptors, called Ly49 in the mouse and killer inhibitory receptors (KIR) in the human, recognize MHC class I molecules and inhibit killing by NK cells by overruling the actions of the killer receptors. This inhibitory signal is lost when host cells do not express MHC class I and perhaps also in cells infected with virus, which might inhibit MHC class I expression or alter its conformation. Normal cells respond to IFN- α and β by increasing levels of MHC class I expression, making them resistant to activated NK killing. The infected cells can fail to increase MHC class I expression, making them targets for activated NK cells.




Figure 5. Mechanism by which NK cells can distinguish infected from non-infected cells (27)

13. NK cell response to HIV infection

The role of NK cells in controlling HIV replication and spread of the viruses has not been fully delineated. Several studies have correlated high NK cell activity with the reduced susceptibility of certain individuals to HIV infection (175); and loss of NK cell activity and frequency had been correlated with HIV disease progression, particularly in individuals with opportunistic infections (14, 176, 177). As the result of the rapid mutation in HIV-specific CTL epitopes, HIV escape mutants were discovered. Moreover, Nef protein downregulates MHC class I molecules expression on HIV infected cells and escape from CTL destruction (9, 86, 178). Antigenic variation of HIV may render themselves more sensitive to NK cells than CTL mediated cytotoxicity. So NK cell may thus serve as a second line of defense against HIV-infected cells evading the T cell mediated immune response as well as a protective mechanism against an array of secondary opportunistic infections and malignant manifestation of HIV infection.

Peripheral blood mononuclear cells (PBMCs) with or without AIDS or AIDS related complex (ARC) exhibit poor NK cytotoxic activity in the ^{51}Cr release assay (80, 85, 179-182). While Ironson *et al.*, demonstrated that there was no difference in NK cell cytotoxic activity between the HIV seronegative subjects and asymptomatic HIV seropositive cases with low CD4^+ T cell count (83). In particular, NK cells are thought to be important in early host defense against viruses, including HIV. Studies of NK cells in HIV infection have shown variable levels of deficiency in both NK cell number and function. Ullum *et al.*, demonstrated rapid disease progression after HIV infection also has been correlated with decrease NK cell activity (86).

Scott-Algara *et al.* reported that the defective mechanism underlying the observed defective NK cytotoxic activity could be the results from lack or decreased frequency of effector cells, inability to recognize and bind the target cells, failure to be activated for the release and/or synthesis of NK cytotoxic factors (182). The HIV specific ADCC mechanism also declined with disease progression due to both a defect in the ability to generate anti-HIV antibodies required for ADCC as well as the diminished lytic capacity of the NK cells (13, 149).

14. Measurement of NK cell cytotoxic activity

Several studies on NK cytotoxic activity in HIV infection were based on the standard protocol ^{51}Cr -release assay. NK cell cytotoxic activity was expressed as number of lytic unit (LU20) / 10^6 PBMCs. The LU20 / 10^6 PBMCs is defined as lytic activity of effector cells which are required to lyse 20% of 5×10^3 target cells. LU20 / 10^6 PBMCs was calculated based on the equation (183) :

$$\text{Number of lytic units}/10^6 \text{ PBMC} = 10^6 / T \cdot X_p$$

T = number of target cells i.e., 5000 cells/well

X_p = E : T ratio required to lyse p% of target cells

p = reference lysis level (i.e., 20%)

As 5 E : T ratios were performed, it will be easier to make calculation by LU computer program (184).

Determination of NK cytotoxic activity was performed by using fresh PBMCs without prior *in vitro* sensitization. The K562 (erythroleukemic cell line) is used for the standard target cells for measuring in this assay. Cytotoxicity of NK cells can be measured by assaying the release of ^{51}Cr from labeled target cells. ^{51}Cr is thought to bind to cytoplasmic proteins of the cell. Following NK cell attack on the target cell, these proteins are released to the surrounding area, providing a marker for the cytoplasmic events of apoptosis. PBMCs usually are used as a source of immune effector (NK) cells. After incubation of the effector and tumor target cells, the radioisotope was relatively release from the lysed of target cells into the supernatant. Measurement of releasing radioisotope is performed by using automated radioisotope counter (184).

CHAPTER IV

MATERIALS AND METHODS

1. Subjects

The subjects comprised 3 major groups in 3 different studies as follows:

1) Determination of lymphocyte immunophenotype reference ranges : A total of 125 HIV seronegative healthy subjects including 67 males and 58 females at age range of 18-55 years (mean \pm SD = 30.9 \pm 10.1, median = 27.0 years)

2) Determination of intracellular cytokine/ chemokine production :

2.1 A total of 10 HIV seronegative healthy subjects including 2 males and 8 females at age range of 23-31 years (mean \pm SD = 26.0 \pm 2.5, median = 25.5 years)

2.2 A total of 28 HIV seropositive subjects with different ranges of CD4⁺T lymphocyte count as follows:

2.2.1 nine HIV seropositive subjects with CD4<14% including 9 males and 1 female at age range of 24-40 years (mean \pm SD= 30.1 \pm 5.1, median= 28.0 years)

2.2.2 19 HIV seropositive subjects with CD4 \geq 14% including 11 males and 8 females at age range of 21-53 years (mean \pm SD= 31.9 \pm 10.0, median= 28.0 years)

3) Determination of NK cell cytotoxic activity :

3.1 A total of 69 HIV seronegative, healthy subjects including 38 males and 31 females at age range of 21-55 years (mean \pm SD = 32.8 \pm 10.0, median = 30.0 years) and 66 HIV-1 subtype E and 3 HIV-1 subtype B

3.2 A total of 107 HIV seropositive subjects categorized on the percentages of CD4⁺T lymphocyte count as follows:

3.2.1 46 HIV seropositive subjects with CD4<14% including 26 males and 20 females at age range of 18-50years (mean \pm SD= 29.7 \pm 6.9, median= 28.5 years)

3.2.2 61 HIV seropositive subjects with $CD4 \geq 14\%$ including 20 males and 41 females at age range of 19-53 years (mean \pm SD= 29.4 \pm 7.80, median= 27.0 years)

All of HIV infected subjects attended the HIV Clinic, Department of Preventive and Social Medicine, Faculty of Medicine Siriraj Hospital during the period between July 1999-June 2002. All were naïve for treatment with anti-retroviral drug at the time of enrollment. The subjects understood the objective of this study and signed in the consent form.

2. Blood specimen collection

For each subject, 25 ml of the venous blood were collected in EDTA vacutainer tubes for NK cell cytotoxicity assay; and the other 2 ml were collected in a sodium heparinized vacutainer tube (Becton-Dickinson Vacutainer System, Franklin Lakes, NJ). EDTA blood sample was further divided into 3 aliquots for the following purposes:

2.1 Determination for white blood cell count and differential cell count by automated machine (Beckman-Coulter STKS, CA, USA.) at the Department of Clinical Pathology, Faculty of Medicine Siriraj Hospital

2.2 Determination for lymphocyte subsets

2.3 Determination for intracellular cytokine and chemokine expressing cells

3. Immunophenotype staining

Lymphocyte subpopulation present in EDTA blood sample were enumerated by two color staining of the surface markers using matched combination of fluorescein isothiocyanate and phycoerythrin conjugated mAbs (Becton Dickinson). The lymphocyte subsets determined were total CD3⁺T lymphocytes, CD3⁻CD19⁺B lymphocytes, CD3⁺CD4⁺T helper/inducer lymphocytes, CD3⁺CD8⁺T suppressor/cytotoxic lymphocytes, CD3⁻CD16⁺ and/or CD56⁺ NK lymphocytes, CD3⁻CD16⁺ and CD3⁻CD56⁺NK lymphocyte subpopulation. The characterization of lymphocyte immunophenotype was shown in Table 6. All lymphocyte subsets were measured by flow cytometer (FACScan; Becton Dickinson Immunocytometry System San Jose, CA, USA) using a panel of mAbs and reagent as follow:

1. Simultest LeucoGATE reagent (CD45/CD14, Anti-Hle-I/Leu-M3, Catalog No. 340040) which contains FITC labeled anti-CD45 mAb and PE labeled anti-CD14 mAb
2. Simultest Control reagent $\gamma 1/ \gamma 1$ (IgG₁/IgG₁, Catalog No. 349526) which contains FITC labeled anti-IgG₁ mAb and PE labeled anti-IgG₁ mAb
3. Simultest CD3/CD19 reagent (Leu-4/12, Catalog No. 349211) which contain FITC label anti-CD3 mAb and PE labeled anti-CD19 mAb
4. Simultest CD3/CD4 reagent (Leu-4/3a, Catalog No. 340043) which contains FITC labeled anti-CD3 mAb and PE labeled anti-CD4 mAb
5. Simultest CD3/CD8 reagent (Leu-4/2a, Catalog No. 340044) which contains FITC labeled anti-CD3 mAb and PE labeled anti-CD8 mAb
6. Simultest CD3/CD16+CD56 reagent (Leu-4/11c+19, Catalog No. 340042) which contains FITC labeled anti-CD3 mAb and PE labeled anti-CD16 and CD56 mAb
7. CD3-FITC reagent (Leu-4 FITC, Catalog No. 349201) which contains FITC labeled anti- CD3 mAb
8. CD16-PE reagent (Leu-11c, Catalog No. 347617) which contains PE labeled anti-CD16 mAb
9. CD56-PE reagent (Leu-19, Catalog No. 347747) which contains PE labeled anti-CD56 mAb

10. FACS Lysing Solution 10X concentration (Catalog No. 349202) which contains reagent for lysing the erythrocytes under gentle hypotonic condition and preserving the leucocytes

3.1 Staining procedure

1. Set up eight 12x75 mm tubes for one blood sample
2. Gently mix thoroughly the EDTA blood and place each 50 μ l of EDTA blood into each tube
3. Add 5-10 μ l of each pair of mAb into each tube as follow:
 - 3.1 Simultest CD45/14
 - 3.2 Simultest γ 1/ γ 1 (IgG₁/IgG₁)
 - 3.3 Simultest CD3/CD19
 - 3.4 Simultest CD3/CD4
 - 3.5 Simultest CD3/CD8
 - 3.6 Simultest CD3/CD16+CD56
 - 3.7 CD3 and CD16
 - 3.8 CD3 and CD56
4. Vortex gently and incubate the tubes for 15 to 30 minutes at room temperature in the dark
5. Add 2 ml of 1X FACS Lysing solution into each tube. Immediately vortex gently and incubate for 10 minutes at room temperature in the dark
6. Centrifuge the tubes at 200xg for 5 minutes
7. Aspirate the supernatant by leaving approximately 50 μ l of residual fluid; avoid disturbing the pellet
8. Add 2 ml of phosphate buffer saline (PBS) into each tube. If samples contain too many red blood cells, add 2 to 3 drops of distilled water before adding PBS immediately
9. Vortex the tubes gently and centrifuge at 200xg for 5 minutes
10. Aspirate the supernatant and then resuspend the cell pellet in 0.5 ml of 1.0% paraformaldehyde in PBS

The stained cells were analyzed by a flow cytometer using SimulSET Software (Becton Dickinson), within 24 hours after staining.

Table 6. Characterization of lymphocyte immunophenotypes

Reagents (FITC/PE)	CD number (FITC/PE)	Functional identity of relevant subsets
LeucoGATE	CD45/CD14	Establishment of an optimal lymphocyte gate for immunophenotyping of erythrocyte lysed whole blood (lymphocyte monocytes and granulocytes)
Isotype control	IgG1/IgG1	Direct conjugate control for non-specific staining
Leu-4/Leu-12	CD3/CD19	T (CD3+CD19-) and B (CD3-CD19+) lymphocytes
Leu-4/Leu-3	CD3/CD4	CD4+ helper/inducer T cells
Leu-4/Leu-2	CD3/CD8	CD8+ cytotoxic/suppressor T cells
Leu-4/Leu-11+ Leu-19	CD3/CD16+CD56	NK cells (CD3-CD16+ and/or CD56+)
Leu-4/Leu-11	CD3/CD16	NK cell subsets (CD3-CD16+)
Leu-4/Leu-19	CD3/CD56	NK cell subsets (CD3-CD56+)

4. Intracellular cytokine staining

This technique is used for determination of cytokine production in heterogeneous cell population based on the ability to detect a single cytokine expressing cell. This rapid method determines the intracellular cytokine production by activating PBMCs with mitogen in a short incubation time (4 hours). Subsequently, addition of brefeldin A to block the secretory pathway, allows the accumulation of cytokines that can be detected by flow cytometer using a panel of mAbs. Reagents No.1-12 were purchased from Becton Dickinson, San Jose, CA, USA, and No.13-15 were purchased from Sigma Chemical Co., St Louis, MO, USA

1. IgG₁-PE (Isotype control, Catalog No. 349043) which contains PE labeled anti-IgG₁ mAb
2. CD3-FITC reagent (Leu-4 FITC, Catalog No. 349201) which contains FITC labeled anti-CD3 mAb
3. CD3-PerCP reagent (Leu-4 PerCP, Catalog No. 347344) which contains PerCP labeled anti-CD3 mAb
4. CD8-PerCP reagent (Leu-2a PerCP, Catalog No. 347314) which contains PerCP labeled anti-CD8 mAb
5. CD16-FITC reagent (Leu11a FITC, Catalog No. 347523) which contains FITC labeled anti-CD16 mAb
6. CD56-FITC reagent (NCAM16.2, Catalog No. 340410) which contains FITC labeled anti-CD56 mAb
7. Anti-human IFN- γ mAb-PE (Catalog No. 18905A) which contains PE labeled anti-human IFN γ mAb
8. Anti-human RANTES mAb-PE (Catalog No. 20975A) which contains PE labeled anti-human RANTES mAb
9. Anti-human MIP-1 α mAb-PE (Catalog No. 20955A) which contains PE labeled anti-human MIP-1 α mAb
10. Anti-human MIP-1 β mAb-PE (Catalog No. 23855A) which contains PE labeled anti-human MIP-1 β mAb
11. FACS Lysing Solution 10X concentration (Catalog No. 349202)
12. FACS Permeabilizing Solution 10X concentration (Catalog No. 340457)

which contains the solution for the permeabilization of lymphocyte membranes prior to intracellular immunofluorescence staining with mAb

13. Ionomycin Calcium salt (I) 1 mg (Catalog No. I-0634)
14. Phorbol 12-myristate 13-acetate (PMA) 1 mg (Catalog No. P-8139)
15. Brefeldin A (BFA) 5 mg (γ -4-Dihydroxy-2[6-hydroxy-1-heptenyl]-4-cyclopentane-crotonic acid λ -lactone, Catalog No. B-7651)

4.1 Cell stimulation for cytokine or chemokine production

1. Dilute 500 μ l of whole heparinized blood in 500 μ l RPMI 1640 (GIBCO, Paisley Scotland, USA) in a 15 ml conical centrifuge tube
2. Add 25 μ l of 25 ng/ml PMA, 20 μ l of 1 μ g/ml I and 10 μ l of 0.5 mg/ml BFA for cell stimulation into each tube (Tubes of unstimulated samples are set up in parallel by adding BFA but without PMA and I)
3. Mix the tube by vortex and incubate at 37 °C in 5 % CO₂ humidified atmosphere for 4 hours
4. Add 100 μ l of 20 mM EDTA into each tube
5. Vortex the tube and incubate at room temperature for 15 minutes
6. Add 4.5 ml of FACS Lysing solution into each tube
7. Gently mix the tube and incubate at room temperature for 10 minutes
8. Wash with 2% fetal bovine serum (FBS) (GIBCO) in PBS as washing buffer
9. Centrifuge at 500xg for 5 minutes
10. Discard the supernatant
11. Add 2.5 ml of FACS permeabilizing solution into each tube and incubate at room temperature for 10 minutes
12. Wash with washing buffer by centrifuge at 500 x g for 5 minutes and discard the supernatant
13. Resuspend the cell pellet with 0.5 ml of washing buffer

4.2 Intracellular cytokine and chemokine staining

The stimulated cells were stained using triple colors as the following protocol:

1. Set up nine 12x75 mm tubes for 1 blood sample
2. Add 5 μ l of each mAb into each tube as follow;
 - 1.1 CD16 plus CD56-FITC/IgG₁-PE/CD3-PerCP
 - 1.2 CD3-FITC/IFN- γ -PE/CD8-PerCP
 - 1.3 CD3-FITC/RANTES-PE/CD8-PerCP
 - 1.4 CD3-FITC/MIP-1 α -PE/CD8-PerCP
 - 1.5 CD3-FITC/MIP-1 β -PE/CD8-PerCP
 - 1.6 CD16 plus CD56-FITC/ IFN- γ -PE/CD3-PerCP
 - 1.7 CD16 plus CD56-FITC/ RANTES -PE/CD3-PerCP
 - 1.8 CD16 plus CD56-FITC/ MIP-1 α -PE/CD3-PerCP
 - 1.9 CD16 plus CD56-FITC/ MIP-1 β -PE/CD3-PerCP
3. Add 100 μ l of stimulated cells into each tube
4. Incubate all samples in room temperature for 30 minutes in the dark
5. Wash the stained cell sample with washing buffer
6. Centrifuge at 500xg for 5 minutes
7. Aspirate the supernatant and resuspend in 1% paraformaldehyde in PBS
8. Analyse by flow cytometer using CellQuest software (Becton

Dickinson)

Table 7. Monoclonal antibodies to characterize immune cells expressing cytokine and chemokine

Tube No.	Mab conjugated with FITC or PE or PerCP	Functional identity of relevant subset
1	CD16+CD56/IgG ₁ /CD3	Isotype-matched negative control
2	CD3/IFN- γ /CD8	IFN- γ expressing CD3 and CD8 T cells
3	CD3/RANTES/CD8	RANTES expressing CD3 and CD8 T cells
4	CD3/MIP-1 α /CD8	MIP-1 α expressing CD3 and CD8 T cells
5	CD3/MIP-1 β /CD8	MIP-1 β expressing CD3 and CD8 T cells
6	CD16+CD56/ IFN- γ /CD3	IFN- γ expressing CD3 T and NK cells
7	CD16+CD56/ RANTES /CD3	RANTES expressing CD3 T and NK cells
8	CD16+CD56/ MIP-1 α /CD3	MIP-1 α expressing CD3 T and NK cells
9	CD16+CD56/ MIP-1 β /CD3	MIP-1 β expressing CD3 T and NK cells

5. Flow cytometric analysis

Samples were analysed using a FACScan or FACSort flow cytometer (Becton Dickinson) which had been calibrated by using CaliBRITE Beads and Auto Comp Software (Becton Dickinson). Two color immunophenotype analyses were accomplished with the SimulSET software (Becton Dickinson). The LeucoGATE mAb (CD45/CD14) identified lymphocytes, monocytes and granulocytes by light scatter gating; and the purity of at least 95% lymphocytes was obtained for immunophenotype analyses. The SimulSET software was programmed to enumerate 15,000 lymphocyte cells in a measurement.

Three color flow cytometric analyses was performed with CELLQuest software (Becton Dickinson). The results were analysed by setting up the templates which used the plots and histograms for each sample collection. Lymphocytes were initially gated by forward and side scatter, secondary gating was set on the basis of staining with isotypic control mAb. The quadrant markers used to determine the percentage of lymphocytes producing cytokine and chemokine were set by basing on unstimulated sample. The percentages of antigen specific cytokine and chemokine response were calculated by subtracting the values of the stimulated sample with that of the unstimulated match sample.

6. Evaluation of natural killer cell cytotoxic activity

NK cell cytotoxic activity was measured by a standard chromium 51 (^{51}Cr) release assay. Determination of NK cell cytotoxic activity was performed by using fresh PBMCs without prior *in vitro* sensitization. The cell line K-562 (kindly provided by Dr. Chitraporn Karnasuta, Armed Forces Research Institute of Medical Sciences) is used as the standard target cells.

6.1 Preparation of effector cells

Effector cells were PBMCs isolated from EDTA blood samples by standard ficoll hypaque (Robbins Scientific, Sunnyvale, CA, USA) density gradient as follows:

1. Centrifuge the tube of EDTA blood at 200xg for 10 minutes at room temperature and remove the plasma
2. Dilute the packed blood with 2 volumes of steriled PBS, for example, resuspend 4 ml of blood in 4 ml of PBS
3. Slowly overlay 8 ml of diluted blood on to 4 ml of ficoll-hypaque in a 15 ml conical centrifuge tube and centrifuge at 500xg for 20 minutes at room temperature with the brake off
4. After centrifuging, carefully collect PBMCs in white band at interface. Upper part of the tube contains plasma, platelets, and PBS, and the lower part contains ficoll-hypaque, red blood cells and granulocytes
5. Wash the cells once with PBS and twice with RPMI 1640 with antibiotic by centrifuging at 200xg for 10 minutes at room temperature with brake on

6. Resuspend separated PBMCs in growth medium (RPMI 1640 plus 10% fetal bovine serum) to the concentration 5×10^6 cells in 5 ml. In order to enrich lymphocyte by depletion of the adherent macrophages from PBMC population, remove adherent macrophages by placing the PBMC suspension in a 25 cm² polystyrene tissue culture flask (Costar, Corning, NY)

7. Incubate the cells at 37 C in 5% CO₂ incubator for overnight. The flask should be laid down flat in the incubator

8. After incubation, collect the nonadherent lymphocyte and dispense into 15 ml centrifuge tube

9. Centrifuge the tube at 1,500rpm for 10 minutes and decant the supernatant

10. Resuspend cell pellet in growth medium and adjust to the concentration of 5×10^6 cells/ml

6.2 Preparation of target cells

The cell line K562, derived from a patient with chronic myelogenous leukemia in blast crisis, is the standard target cells for measuring NK cytotoxic activity. This target cells are the continuous cell line and could multiple indefinitely. The K562 cells were cultered in 10% FCS RPMI growth medium and incubated at 37°C in a CO₂ incubator. In the assay, the K562 target cells should be in log phase and more than 95% viable. The viability of the cells were determined by tryphan blue exclusion. They were fed with 10% FCS RPMI growth medium one day prior to the test. One million K562 cells are sufficient for setting up assay for 3-4 subjects.

The K562 cell lines should be subpassaged twice a week. The cell line could be continuous maintained or cryopreserved in the freezing medium, and stored in liquid nitrogen.

6.3 Cryopreservation of cells

Cells should be frozen at concentration about 10×10^6 cells in 1 ml freezing medium; and their viability should be more than 95%. RPMI 1640 containing 20% FCS and 10% dimethyl sulfoxide (DMSO)

1. Resuspend the cells to be frozen in 0.5 ml of cold RPMI 1640 supplemented with 40% FCS in a cryovial (Costar, Corning, NY)
2. Gradually add dropwise of 0.5 ml cooled RPMI 1640 containing 20% DMSO into the vial of cells with gently mix
3. Immediately place cryovials in a storage foam box in a -70°C freezer for overnight, then transfer the vial into a liquid nitrogen tank

Notice: Cap of the vial should be the type of inner thread with O-ring. Volume of cell suspension added into the cryotube should not be higher than the indicated line on the vial wall.

6.4 Thawing frozen cells from liquid nitrogen tank

1. Transfer cryovial from the liquid nitrogen tank to a -70°C freezer for 1 hour
2. Place cryovial into an ice box with closed lid for 1 hour in order to let liquid nitrogen evaporated. If the screw cap was immediately loosen while liquid nitrogen gas still remain at the joint between the cap and the body of vial, the cryovial might be bursted out.
3. After thawing, put the cell suspension into a tissue culture flask, and add with 10 ml of growth medium
4. Incubate the flask overnight at 37°C in a CO_2 incubator
5. Replace with new growth medium and observe daily for growth

With this freezing-thawing protocol, viability of the reconstituted K562 cells is always higher than 95%.

6.5 Labeling of K562 target cells

1. Place 10^6 K562 cells in log phase in a 15 ml conical tube
2. Centrifuge the tube at 1500 rpm for 10 minutes at 25°C
3. Discard the supernatant and resuspend the cell pellet with 1 ml of growth medium
4. Add $200\ \mu\text{Ci}$ of $\text{Na}_2^{51}\text{CrO}_4$ (Amersham Pharmacia Biotech UK limited, Buckingham shire, England) per 1×10^6 K562 cells (the volume of ^{51}Cr needed is adjusted according to the decay of the isotope)
5. Incubate the target cells for 1-1.5 hour at 37°C with shaking every 15 min to ensure uniform labeling
6. Wash the ^{51}Cr -labeled K562 target cells with 10 ml of 1x RPMI 1640 by centrifugation at 1500 rpm for 10 minutes for 3 times
7. Resuspend the cells in 1 ml of growth medium
8. Count cells by using a hemacytometer, and adjust to the cell concentration of 5×10^6 viable ^{51}Cr -labeled K562 cells/ml. This is the stock required for plating the target cells

6.6 Plating of effector and target cells

The test was run in a triplicate in the microtiter U-shape (Costar, Corning, NY). Effector cells were mixed with target cells at the effector (E): target (T) ratios of 100:1, 50:1, 12.5:1 and 6.25:1

The U-bottom microtiter plate was set as below:

		Subject 1			Subject 2			Subject 3			Subject 4		
E : T		1	2	3	4	5	6	7	8	9	10	11	12
100 : 1	A												
50 : 1	B												
25 : 1	C												
12.5 : 1	D												
6.25 : 1	E												
	F	Spontaneous release											
	G												
	H	Maximum release											

6.7 Running NK cytotoxic activity assay

1. Add 100 μ l of 10% FCS RPMI media into the wells B1-B3, C1-C3, D1-D3, E1-E3 and F1-F3
2. Add 200 μ l of effector cells at concentration of 10^6 cells/ml in triplicate into wells A1-A3 of the first row
3. Mix cells in wells A1-A3 thoroughly, and perform serial two fold dilution by transfer 100 μ l of the cell suspension from the wells A1-A3 to the wells B1-B3. The remaining amount of the effector cells in the wells A1-A3 is 5×10^5 cells/well

4. Mix and transfer 100 μ l of the cell suspension from the wells B1-B3 to the wells C1-C3. The remaining amount of the effector cells in the wells B1-B3 is 2.5×10^5 cells/well

5. Mix and transfer 100 μ l of the cell suspension from the wells C1-C3 to the wells D1-D3. The remaining amount of the effector cells in the wells C1-C3 is 12.5×10^4 cells/well

6. Mix and transfer 100 μ l of the cell suspension from the wells D1-D3 to the wells E1-E3. The remaining amount of the effector cells in the wells D1-D3 is 6.25×10^4 cells/well

7. Mix and discard 100 μ l of the cell suspension from the wells E1-E3. The remaining amount of the effector cells in the wells E1-E3 is 3.125×10^4 cells/well

8. Add 100 μ l of 10% FCS RPMI media into the wells F1-F3, while add 100 μ l of sterilized 10% sodium dodecyl sulfate (SDS) (Sigma, MO, USA) into the wells H1-H3

9. Add 100 μ l of ^{51}Cr -labeled K562 target cells at concentration of 5×10^4 cells/ml into every well of rows A-F and H, column 1-3 (for one subject), to give effector: target cells ratios of 100:1 in row A, 50 :1 in row B, 25:1 in row C, 12.5:1 in row D and 6.25:1 in row E. The wells F1-F3 represent spontaneous release, while the wells H1-H3 represent maximum release

The experiment was done in triplicate and the component of the test wells were as below:

Row A; column 1-3 = Effector 5×10^5 cells + target 5×10^3 cells

Row B; column 1-3 = Effector 2.5×10^5 cells + target 5×10^3 cells

Row C; column 1-3 = Effector 1.25×10^5 cells + target 5×10^3 cells

Row D; column 1-3 = Effector 6.25×10^4 cells + target 5×10^3 cells

Row E; column 1-3 = Effector 3.125×10^4 cells + target 5×10^3 cells

Row F; column 1-3 = 100 μ l of the growth medium + target 5×10^3 cells

Row H; column 1-3 = 100 μ l of 10% SDS (detergent) + target 5×10^3 cells

10. Incubate the reaction plate for 4 hours at 37 $^{\circ}\text{C}$ in a CO_2 incubator to allow close contact between effector cells and the target cells

6.8 Harvesting the assay plate

1. Carefully remove the plate from the incubator and carefully transfer 35 μ l of culture supernatant from each well of the reaction plate to a scintillation fluid coated 96 well Luma plate (Packard Bioscience Company, CT, USA) in the corresponding well position
2. Decontaminate the infectiousness of the test wells by adding 15 μ l of 100% commercial bleach to every well of the Luma plate and incubate overnight in order to let the fluid evaporated
3. Measure radioactivity present in the Luma plate by Beta-counter (Packard Bioscience Company, Model Top Count, CT, USA)

6.9 Data interpretation

Incubation of effector cells with ^{51}Cr -labeled K562 target cells will lead to target cell lysis and a release of the cytosol content together with ^{51}Cr into the culture supernatant. Measurement of releasing radioisotope is performed by using automated radioisotope Beta-counter. Percent lysis of each kind of effector cell was calculated based on the equation below.

$$\% \text{ Lysis} = \frac{\text{mean test cpm} - \text{mean spontaneous cpm}}{\text{mean max cpm} - \text{mean spontaneous cpm}} \times 100$$

cpm = counts per minute (average cpm of triplicate wells)

test cpm = cpm released by the target cells in the presence of the effector cells

spontaneous cpm = cpm released by the target cells in the absence of the effector cells

maximum cpm = cpm released by the target cells in the presence of 10% SDS

NK cell cytotoxic activity was expressed as number of lytic unit (LU_{20}) / 10^6 PBMC. The $LU_{20} / 10^6$ PBMCs is defined as lytic activity of effector cells which are required to lyse 20% of 5×10^3 target cells (184). $LU_{20} / 10^6$ PBMC was calculated by using computer software, kindly provided by Dr. Chitraporn Karnasuta.

7. Anti-HIV testing

Subjects enrolled in this study were tested for HIV-1 infection. Sera or plasma samples were assayed for anti-HIV antibody by ELISA (Organon Teknika BV, Netherland) and gel partical agglutination (Serodia-HIV-FUJIREBIO Inc, TOKYO, Japan). Reaction samples were confirmed by three screening test of different principles. The assay were performed as the diagnostic service of the Division of Virology, Faculty of Medicine Siriraj Hospital.

8. Statistical analysis

Data of each lymphocyte subset and percentage of cytokine and chemokine expressing cells and NK cytotoxic activity obtained from all subsets were analysed for difference between HIV-1 seronegative and seropositive subjects. Komogorov Smirnov test was used to verified the pattern of distribution of a dataset in order to determine the goodness of fit. When normal distribution of dataset was obtained ($p \geq 0.05$), parametric analyses would be employed, i.e., determination for sex difference of lymphocyte subsets would be performed by an independent t-test. On another hand, if distribution of dataset was not normal ($p < 0.05$), non-parametric assay would be employed, i.e., the difference of lymphocyte subsets would be determined by Mann whitney-U test. In addition, age variation was determined by simple linear regression test and the correlation between lymphocyte subset or lytic unit vs NK cell number was determined by pearson correlation coefficients.

CHAPTER V

RESULTS

1. Leukocyte differential count in HIV seronegative, healthy subjects

A total of 125 HIV seronegative, healthy subjects were determined for the reference ranges of leukocyte differential counts by using an automated machine (Beckman-Coulter Counter STKS, CA, USA).

The results demonstrated that number of white blood cell (WBC) in males was significantly higher than in females (independent t-test; $p=0.005$); but there was no statistically significant difference in percentage of lymphocyte or monocyte count between both genders ($p\geq 0.05$) as analysed by independent t-test and Mann Whitney U-test, respectively (Table 8).

Table 8. Leukocyte differential count of the HIV seronegative, healthy subjects

Cell type	Sex	N	Number of cell count		
			Mean \pm SD	Median	95% Reference range
WBC x 10 ³ cells/ μ l	M	67	7.3 \pm 2.0*	7.1	6.8 - 7.8
	F	58	6.4 \pm 1.5*	6.2	6.0 - 6.8
	Both	125	6.9 \pm 1.9	6.6	6.6 - 7.2
%Lymphocyte	M	67	30.7 \pm 7.7	30.0	28.8 - 32.6
	F	58	31.1 \pm 8.0	30.5	29.0 - 33.2
	Both	125	30.9 \pm 7.8	30.0	29.5 - 32.3
%Monocyte	M	67	7.2 \pm 2.5	7.0	6.6 - 7.8
	F	58	7.2 \pm 2.3	7.0	6.6 - 7.8
	Both	125	7.2 \pm 2.4	7.0	6.8 - 7.6

* There was statistically significant difference as number of cell count of both sexes was compared (independent t-test; p=0.005)

2. Reference ranges of lymphocyte immunophenotype subsets with respect to sex and age

Reference ranges for lymphocyte subsets were determined in 125 healthy Thai adults by flow cytometric analysis using a panel of monoclonal antibodies conjugated with either fluorescein isothiocyanate or phycoerythrin to identify cell-specific surface antigens. The reference range of each lymphocyte subpopulation as presented in term of percentage and absolute number are shown in Tables 9 and 10, respectively.

The result demonstrated that males had higher percentages and higher absolute numbers of NK cells than females (independent t-test; $p=0.008$ and 0.003 , respectively). Discordant results between percentages and absolute number were observed with $CD3^{+}T$ cell count. As absolute number was partly derived from WBC count, percentage value may better yield the actual result of comparison. Our study showed that females had significantly higher percentages of $CD3^{+}T$ cells (independent t-test; $p=0.018$).

There was no discrepancy by sex as tested on various parameters: $CD4^{+}T$, $CD8^{+}T$, $CD19^{+}B$ lymphocytes and $CD4/CD8$ ratio (independent t-test; $p\geq 0.05$). Regarding the variation in values of all lymphocyte subsets by age as analysed by simple linear regression analysis, the difference were shown only in $CD3^{+}CD4^{+}T$ cells with an increase of 1.1% and 53.8 cells/ μ l per decade (Tables 9 and 10).

Table 9. Lymphocyte immunophenotype reference ranges with respect to sex and age by percentages

Lymphocyte immunophenotype	Sex	N	Mean	SD	Median	95% Reference range	Sex difference	Age variation
CD3⁺ (total T cells)	M	67	63.4	8.4	64.0	61.3-65.4	3.4 %	
	F	58	66.8	7.4	66.0	64.9-68.7	F > M	NS
	Both	125	65.0	8.1	65.0	63.5-66.4	(p=0.018)	
CD3⁻ CD19⁺ (B cells)	M	67	12.1	3.8	12.0	11.2-13.0		
	F	58	13.0	4.9	13.0	11.7-14.3	NS	NS
	Both	125	12.5	4.3	12.0	11.8-13.3		
CD3⁺ CD4⁺ (T helper/inducer)	M	67	31.7	6.0	32.0	30.2-33.2		1.1 %
	F	58	32.6	5.8	32.5	31.0-34.1	NS	per
	Both	125	32.1	5.9	32.0	31.1-33.1		decade
CD3⁺ CD8⁺ (T suppressor/cytotoxic)	M	67	26.9	7.1	26.0	25.2-28.6		
	F	58	28.7	7.5	29.0	26.8-30.7	NS	NS
	Both	125	27.7	7.3	28.0	26.5-29.0		
CD4/CD8 ratios	M	67	1.3	0.5	1.2	1.2-1.4		
	F	58	1.3	0.5	1.1	1.1-1.4	NS	NS
	Both	125	1.3	0.5	1.2	1.2-1.4		
CD3⁻ CD16⁺ and/or CD56⁺ (Total NK cells)	M	67	23.2	9.1	21.0	21.0-25.5	4.0 %	NS
	F	58	19.2	7.5	19.0	17.2-21.2	M > F	
	Both	125	21.4	8.6	20.0	19.8-22.9	(p=0.008)	

NS = no significant difference

Table 10. Lymphocyte immunophenotype reference ranges with respect to sex and age by absolute numbers

Lymphocyte immunophenotype	Sex	N	Mean	SD	Median	95% Reference range	Sex difference	Age Variation
CD3⁺ (total T cells)	M	67	1,355.5	401.9	1,307.0	1,257.4-1,453.5		
	F	58	1,308.1	421.7	1,198.0	1,197.3-1,419.0	NS	NS
	Both	125	1,333.5	410.2	1,269.0	1,260.9-1,406.1		
CD3⁻ CD19⁺ (B cells)	M	67	268.9	124.5	252.0	238.5-299.3		
	F	58	257.4	128.2	234.5	223.7-291.1	NS	NS
	Both	125	263.6	125.9	245.0	241.3-285.8		
CD3⁺ CD4⁺ (T helper/inducer)	M	67	689.8	240.8	663.0	631.1-784.5		53.8
	F	58	633.8	198.7	584.5	581.6-686.1	NS	cells/ μ l
	Both	125	663.8	223.1	624.0	624.3-703.3		per decade
CD3⁺ CD8⁺ (T suppressor/cytotoxic)	M	67	587.5	229.6	542.0	531.5-643.6		
	F	58	570.3	277.4	527.5	497.4-643.3	NS	NS
	Both	125	579.6	252.0	531.0	534.9-624.2		
CD4/CD8 ratios	M	67	1.3	0.5	1.2	1.2-1.4		
	F	58	1.3	0.5	1.1	1.1-1.4	NS	NS
	Both	125	1.3	0.5	1.2	1.2-1.4		
CD3⁻ CD16⁺ and/or CD56⁺ (Total NK cells)	M	67	533.3	352.1	455.0	447.4-619.1	152.3	
	F	58	381.0	202.3	366.5	327.8-434.1	cells/ μ l	NS
	Both	125	462.6	301.0	403.0	409.3-515.9	M > F (p=0.003)	

NS = no significant difference

3. Determination for the influence of sex on the number of NK cells and their subpopulations

In addition to determination of the total number of NK cells (cells which expressed either $CD3^+CD16^+CD56^+$ or $CD3^+CD16^+CD56^-$ or $CD3^+CD16^-CD56^+$), we also enumerated NK cell subset which expressed CD16 molecules (either with or without co-expression of the CD56 marker), and NK subset which expressed CD56 molecules (either with or without co-expression of CD16 marker). Based on the reference ranges for both sexes of 19.8-22.9% for total NK cells, 18.1-20.9% for the CD16 NK cell subset and 17.0-19.8% for the CD56 NK cell subset, we can estimate by calculation that approximately 86% of NK cells carried both CD16 and CD56 molecules on the same cells (Table 11). Generally, males have a higher number of NK cells than females (independent t-test; $p=0.008$). In addition, age variation as analysed by simple linear regression analysis, has not been found with NK cell number.

Table 11. NK cell subpopulations by percentages and absolute numbers

NK cells Immunophenotype	Sex	N	Mean	SD	Median	95% Reference range	Sex difference	Age Variation
Immunophenotypes by percentages								
CD3 ⁻ CD16 ⁺ and/or CD56 ⁺	M	67	23.2	9.1	21.0	21.0-25.5	4.0 %	NS (p=0.008)
	F	58	19.2	7.5	19.0	17.2-21.2	M > F	
	Both	125	21.4	8.6	20.0	19.8-22.9		
CD3 ⁻ CD16 ⁺	M	65	21.0	7.9	20.0	19.0-22.9	3.1 %	NS (p=0.028)
	F	56	17.9	7.3	18.0	15.9-19.8	M > F	
	Both	121	19.5	7.8	19.0	18.1-20.9		
CD3 ⁻ CD56 ⁺	M	65	19.6	8.1	17.0	17.6-21.6		NS NS
	F	56	17.0	7.3	17.5	15.0-18.9	NS	
	Both	121	18.4	7.8	17.0	17.0-19.8		
Immunophenotypes by absolute numbers								
CD3 ⁻ CD16 ⁺ and/or CD56 ⁺	M	67	533.3	352.1	455.0	447.4-619.1	152.3	NS (p=0.003)
	F	58	381.0	202.3	366.5	327.8-434.1	cells/μl	
	Both	125	462.6	301.0	403.0	409.3-515.9	M > F	
CD3 ⁻ CD16 ⁺	M	65	481.7	302.8	429.0	406.7-556.7	124.5	NS (p=0.009)
	F	56	357.2	192.9	349.5	305.5-408.9	cells/μl	
	Both	121	424.1	264.2	377.0	376.5-471.6	M > F	
CD3 ⁻ CD56 ⁺	M	65	447.7	289.7	351.0	375.9-519.5	106.9	NS (p=0.022)
	F	56	340.8	200.5	328.0	287.1-394.5	cells/μl	
	Both	121	398.2	257.0	340.0	352.0-444.5	M > F	

NS = no significant difference

4. Reference ranges of CD4⁺T cells in healthy Thai adults based on absolute number of CD4⁺T cell count

Absolute number of CD4⁺T cells of healthy subjects were classified into three ranges of ≤ 399 , 400-499, ≥ 500 cells/ μl . The results demonstrated that 26.4 % of our subjects had CD4⁺T cell count below 500 cells / μl ; and 13.6% had CD4⁺T cell count below 400 cells/ μl (Table 12). CD4⁺T cell count of these subjects ranged between 281 and 1,458 cells/ μl (data not shown), and with 95 % reference range of 624.3-703.3 cells/ μl (Table 10).

Table 12. Distribution of subjects with different ranges of CD4⁺ T cells

Sex	Number of subjects with CD4 ⁺ T cells of			Total
	≤ 399 cells/μl	400-499 cells/μl	≥ 500 cells/μl	
Male	9	9	49	67
Female	8	7	43	58
Both	17 (13.6%)	16 (12.8%)	92 (73.6%)	125 (100%)

5. Correlation between level of CD4⁺T cell count and changes in level of CD19⁺B, CD8⁺T or NK lymphocytes in HIV seropositive subjects

It has been well accepted that number of CD4⁺T cells decreases as the HIV disease progress. And as number of CD4⁺T cell decrease, number of some lymphocyte subset(s) should increase in order to reach the total count of 100 %. The present study assayed for number of CD8⁺T cells, CD19⁺B cells and NK cells in HIV infected cases with CD4⁺T cells < 14% or ≥ 14% in order to determine number of lymphocyte subset (s) which was affected when number of CD4⁺T cells changes during course of the disease.

Statistical analyses using Pearson correlation and linear regression demonstrated that change on CD4⁺T cell count had statistically significant effect on level of CD8⁺T cell and NK cell counts in term of inverse correlation. Meanwhile, the change on CD4⁺T cell count had no statistical effect on level of CD19⁺B cell count (Figs. 6-8 and, Table 13).




Figure 6. Correlation between level of CD4⁺T lymphocyte and CD19⁺B lymphocyte count in HIV seropositive subjects with ;
 a) CD4 < 14% (n=48) b) CD4 ≥ 14% (n=64) c) total subjects (n=112)




Figure 7. Correlation between level of CD4⁺T lymphocyte and CD8⁺T lymphocyte count in HIV seropositive subjects with ;
 a) CD4 <14% (n=48) b) CD4 ≥ 14% (n=64) c) total subjects(n=112)




Figure 8. Correlation between level of CD4⁺T lymphocyte and NK lymphocyte count in HIV seropositive subjects with ;
 a) CD4 <14% (n=48) b) CD4 ≥14% (n=64) c) total subjects (n=112)

Table 13. Correlation between level of CD4⁺ T cells and CD19⁺B, CD8⁺ T cells or NK lymphocytes in HIV seropositive subjects

Parameter	Correlation with level of CD4 ⁺ T cells		
	CD4<14% (n=48)	CD4 ≥ 14% (n=64)	All (n=112)
CD19⁺ B lymphocyte count	r = -0.095 p = 0.522	r = -0.139 p = 0.274	r = 0.058 p = 0.543
CD8⁺ T lymphocyte count	r = -0.178 p = 0.225	r = -0.375 p = 0.002	r = -0.467 p = 0.000
NK lymphocyte count	r = -0.071 p = 0.633	r = -0.075 p = 0.556	r = -0.220 p = 0.020

6. Synthesis of IFN γ , RANTES, MIP-I α and MIP-I β in different lymphocyte subsets of HIV seronegative and HIV seropositive subjects

Flow cytometry was used to analyse for expression of cell surface markers and intracellular cytokine and chemokine production in CD3⁺ and CD8⁺T cells and NK cells upon activation with PMA and ionomycin. The study included 10 HIV seronegative subjects and 28 HIV seropositive subjects (nine had CD4⁺T cells <14%; and 19 had CD4⁺T cells \geq 14 %). Freshly heparinized whole blood specimens were stimulated with PMA and ionomycin for 4 hours in the presence of BFA. Unstimulated control (without the stimulation of PMA and ionomycin) were included to verify the staining specificity, as a guide for setting markers to delineate positive and negative populations and to assess the level of background of cytokine synthesis.

The stimulated and unstimulated control blood samples were stained by panel of monoclonal antibodies that determined lymphocyte subset and intracellular cytokines : IFN γ , RANTES, MIP-I α , MIP-I β . The frequencies of cytokine expressing cells were enumerated by 3-color flow cytometric analysis. Gating of lymphocyte subsets and fluorescence intensity of the stained cells are shown in Figures 9-12.




Figure 9. Flow cytometric three-color analysis of IFN γ expressing CD3, CD8 and NK cells from A: unstimulated control; and B: PMA and ionomycin stimulation




Figure 10. Flow cytometric three-color analysis of RANTES expressing CD3, CD8 and NK cells from A): unstimulated control; and B): PMA and ionomycin stimulation





Figure 11. Flow cytometric three-color analysis of MIP-1 α expressing CD3, CD8 and NK cells from A): unstimulated control and; B: PMA and ionomycin stimulation

A. no stimulation



B. stimulation with PMA and ionomycin




Figure 12. Flow cytometric three-color analysis of MIP-1 β expressing in CD3, CD8 and NK cells from A): unstimulated control and; B): PMA and ionomycin stimulation

6.1 Spontaneous production of cytokine/ chemokine without antigenic stimulation

Staining of the unstimulated cultures showed that spontaneous synthesis of IFN γ , RANTES, MIP-I α , MIP-I β was clearly observed in CD8⁺T cells, but not so in CD3⁺T cells and NK cells (Figs. 13-15 and Table 14). Percentages of cells spontaneously synthesized cytokine/ chemokines was the background value which will be subtracted from the percentages of stained cells of the stimulated culture to yield the corrected percentages of cells synthesizing cytokine/ chemokines as the result of PMA and ionomycin activation.

Table 14. The background value of cytokine expressing cell without antigenic stimulation

	Study groups	% of Mean \pm SD		
		CD3 producing cells	CD8 producing cells	NK producing cells
IFN γ	HIV seronegative subjects (n=10)	0.6 \pm 1.2	15.5 \pm 20.4	0.1 \pm 0.1
	HIV seropositive subjects (n=28)	0.8 \pm 1.9	7.6 \pm 7.7	0.0 \pm 0.0
RANTES	HIV seronegative subjects (n=10)	1.7 \pm 2.3	6.8 \pm 6.0	1.0 \pm 2.2
	HIV seropositive subjects (n=28)	1.7 \pm 2.8	10.5 \pm 8.4	0.4 \pm 0.7
MIP-Iα	HIV seronegative subjects (n=10)	0.5 \pm 0.1	5.4 \pm 5.3	0.0 \pm 0.1
	HIV seropositive subjects (n=28)	1.0 \pm 2.4	8.3 \pm 8.1	0.0 \pm 0.0
MIP-Iβ	HIV seronegative subjects (n=10)	0.8 \pm 1.2	6.7 \pm 5.8	0.3 \pm 0.4
	HIV seropositive subjects (n=28)	1.5 \pm 3.4	8.7 \pm 7.6	0.3 \pm 0.4




Figure 13. Production of intracellular cytokines, e.g., IFN γ , RANTES, MIP-I α and MIP-I β in non-activated CD3⁺T cells from HIV seronegative (n=10) and seropositive subjects (n=28)




Figure 14. Production of intracellular cytokines, e.g., IFN γ , RANTES, MIP-I α and MIP-I β in non-activated CD8⁺ T cells from HIV seronegative (n=10) and seropositive subjects (n=28)

6.2. Production of cytokine/ chemokine upon stimulation with PMA and ionomycin

Among CD3⁺T, CD8⁺T and NK cells from three groups of subjects: HIV seronegative subjects, HIV seropositive cases with CD4⁺ T cells < 14% and HIV seropositive cases with CD4⁺T cells \geq 14%, it has been shown upon stimulation that CD8⁺T cells and NK cells were good sources of IFN γ and MIP-I β ; whereas MIP-I α and RANTES were mainly produced from CD8⁺T cells. Comparison was also conducted between the HIV seronegative subjects and HIV infected subjects among IFN γ , RANTES, MIP-I α and MIP-I β expressing in CD3⁺T, CD8⁺T and NK cells. The study demonstrated that IFN γ , RANTES, MIP-I α and MIP-I β expressing CD8⁺T and NK cells were not significantly different as compared between the HIV seronegative subjects and HIV infected subjects. (as analysed by independent t-test and Mann Whitney U-test).

The results of cytokine/ chemokine production in CD3⁺T cells, CD8⁺T cells and NK cells of HIV seronegative and HIV seropositive subjects are shown in Figure. 16 together with the summarization in Table 15.




Figure 16. Corrected percentages of IFN γ , RANTES, MIP-1 α and MIP-1 β expressing CD3⁺T, CD8⁺T and NK lymphocytes in HIV seronegative (n=10) and seropositive subjects (n=28)

Table 15. Flow cytometric analyses on cytokine and chemokine producing cells upon stimulation by PMA and ionomycin

	Study groups	CD3 producing cells	CD8 producing cells	NK producing cells
IFN γ	HIV seronegative subjects (n=10)	23.2 \pm 6.2 ^b	30.3 \pm 22.2 ^b	20.1 \pm 10.8 ^b
	HIV seropositive subjects with ; CD4<14% (n=9)	37.2 \pm 31.5 ^b	34.9 \pm 22.7 ^b	30.1 \pm 28.7
	CD4 \geq 14% (n=19)	29.3 \pm 23.4 ^b	31.2 \pm 26.9 ^b	23.1 \pm 18.7 ^b
RANTES	HIV seronegative subjects (n=10)	0.2 \pm 0.3 ^a	12.8 \pm 9.1 ^b	0.4 \pm 0.8 ^a
	HIV seropositive subjects with ; CD4<14% (n=9)	6.9 \pm 20.5 ^a	19.2 \pm 22.2 ^b	5.5 \pm 15.6 ^a
	CD4 \geq 14% (n=19)	1.6 \pm 2.3 [*]	8.2 \pm 12.9 ^b	0.8 \pm 1.2 ^a
MIP-Iα	HIV seronegative subjects (n=10)	0.6 \pm 1.5 ^b	12.8 \pm 8.1 ^a	0.3 \pm 0.8 ^b
	HIV seropositive subjects with ; CD4<14% (n=9)	6.7 \pm 19.7 ^b	19.9 \pm 22.2 ^a	6.3 \pm 18.2 ^b
	CD4 \geq 14% (n=19)	1.5 \pm 2.2 ^b	8.2 \pm 12.0 ^a	0.5 \pm 1.0 ^b
MIP-Iβ	HIV seronegative subjects (n=10)	35.0 \pm 15.5 ^b	28.5 \pm 22.1 ^a	35.1 \pm 15.0 ^b
	HIV seropositive subjects with ; CD4<14% (n=9)	35.7 \pm 33.0 ^b	19.1 \pm 20.7 ^b	32.3 \pm 30.5 ^b
	CD4 \geq 14% (n=19)	32.9 \pm 23.3 ^b	17.5 \pm 21.4 ^a	32.1 \pm 23.5 ^b

* = There was statistically significant difference as compared between HIV seronegative and seropositive subjects (p<0.05)

a = data was analysed by Mann Whitney U-test; b = data was analysed by independent t-test

8. NK cell cytotoxicity assay

NK cell cytotoxicity assay was performed in 69 HIV seronegative (38 males, 31 females with age-range of 21-52, 21-55 years; and mean age of 34.05 ± 8.8 , 31.3 ± 11.1 , respectively), and 107 HIV seropositive subjects (46 males, 61 females with age-range of 21-53, 18-50 years and mean age of 32.9 ± 8.13 , 26.97 ± 5.63 , respectively). Fresh, unsensitized PBMCs from the subjects was used as source of NK effector cells (E) and K562 cells labeled with ^{51}Cr were used as the target cells (T). The NK cell cytotoxic activity was measured by chromium release from the radiolabeled target cells. Five different E: T ratios were used in a test assay and the value of NK cytotoxic activity was expressed in term of LU_{20} .

The result on NK cell cytotoxic activity in HIV seronegative subjects showed no sex difference (independent t-test; $p \geq 0.05$) (Table 16) with an average $\text{LU}_{20}/10^6\text{PBMC}$ of 10.5 ± 8.4 . Among HIV seropositive subjects, the study did not find significant difference in NK cell number and NK cell cytotoxic activity as compared between the group with $\text{CD4}^+\text{T}$ cell count $< 14\%$ ($\text{LU}_{20}/10^6\text{PBMC} = 6.4 \pm 5.8$) and the group with $\text{CD4} > 14\%$ ($\text{LU}_{20}/10^6\text{PBMC} = 8.1 \pm 7.1$) (independent t-test; $p \geq 0.05$). However, the study has demonstrated a decrease in NK cell cytotoxicity in both groups of HIV infected cases as compared either one with HIV seronegative controls (Mann Whitney U test; $p < 0.05$). Moreover, a decrease in NK cell number was found in HIV infected cases with $\text{CD4} \geq 14\%$ as compared to HIV seronegative controls (independent t-test; $p = 0.000$) (Table 17).

Similarly, when the mean values of $\text{LU}_{20}/10^6$ NK cells from all subjects was taken into account, a decrease in NK cytotoxic activity could be observed in the late disease stage of HIV infected subjects with $\text{CD4} < 14\%$ as compared to HIV seronegative subjects (Table 17). This study also showed the correlation between number of NK cells or their subpopulation (CD16^+ and CD56^+ NK cells) and NK cytotoxic activity (Figures 17-19). The result suggested that both NK cell subpopulations had cytotoxic function. Moreover, NK cells in late disease stage could retain their function, even though, not as good as NK cells from HIV seronegative subjects. On the other hand, our study had shown an inverse correlation between NK

cell cytotoxic activity and CD4⁺T cell count only in HIV seronegative subjects, whereas, in HIV seropositive subjects, this correlation could not be demonstrated.

Table 16. Determination for the amount of NK cells and their function in HIV seronegative, healthy Thais as classified by sex

Subjects	N	Mean percentages of NK cells \pm SD	Cytotoxic activity \pm SD (LU20/10⁶ PBMC)
Total subjects	69	20.8 \pm 7.8	10.5 \pm 8.4
Males	38	21.8 \pm 8.8	10.7 \pm 9.2 ^a
Females	31	19.5 \pm 6.4	10.1 \pm 7.5 ^a

a = There was no statistically significant difference in NK cell activity as compared between sex (independent t-test; $p \geq 0.05$).

Table 17. Correlation between NK cell number and NK cytotoxic activity in HIV seronegative and HIV seropositive subjects with different ranges of CD4⁺T lymphocyte count

Subjects	N	percentages of NK cells \pm SD	Cytotoxic activity \pm SD (LU20/ 10 ⁶ PBMCs)	Mean LU20/10 ⁶ NK cells
HIV seronegative subjects	69	20.8 \pm 7.8	10.5 \pm 8.4	52.3 \pm 42.9
HIV-1 seropositive subjects with ;				
CD4 \geq 14 %	61	16.0 \pm 7.3 ^{a,b} (p=0.000)	8.1 \pm 7.1 ^{a,b*} (p=0.049)	52.1 \pm 46.8 ^{c*,d} (p=0.006)
CD4 < 14 %	46	19.0 \pm 8.5 ^{c, a}	6.4 \pm 5.8 ^{a, b*} (p=0.002)	32.2 \pm 26.0 ^{b*,d} (p=0.005, 0.006)

a, a* = There was no statistically significant difference as compared between HIV seropositive subjects with CD4<14% and \geq 14% (p \geq 0.05).

b, b* = There was statistically significant difference as compared between HIV seronegative and HIV seropositive subjects (p<0.05).

c, c* = There was no statistically significant difference as compared between HIV seronegative and HIV seropositive subjects (p \geq 0.05).

d = There was statistically significant difference as compared between HIV seropositive subjects with CD4<14% and \geq 14% (p<0.05).

a, b, c, d = analysed by independent t-test; a*, c* = analysed by Mann Whitney U-test




Figure 17. Correlation between LU20/ 10⁶ PBMC and NK cell numbers in; a) HIV seronegative subjects; b) HIV seropositive subjects with CD4_≥14% and; c) HIV seropositive subjects with CD4<14%




Figure 18. Correlation between LU20 / 10⁶ PBMC and %CD16⁺NK cell subpopulation in; a) HIV seronegative subjects ; b) HIV seropositive subjects with CD4_≥14% and; c) HIV seropositive subjects with CD4<14%




Figure 19. Correlation between LU20 / 10⁶ PBMC and %CD56⁺NK cell subpopulation in; a) HIV seronegative subjects; b) HIV seropositive subjects with CD4_≥14% and; c) HIV seropositive subjects with CD4_<14%




Figure 20. Correlation between LU20/10⁶ PBMC and %CD4⁺T cell count in; a) HIV seronegative subjects; b) HIV seropositive subjects with CD4<14% and ; c) HIV seropositive subjects with CD4≥14%

CHAPTER VI

DISCUSSION

Immune mechanisms are mediated by multiple white blood cell types: granulocytes, monocytes and lymphocytes. A functional classification of lymphocytes can further distinguish B, T and NK cells, utilizing the expression of specific cell surface glycoproteins and/ or the specific mechanism by which the cell protects the host from foreign antigens. The number and functional capacity of lymphocyte subsets reflect the overall state of immune competence of an individual. Enumeration of lymphocyte subsets by flow cytometric method has made possible the precise identification of lymphocyte immunophenotypes. In a variety of clinical situation, number of lymphocyte subset have become important in the disease diagnosis and monitoring the immunorestitution treatments. For example, the relative proportion of lymphocyte subsets may be altered in certain disease states, such as depletion of CD4⁺T lymphocytes during the course of HIV infection (2).

Phenotyping for reference ranges of lymphocyte subsets can be influenced by time of blood collection, duration and temperature of specimen storage before staining, flow cytometric analysis, and also by methodologies (62-65). Previous multi-center study reported the effects of time lapse from phlebotomy to staining and fixing, and also time lapse from staining and fixing to analysis by flow cytometry (64). In addition, the Japanese study suggested circadian variation in circulating T cell subsets as evaluated in whole blood by flow cytometry technique. Rhythmicity with high values of lymphocyte numbers was shown during night time and low values during daytime (62).

In order to control the variations from the above technical practices, we performed phlebotomy during 9.00-11.00 a.m., followed by staining and fixing within 3 hours; then, flow cytometrically analyzed within 24 hours after staining and fixing. We also kept blood specimens in air-conditioned room (25⁰ C) until stained.

Previous investigators demonstrated that changes in absolute number of lymphocyte subset were not always consistent with changes in percentages values (55). It should be kept in mind that absolute number was derived from three variable values: white blood cell count, total lymphocyte count and percentages of lymphocyte subset. Therefore, the value presented by percentages may not correlated with that of the absolute number (Tables 9 and 10). In addition, with respect to the difference in the values of WBC count by sex (Table 8), we suggested that statistical analyses of the datasets presented in term of percentage values should be more accurate, because the result obtained is independent from other variable confounding factors.

The result of our study was compared to previous works done in Thai blood donors as reported by 1) Webster *et al.* (10) in the multi-site study in Bangkok, i.e., Siriraj Hospital, Chulalongkorn Hospital and Armed Forces Research Institute of Medical Sciences (AFRIMS) in 1996, and 2) Vithayasai *et al.* (58) in the Chiang Mai study. The different studies presented their data in different ways and parameters. Collectively, some points of similarity and some points of discrepancy were found. In overall, all two studies demonstrated that Thai people had relative low CD4⁺T cell count. The CD4⁺T cell reference values of the present study and of that from Chulalongkorn Hospital were comparable *viz*, mean = 32.1 *v.s.* 32.8 % or 663.8 *v.s.* 730 cells/ μ l for both sexes, but both values were relatively lower than those pooled data generated by Siriraj and AFRIMS Laboratories *viz*, 37.4% or 840 cells/ μ l, and lower than those reported by Chiang Mai Laboratory *viz*, 36.1% or 910 cells/ μ l. The mean CD4/CD8 ratio in our study was about the same as that demonstrated by Webster *et al.* (10) *viz* 1.3 *v.s.* 1.35, while it was 1.5 by Vithayasai *et al.* (58). Both Webster *et al.* (10) and the present study revealed high NK cell counts especially in male subjects.

Gathering from various reports, the most important finding was variation of lymphocyte reference ranges according to ethnic groups (Table 1). Mongoloids had lower CD4⁺ T cell count, but higher number of NK cells (1, 10, 56). The values of CD4⁺T cells in Mongoloid people from different countries were similar, i.e., 32% in Thai people, 32% in Chinese living in Hong Kong (56), 34.8% in Chinese living in Malaysia and 32.6% in native Malay people (1). It was noteworthy that Indian people in Malaysia had higher CD4⁺T cells count; Indians are classified as Caucasian ethnic group. Mongoloid ethnic with relatively high CD4⁺T cell count was observed in Japanese (72). Nevertheless, the count was still lower than those reported in Caucasians. Reports for Caucasians from Western countries demonstrated higher CD4⁺ T cell counts varying from 42 to 48% (53-55, 63). Our study found that 26.4% of Thai people had CD4⁺T cell count <500 cells/ μ l, and 13.6% had CD4⁺T cell count < 400 cells/ μ l. The subject who had lowest amount of CD4⁺T cell count of 281 cells/ μ l possessed normal values of WBC and CD4⁺T cell count; however, her lymphocyte count was only 15%. This subject was apparently healthy during the week of blood collection, and had no obvious medical history. Similarly, 2.9% of Chinese living in Hongkong had CD4⁺T cell count <300 cells/ μ l; and 15.9% had CD4⁺T cell count < 500 cells/ μ l (56).

Refer to guidelines from US. and CDC, CD4⁺T cell levels are used as criteria for AIDS case definitions, for initiation of anti-retroviral drugs and for pre-exposure prophylaxis of opportunistic infections (69-71). Treatment of Mongoloid people infected with HIV by basing on level of CD4⁺T cell count recommended for Caucasians should be taken into careful reconsideration.

The compensation for Mongoloids for their low CD4⁺T count is their high NK cell level. We also studied the changes in lymphocyte subset in HIV infected cases by determining the change in number of CD8⁺T cells, CD19⁺B cell and NK cells in correlation with the level of CD4⁺T cell count (Table 13 and Figures 6-8). This study revealed that level of CD4⁺T cell count had inversed correlation with the CD8⁺T and NK lymphocyte count. Nevertheless, we have not seen any significant correlation between CD4⁺T and CD19⁺B lymphocyte counts. This study indicated that during the progression

of HIV infection, a quantitative defect in CD4⁺T cell count had resulted in an increase in expansion of CD8⁺T and NK lymphocyte level.

Activation of NK cells by immunological modulators or drugs may be a supplementary treatment for cancer and viral, especially HIV infections. It has been suggested that NK cells may be a factor protecting the health of HIV seropositive individuals with very low CD4 cell counts(83). A low level of NK cells in HIV infected cases was associated with rapid disease progression, compared to those who progressed more slowly (17). The Los Angeles Center of the Multicenter AIDS Cohort Study demonstrated that percentages of NK cells severely decreased in the presence of HIV infection (81). NK cells are relatively resistant to HIV infection. Whether, the HIV can replicate in NK cells or not is still inconclusive (185-187). NK cells can mediate killing activity which is not antigen specific. Therefore, NK cells can potentially overcome the problem of HIV antigenic diversity, while this is an important factor limiting the roles of specific cytotoxic T lymphocytes and neutralizing antibody in HIV infection. Previous work done by de Souza *et al.* (84) reported that at comparable number, NK cells of Thais had higher cytotoxicity function than North Americans. The role of NK cells in HIV infected Thais and other Mongoloid people deserves further investigation.

The present study looked into those NK cells which expressed CD16 and/or CD56 molecules. CD16 molecules function as Fc receptors that bind IgG (Fc γ receptor III), thus, these receptors can link NK cells to the target cells coated with IgG and lead to target cell destruction *via* the mechanism so called antibody-dependent cell-mediated cytotoxicity (ADCC) (146, 188). CD56⁺ NK cells play an important role in the production of several kinds of cytokines following an induction by monokines (146, 188). It was previously reported that activated NK cells may release RANTES, MIP-1 α and MIP-1 β , the cytokines capable of binding HIV coreceptor, and thus, blocking HIV replication (19). NK cells also release tumor necrosis factor and IFN γ which enhance immunological response (189). Our study demonstrated that 86% of NK cells express both CD16⁺ and CD56⁺ molecules on the same cell; and both subpopulations have cytotoxic function. NK cells also recognize and directly kill target cells through killer activating receptors. These killing activity is usually overridden by killer-inhibitory

receptors on recognition of MHC class I molecule under normal situation. Unfortunately, immunophenotypes of these molecules is not known (188).

NK cell is a critical component of innate immune response and thought to be important during the early stage of HIV infection (148, 190, 191). Although cytokine expressing CD8⁺T cells have been extensively documented, there have been few studies on cytokine production from NK cells.

Previous group of investigators had found that NK cells are as potent as CD8⁺T cells in the mediation of suppression of HIV replication, predominantly by secretion of CC-chemokines (21). In addition, Fehniger *et al.* (191) demonstrated that supernatant from NK cells can inhibit the replication not only of M-tropic HIV-1, but also of T-tropic HIV-1 strain. Moreover, Scharon *et al.* (192) demonstrated that NK cell could produce IFN- γ during the first week of infection, and occurred more rapidly than might be expected for conventional primary T cell response. Oliva *et al.* (19) found the level of RANTES, MIP-1 α and MIP-1 β synthesized by NK cells was comparable to those produced by the stimulated CD8⁺T cells and the study done by Vitale *et al.* (20) reported a higher percentages of IFN- γ producing NK cells as compared to number of IFN- γ producing T cells from both the HIV seronegative and HIV seropositive subjects.

Concerning the methods used for determining cytokine and chemokine production such as ELISA, RT-PCR, T cell cloning and ELISpot, all of these methods are time consuming and labor intensive. Moreover, these techniques require long activation time and cannot determine the frequencies and the phenotypes of the cytokine-producing cells at the single cell level (93, 117, 193-196). The flow cytometric method has been designed to simply and rapidly assess intracellular cytokine production in different leukocyte subsets resided in PBMCs or whole blood samples. These procedures determine the percentages of activated cells and the identification of leukocyte subsets capable of expressing various cytokines and cell surface antigens simultaneously. The ability to assess the key intracellular functional markers by multiparametric flow cytometry offer some unique advantages in a number of clinical applications.

The present study was a further effort to reveal the production of various types of intracellular cytokines in NK and CD8⁺T cells. It was shown that NK and CD8⁺T cells were the expandable sources of cytokine and chemokine production in healthy subjects and HIV-1 infected ones. Few percentages of cytokine or chemokine producing CD8⁺T, and NK cells were observed in unstimulated PBMC samples (Figures 13-15 and Table 14). Our study showed that NK cells and CD8⁺T cells from HIV infected cases were functionally active with regards to the production of cytokine and chemokine in response to PMA and ionomycin as compared to HIV seronegative subjects. Both NK and CD8⁺T cells from HIV seronegative and HIV seropositive subjects were capable of producing high level of IFN- γ and MIP-1 β ; whereas, the production of RANTES and MIP-1 α was mainly produced from CD8⁺T cells (Table 15). Thus, HIV infection of any disease stage may not affect the ability of production of cytokine and chemokine. In contrast, Azzoni *et al.* (119) found that NK cells from HIV-infected cases were impaired in their ability to produce IFN- γ as compared to the HIV seronegative donors. Moreover, Peruzzi *et al.* (197) suggested that the gp120 mediated decrease in IFN- γ production by NK cells may contribute to the cytokine imbalance observed in HIV infection.

NK cell number and function are altered in many diseases, including HIV, certain cancers, viral infection, and autoimmune disease (16, 17). Rapid disease progression after HIV infection also has been correlated with a decrease in NK cell cytotoxic activity (180). Decrease in amount and cytotoxic function of NK cells by HIV infection was also found in several studies (81, 83, 84, 185).

This study also measured NK cell cytotoxic activity by ⁵¹Cr release assay. The significant correlation between the cytotoxic activity and level of NK cells or their subpopulation could be observed. Correlation between the NK cytotoxic activity and level of CD4⁺T cell count was also determined; and an inverse correlation was found only in HIV seronegative subjects. In contrast, there was no any correlation between CD4⁺T cell count and NK cell cytotoxic activity as observed in HIV infected cases. Our subjects included some asymptomatic HIV infected cases with low CD4⁺T cell count who had high NK cell cytotoxic activity (data not shown) which was similarly reported by Ironson *et al.* (83). NK cytotoxic activity may, therefore, be a factor protecting the health of these

people. The host may maintain a degree of protection from HIV-related complications by compensating for the loss of specific T-cell immunity with non-specific immune defense.

Regarding the sex influence on NK cell cytotoxic activity in HIV seronegative subjects, a statistically significant difference was not observed. As compared to HIV seronegative subjects, a decline in NK cell activity was observed in all HIV seropositive subjects with CD4⁺T count <14 and \geq 14 %. There was no significant difference in percentage of NK cells as compared the HIV-1 seronegative subjects with the HIV seropositive subjects who had CD4⁺T count less than 14 % (Table 17). This might be due to the compensation of number of NK cells for the decrease in number of CD4⁺T cells.

Our study was different from what had been reported by de Souza (84) who demonstrated that NK cell number and function of the Thai HIV seronegative and seropositive subjects were not significantly different. In contrast, as compared between the HIV seronegative and seropositive North American subjects, they demonstrated the profound impairment of NK cell number and function (80, 84). They also demonstrated that the number and percentage of NK cells in HIV seronegative Thais and North Americans were similar, but the cytotoxic activity was significantly greater in the Thais. Meanwhile, Douglas *et al.* (80) reported that there was no significant difference in NK cytotoxic activity by race.

Various mechanisms have been proposed to explain why NK cell function is impaired in HIV infection. Bonavida *et al.* (179) demonstrated that defective NK cell activity can be the result from; a) a decreased number of effector cells; b) inability of the NK cells to recognize and interact with NK-sensitive targets; c) failure of NK target cells to trigger and activate the NK cells to release NK cytotoxic factor; and d) failure of the NK cell to synthesize NK cytotoxic factor.

Peruzzi *et al.* (197) demonstrated a direct effect of envelope gp120 which can inhibit NK cytotoxic activity. The addition of gp120 to the cell culture was associated with a significant inhibitory effect on NK lysis and was clearly dose dependent (198). Moreover, Cauda *et al.* (199) showed that synthetic peptides corresponding to amino acid sequences 735-752 and 846-860 of the HIV transmembrane gp41, conjugated to protein

carriers, significantly inhibited the NK cell activity of normal PBMCs against the K562 cell line.

Since IFN- γ can be synthesized by CD4⁺T cells and play an important role in stimulating in innate cell-mediated immunity through NK cell cytotoxic activity (104). The decline of CD4⁺T cells in the progression of HIV infection may affect the level of IFN- γ production and NK cell cytotoxicity activity. IFN- γ and IL-2, the T_H1 cytokines which potently stimulate the NK activity, decrease markedly as the T_H1-T_H2 shift progress during the course of HIV infection (98). In addition, the products of T_H2 cytokine e.g., IL-10 and IL-12 have been found to dramatically inhibit NK cell activity (200). Nevertheless, our study demonstrated that the expression of IFN- γ by NK and CD8⁺T cells in HIV infected cases had no defect.

Some previous studies reported the mechanism responsible for the impairment of NK cell function during HIV infection such that NK cell served as a potential reservoir for HIV-1 or can be infected by HIV-1. Nevertheless, results obtained from different groups of investigators were controversial (185-187, 201, 202).

A second set of NK cells sharing characteristics with NK and T cell has been defined as NKT cells (157-160). NKT cells could be infected by HIV as they express both CD4 and CCR5 molecules. Therefore, resulted in NKT cell death. This event may be the other reason why NK cell cytotoxicity was impaired (165). Unutmas *et al.* (166) found that CD4⁺/CCR5⁺NKT cells were more susceptible to R5-tropic, but less susceptible to X-4 tropic virus *in vitro*. Recent findings have shown that number of NKT cells in PBMCs of HIV-1 infected individual are dramatically reduced as compared with healthy donors; and a positive correlation between high viral load and lower CD4⁺NKT numbers was established (164, 166).

Our study aimed to investigate the role of NK cells in the protection against HIV disease progression based on the advantages that a) Thai people had higher frequency of NK cells as compared to the Caucasians; b) NK cells could retain their activity on chemokine/cytokine production, even though, their cytotoxic activity may decrease; c) NK cell function is not antigen specific; therefore, they can overcome the problem of

HIV escape from humoral and cell-mediated immune response. Therefore, if NK cell activity can be sustained by any immunostimulator, it should provide an alternate approach for HIV treatment.

CHAPTER VII

CONCLUSION

The present study comprised three major parts : 1) flow cytometric analyses to establish the lymphocyte immunophenotype reference ranges for Thai people together with the additional investigation in HIV infected cases on lymphocyte immunophenotype changes according to level of CD4⁺ T cell count; 2) intracellular cytokine/chemokine production in NK cells and CD8⁺T cells of HIV seronegative and HIV seropositive subjects; 3) NK cell cytotoxic activity assayed in HIV seronegative and HIV seropositive subjects

Lymphocyte subpopulation , i.e., T, B and NK cells including NK cell subsets were enumerated by two color-flow cytometry in a total of 125 HIV seronegative Thai adults. The study demonstrated relatively low CD4⁺T cell count in these subjects, i.e., 26.3% of them had a CD4⁺T cell count of less than 500 cells/ μ l. In contrast, their NK cell counts were relatively high. Collective data from several studies including the present one suggested that high NK cell counts may be a compensation for CD4⁺T cell count in Mongoloid people. It was roughly estimated that about 86% of NK cells harbored CD16 and CD56 molecules. Statistical analyses of the percentage values showed that females had significantly higher CD3 (total CD3⁺T cell), but lower NK cell counts as compared to males ($p < 0.05$). Regarding age variation, an increase of 1.1% of CD4⁺T cell count per decade was seen. The change of CD4⁺T cell count had an effect on level of CD8⁺T cell and NK cell in term of inverse correlation, i.e., HIV infected cases with low level of CD4⁺T cell count had higher number of CD8⁺T cells and NK cells. In contrast, change in level of CD4⁺T cell count had no effect on B cell level.

Three color flow cytometric analysis was conducted to determine intracellular cytokine/chemokine production in CD3⁺T, CD8⁺T and NK cells. The study included a total of 10 HIV seronegative subjects and 29 HIV seropositive subjects. The HIV seropositive subjects were further divided into group of CD4<14% (n=9) and group of CD4≥14% (n=19). The study demonstrated that spontaneous synthesis of IFN γ , RANTES, MIP-I α and MIP-I β was clearly observed in CD8⁺T cells at low frequency, but not in NK cells. Moreover, it has been shown upon stimulation with PMA and ionomycin that CD8⁺T cells and NK cells were important sources of IFN γ and MIP-I β ; whereas RANTES and MIP-I α were mainly produced from CD8⁺T cells. The finding hold true in both in HIV seronegative and HIV seropositive subjects. This study suggested that HIV-1 infection has less effect on the ability of NK and CD8⁺T cell in the cytokine/chemokine production.

NK cytotoxic activity in 69 HIV seronegative and 107 HIV seropositive subjects were measured by a standard ⁵¹Cr release assay. It was shown that NK cell cytotoxic activity had no statistically significant difference by sex as observed in HIV seronegative subjects. Furthermore, NK cytotoxic activity of HIV seropositive subjects were significantly lower than that of HIV seronegative subjects.

NK cell possess several advantages in the control of HIV disease progression such as: 1) Thai people had high NK cell number; 2) NK cell recognition is antigen non specific so that it can overcome the problem of as escape of HIV mutant from cytotoxic T lymphocytes and neutralizing antibodies which both of them are antigen specific in function. This study, thus, propose NK cell as a novel approach on HIV disease treatment.

REFERENCES

1. Choong ML, Ton SH, Cheong SK. Influence of race, age and sex on the lymphocyte subsets in peripheral blood of healthy Malaysian adults. *Ann Clin Biochem* 1995;32 (Pt 6):532-9.
2. Kidd PG, Vogt RF, Jr. Report of the workshop on the evaluation of T-cell subsets during HIV infection and AIDS. *Clin Immunol Immunopathol* 1989;52(1):3-9.
3. Hunninghake GW, Crystal RG. Pulmonary sarcoidosis: a disorder mediated by excess helper T-lymphocyte activity at sites of disease activity. *N Engl J Med* 1981;305(8):429-34.
4. Alpert G, Leibovitz L, Danon YL. Analysis of T-lymphocyte subsets in measles. *J Infect Dis* 1984;149(6):1018.
5. Goetzman EA. Flow cytometry: basic concepts and clinical applications in immunodiagnosics. *Clin Lab Sci* 1993;6(3):177-82.
6. Carbone J, Gil J, Benito JM, Navarro J, Munoz-Fernandez A, Bartolome J, et al. Increased levels of activated subsets of CD4 T cells add to the prognostic value of low CD4 T cell counts in a cohort of HIV-infected drug users. *Aids* 2000;14(18):2823-9.
7. Graziosi C, Soudeyns H, Rizzardi GP, Bart PA, Chapuis A, Pantaleo G. Immunopathogenesis of HIV infection. *AIDS Res Hum Retroviruses* 1998;14 Suppl 2:S135-42.
8. Phillips RE, Rowland-Jones S, Nixon DF, Gotch FM, Edwards JP, Ogunlesi AO, et al. Human immunodeficiency virus genetic variation that can escape cytotoxic T cell recognition. *Nature* 1991;354(6353):453-9.
9. Collins KL, Chen BK, Kalams SA, Walker BD, Baltimore D. HIV-1 Nef protein protects infected primary cells against killing by cytotoxic T lymphocytes. *Nature* 1998;391(6665):397-401.

10. Webster HK, Pattanapanyasat K, Phanupak P, Wasi C, Chuenchitra C, Ybarra L, et al. Lymphocyte immunophenotype reference ranges in healthy Thai adults: implications for management of HIV/AIDS in Thailand. *Southeast Asian J Trop Med Public Health* 1996;27(3):418-29.
11. Loubeau M, Ahmad A, Toma E, Menezes J. Enhancement of natural killer and antibody-dependent cytolytic activities of the peripheral blood mononuclear cells of HIV-infected patients by recombinant IL-15. *J Acquir Immune Defic Syndr Hum Retrovirol* 1997;16(3):137-45.
12. Katz JD, Mitsuyasu R, Gottlieb MS, Lebow LT, Bonavida B. Mechanism of defective NK cell activity in patients with acquired immunodeficiency syndrome (AIDS) and AIDS-related complex. II. Normal antibody-dependent cellular cytotoxicity (ADCC) mediated by effector cells defective in natural killer (NK) cytotoxicity. *J Immunol* 1987;139(1):55-60.
13. Ahmad A, Menezes J. Antibody-dependent cellular cytotoxicity in HIV infections. *Faseb J* 1996;10(2):258-66.
14. Ahmad A, Menezes J. Positive correlation between the natural killer and gp 120/41-specific antibody-dependent cellular cytotoxic effector functions in HIV-infected individuals. *J Acquir Immune Defic Syndr Hum Retrovirol* 1995;10(2):115-9.
15. Whiteside TL, Herberman RB. Role of human natural killer cells in health and disease. *Clin Diagn Lab Immunol* 1994;1(2):125-33.
16. Brenner BG, Dascal A, Margolese RG, Wainberg MA. Natural killer cell function in patients with acquired immunodeficiency syndrome and related diseases. *J Leukoc Biol* 1989;46(1):75-83.
17. Bruunsgaard H, Pedersen C, Skinhoj P, Pedersen BK. Clinical progression of HIV infection: role of NK cells. *Scand J Immunol* 1997;46(1):91-5.
18. Rogers AS, Schwartz DF, Weissman G, English A. A case study in adolescent participation in clinical research: eleven clinical sites, one common protocol, and eleven IRBs. *Irb* 1999;21(1):6-10.

19. Oliva A, Kinter AL, Vaccarezza M, Rubbert A, Catanzaro A, Moir S, et al. Natural killer cells from human immunodeficiency virus (HIV)-infected individuals are an important source of CC-chemokines and suppress HIV-1 entry and replication in vitro. *J Clin Invest* 1998;102(1):223-31.
20. Vitale M, Caruso A, Licenziati S, Rodella L, Fiorentini S, Zauli G, et al. Differential production of IFN-gamma, analyzed at the single-cell level, by specific subsets of human NK and T cells from healthy and HIV(+) subjects. *Cytometry* 2000;39(3):189-94.
21. Kottlilil S, Chun TW, Moir S, Liu S, McLaughlin M, Hallahan CW, et al. Innate immunity in human immunodeficiency virus infection: effect of viremia on natural killer cell function. *J Infect Dis* 2003;187(7):1038-45.
22. Earl PL, Doms RW, Moss B. Oligomeric structure of the human immunodeficiency virus type 1 envelope glycoprotein. *Proc Natl Acad Sci U S A* 1990;87(2):648-52.
23. McCune JM, Rabin LB, Feinberg MB, Lieberman M, Kosek JC, Reyes GR, et al. Endoproteolytic cleavage of gp160 is required for the activation of human immunodeficiency virus. *Cell* 1988;53(1):55-67.
24. Chanh TC, Dreesman GR, Kanda P, Linette GP, Sparrow JT, Ho DD, et al. Induction of anti-HIV neutralizing antibodies by synthetic peptides. *Embo J* 1986;5(11):3065-71.
25. Levy J. HIV and the pathogenesis of AIDS. 2 ed: Washington, D.C.:American Society for Microbiology; 1988.
26. Abbas AK, Lichtman AH, Pober JS. Cellular and molecular immunology. 4 ed. Philadelphia: W.B. Saunders Company; 2000.
27. Janeway CA, Travers P, Wipport M, Capra JD. Immunobiology of the immune system in health and disease. 4 ed. New York: Current Biology Publication/ Garland Publishing; 1999.
28. Brook G. Medical Microbiology. Twenty-second ed. New York: Lange Medical Books/ McGraw; 2001.

29. Cheingsong-Popov R, Osmanov S, Pau CP, Schochetman G, Barin F, Holmes H, et al. Serotyping of HIV type 1 infections: definition, relationship to viral genetic subtypes, and assay evaluation. UNAIDS Network for HIV-1 Isolation and Characterization. *AIDS Res Hum Retroviruses* 1998;14(4):311-8.
30. Subbarao S, Limpakarnjanarat K, Mastro TD, Bhumisawasdi J, Warachit P, Jayavasud C, et al. HIV type 1 in Thailand, 1994-1995: persistence of two subtypes with low genetic diversity. *AIDS Res Hum Retroviruses* 1998;14(4):319-27.
31. Gao F, Robertson DL, Morrison SG, Hui H, Craig S, Decker J, et al. The heterosexual human immunodeficiency virus type 1 epidemic in Thailand is caused by an intersubtype (A/E) recombinant of African origin. *J Virol* 1996;70(10):7013-29.
32. Carr JK, Salminen MO, Koch C, Gotte D, Artenstein AW, Hegerich PA, et al. Full-length sequence and mosaic structure of a human immunodeficiency virus type 1 isolate from Thailand. *J Virol* 1996;70(9):5935-43.
33. Rosenberg ZF, Fauci AS. The immunopathogenesis of HIV infection. *Adv Immunol* 1989;47:377-431.
34. Rosenberg ZF. Immunopathogenesis of HIV infection. In *AIDS: Etiology, diagnosis, treatment and prevention*. 3rd ed. Philadelphia: J.B. Lippincott; 1992.
35. Blacklaws BA. Quantification of the reservoir of HIV-1. *Trends Microbiol* 1997;5(6):215-6.
36. Pantaleo G, Graziosi C, Butini L, Pizzo PA, Schnittman SM, Kotler DP, et al. Lymphoid organs function as major reservoirs for human immunodeficiency virus. *Proc Natl Acad Sci U S A* 1991;88(21):9838-42.
37. Heath SL, Tew JG, Szakal AK, Burton GF. Follicular dendritic cells and human immunodeficiency virus infectivity. *Nature* 1995;377(6551):740-4.

38. An P, Nelson GW, Wang L, Donfield S, Goedert JJ, Phair J, et al. Modulating influence on HIV/AIDS by interacting RANTES gene variants. *Proc Natl Acad Sci U S A* 2002;99(15):10002-7.
39. Walker L, Wilks D, O'Brien J, Habeshaw J, Dalgleish A. Localized conformational changes in the N-terminal domain of CD4 identified in competitive binding assay of monoclonal antibodies and HIV-1 envelope glycoprotein. *AIDS Res Hum Retroviruses* 1992;8(6):1083-90.
40. Stantchev TS, Broder CC. Human immunodeficiency virus type-1 and chemokines: beyond competition for common cellular receptors. *Cytokine Growth Factor Rev* 2001;12(2-3):219-43.
41. Hogan CM, Hammer SM. Host determinants in HIV infection and disease. Part 1: cellular and humoral immune responses. *Ann Intern Med* 2001;134(9 Pt 1):761-76.
42. Turner BG, Summers MF. Structural biology of HIV. *J Mol Biol* 1999;285(1):1-32.
43. Pantaleo G, Graziosi C, Fauci AS. New concepts in the immunopathogenesis of human immunodeficiency virus infection. *N Engl J Med* 1993;328(5):327-35.
44. Collins KL, Baltimore D. HIV's evasion of the cellular immune response. *Immunol Rev* 1999;168:65-74.
45. Daar ES, Moudgil T, Meyer RD, Ho DD. Transient high levels of viremia in patients with primary human immunodeficiency virus type 1 infection. *N Engl J Med* 1991;324(14):961-4.
46. Graziosi C, Pantaleo G, Kotler DP, et al. Dissociation between HIV expression in peripheral blood versus lymphoid organs of the same patients. *Clin Res* 1992;40:333A.
47. Fauci AS. The human immunodeficiency virus: infectivity and mechanisms of pathogenesis. *Science* 1988;239(4840):617-22.
48. Fauci AS, Schnittman SM, Poli G, Koenig S, Pantaleo G. NIH conference. Immunopathogenic mechanisms in human immunodeficiency virus (HIV) infection. *Ann Intern Med* 1991;114(8):678-93.

49. Phair JP. Keynote address: variations in the natural history of HIV infection. *AIDS Res Hum Retroviruses* 1994;10(8):883-5.
50. Cavacini LA, Emes CL, Power J, Underdahl J, Goldstein R, Mayer K, et al. Loss of serum antibodies to a conformational epitope of HIV-1/gp120 identified by a human monoclonal antibody is associated with disease progression. *J Acquir Immune Defic Syndr* 1993;6(10):1093-102.
51. Lifson AR, Buchbinder SP, Sheppard HW, Mawle AC, Wilber JC, Stanley M, et al. Long-term human immunodeficiency virus infection in asymptomatic homosexual and bisexual men with normal CD4+ lymphocyte counts: immunologic and virologic characteristics. *J Infect Dis* 1991;163(5):959-65.
52. De Paoli P, Battistin S, Santini GF. Age-related changes in human lymphocyte subsets: progressive reduction of the CD4 CD45R (suppressor inducer) population. *Clin Immunol Immunopathol* 1988;48(3):290-6.
53. Reichert T, DeBruyere M, Deneys V, Totterman T, Lydyard P, Yuksel F, et al. Lymphocyte subset reference ranges in adult Caucasians. *Clin Immunol Immunopathol* 1991;60(2):190-208.
54. Erkeller-Yuksel FM, Deneys V, Yuksel B, Hannet I, Hulstaert F, Hamilton C, et al. Age-related changes in human blood lymphocyte subpopulations. *J Pediatr* 1992;120(2 Pt 1):216-22.
55. Hulstaert F, Hannet I, Deneys V, Munhyeshuli V, Reichert T, De Bruyere M, et al. Age-related changes in human blood lymphocyte subpopulations. II. Varying kinetics of percentage and absolute count measurements. *Clin Immunol Immunopathol* 1994;70(2):152-8.
56. Kam KM, Leung WL, Kwok MY, Hung MY, Lee SS, Mak WP. Lymphocyte subpopulation reference ranges for monitoring human immunodeficiency virus-infected Chinese adults. *Clin Diagn Lab Immunol* 1996;3(3):326-30.
57. Motley D, Meyer MP, King RA, Naus GJ. Determination of lymphocyte immunophenotypic values for normal full-term cord blood. *Am J Clin Pathol* 1996;105(1):38-43.

58. Vithayasai V, Sirisanthana T, Sakonwasun C, Suvanpiyasiri C. Flow cytometric analysis of T-lymphocytes subsets in adult Thais. *Asian Pac J Allergy Immunol* 1997;15(3):141-6.
59. Comans-Bitter WM, de Groot R, van den Beemd R, Neijens HJ, Hop WC, Groeneveld K, et al. Immunophenotyping of blood lymphocytes in childhood. Reference values for lymphocyte subpopulations. *J Pediatr* 1997;130(3):388-93.
60. Shahabuddin S, Al-Ayed I, Gad El-Rab MO, Qureshi MI. Age-related changes in blood lymphocyte subsets of Saudi Arabian healthy children. *Clin Diagn Lab Immunol* 1998;5(5):632-5.
61. Ohta Y, Fujiwara K, Nishi T, Oka H. Normal values of peripheral lymphocyte populations and T cell subsets at a fixed time of day: a flow cytometric analysis with monoclonal antibodies in 210 healthy adults. *Clin Exp Immunol* 1986;64(1):146-9.
62. Miyawaki T, Taga K, Nagaoki T, Seki H, Suzuki Y, Taniguchi N. Circadian changes of T lymphocyte subsets in human peripheral blood. *Clin Exp Immunol* 1984;55(3):618-22.
63. Giorgi JV, Cheng HL, Margolick JB, Bauer KD, Ferbas J, Waxdal M, et al. Quality control in the flow cytometric measurement of T-lymphocyte subsets: the multicenter AIDS cohort study experience. The Multicenter AIDS Cohort Study Group. *Clin Immunol Immunopathol* 1990;55(2):173-86.
64. Parker JW, Adelsberg B, Azen SP, Boone D, Fletcher MA, Gjerset GF, et al. Leukocyte immunophenotyping by flow cytometry in a multisite study: standardization, quality control, and normal values in the Transfusion Safety Study. The Transfusion Safety Study Group. *Clin Immunol Immunopathol* 1990;55(2):187-220.
65. Ekong T, Kupek E, Hill A, Clark C, Davies A, Pinching A. Technical influences on immunophenotyping by flow cytometry. The effect of time and temperature of storage on the viability of lymphocyte subsets. *J Immunol Methods* 1993;164(2):263-73.

66. Hoover DR, Graham NM, Chen B, Taylor JM, Phair J, Zhou SY, et al. Effect of CD4+ cell count measurement variability on staging HIV-1 infection. *J Acquir Immune Defic Syndr* 1992;5(8):794-802.
67. Ginns LC, Goldenheim PD, Miller LG, Burton RC, Gillick L, Colvin RB, et al. T-lymphocyte subsets in smoking and lung cancer: Analysis of monoclonal antibodies and flow cytometry. *Am Rev Respir Dis* 1982;126(2):265-9.
68. Lapolla A, Sanzari MC, Zancanaro F, Masin M, Guerriero A, Piva I, et al. A study on lymphocyte subpopulation in diabetic mothers at delivery and in their newborn. *Diabetes Nutr Metab* 1999;12(6):394-9.
69. CDC. US Department of Health & Human Service. CDC guidelines for national human immunodeficiency virus case surveillance, including monitoring for human immunodeficiency virus infection and acquired immunodeficiency syndrome. Centers for Disease Control and Prevention. *MMWR* 1999;48(RR-13):1-27, 29-31.
70. CDC. US Department of Health & Human Service. 1993 revised classification system for HIV infection and expanded surveillance case definition for AIDS among adolescents and adults. *MMWR* 1992;41(RR-17):1-19.
71. CDC. US Department of Health & Human Service. 1999 USPHS/IDSA guidelines for the prevention of opportunistic infections in persons infected with human immunodeficiency virus. U.S. Public Health Service (USPHS) and Infectious Diseases Society of America (IDSA). *MMWR* 1999;48(RR-10):1-59, 61-6.
72. Webster HK, Kawai Y, Haney D, Nakahara K, Watanabe K, Ogawa E, et al. Determination of laboratory reference ranges for lymphocyte immunophenotype subsets in healthy adult Japanese using flow cytometry. *Asian Pac J Allergy Immunol* 2003;21:161-9.
73. Lee B, Yap H, Chew F, Quah T, Prabhakaran K, Chan GSH, et al. Age-and sex-related changes in lymphocyte subpopulation of healthy Asian subjects:from birth to adulthood. *Cytometry* 1996; 26:8-15.

74. Appay V, Papagno L, Spina CA, Hansasuta P, King A, Jones L, et al. Dynamics of T cell responses in HIV infection. *J Immunol* 2002;168(7):3660-6.
75. Vuillier F, Bianco NE, Montagnier L, Dighiero G. Selective depletion of low-density CD8+, CD16+ lymphocytes during HIV infection. *AIDS Res Hum Retroviruses* 1988;4(2):121-9.
76. Ullum H, Palmo J, Halkjaer-Kristensen J, Diamant M, Klokke M, Kruuse A, et al. The effect of acute exercise on lymphocyte subsets, natural killer cells, proliferative responses, and cytokines in HIV-seropositive persons. *J Acquir Immune Defic Syndr* 1994;7(11):1122-33.
77. Fuchs M, Stadler BM, Malinverni R, de Weck AL, Pichler WJ. [Relationship between T-subsets and clinical aspects of HIV-associated diseases]. *Schweiz Med Wochenschr* 1991;121(3):57-66.
78. Sampalo A, Lopez-Gomez M, Jimenez-Alonso J, Ortiz F, Samaniego F, Garrido F. CD5+ B lymphocytes in HIV infection: relationship to immunological progression of disease. *Clin Immunol Immunopathol* 1993;66(3):260-8.
79. Bryson GJ, Keegan RP, Quinn CP, Collins RJ. Evaluation of semiautomated procedure for lymphocyte subset analysis. *J Clin Pathol* 1990;43(4):337-41.
80. Douglas SD, Durako SJ, Tustin NB, Houser J, Muenz L, Starr SE, et al. Natural killer cell enumeration and function in HIV-infected and high-risk uninfected adolescents. *AIDS Res Hum Retroviruses* 2001;17(6):543-52.
81. Hu PF, Hultin LE, Hultin P, Hausner MA, Hirji K, Jewett A, et al. Natural killer cell immunodeficiency in HIV disease is manifest by profoundly decreased numbers of CD16+CD56+ cells and expansion of a population of CD16dimCD56- cells with low lytic activity. *J Acquir Immune Defic Syndr Hum Retrovirol* 1995;10(3):331-40.
82. Muller C, Szangolies M, Kukel S, Kiehl M, Sorice M, Griggi T, et al. Characterization of autoantibodies to natural killer cells in HIV-infected patients. *Scand J Immunol* 1996;43(5):583-92.

83. Ironson G, Balbin E, Solomon G, Fahey J, Klimas N, Schneiderman N, et al. Relative preservation of natural killer cell cytotoxicity and number in healthy AIDS patients with low CD4 cell counts. *Aids* 2001;15(16):2065-73.
84. de Souza MS, Karnasuta C, Brown AE, Markowitz LE, Nitayaphan S, Garner RP, et al. A comparative study of the impact of HIV infection on natural killer cell number and function in Thais and North Americans. *AIDS Res Hum Retroviruses* 2000;16(11):1061-6.
85. Jason J, Archibald L, McDonald LC, Hart WM, Rheapumikankit S, Tansuphwaswadikul S, et al. Immune determinants of organism and outcome in febrile hospitalized Thai patients with bloodstream infections. *Clin Diagn Lab Immunol* 1999;6(1):73-8.
86. Ullum H, Cozzi Lepri A, Aladdin H, Katzenstein T, Victor J, Phillips AN, et al. Natural immunity and HIV disease progression. *Aids* 1999;13(5):557-63.
87. CDC.1993. Revised classification system for HIV infection and expanded surveillance case definition for AIDS among adolescents and adults. *MMWR* 1992;41(No.RR-17):1-19.
88. Mendel A, Tenant-Flowers M. Natural history and management of early HIV infection. *BMJ* 2001;322:1290-3.
89. Delves PJ, and Roitt IM. The immune system. *N Eng J Med* 1992;327:1789-92.
90. Mackay I, and Rosen FS. Innate immunity. *N Eng J Med* 2000;343:338-44.
91. Guidotti LG, Chisari FV. Noncytolytic control of viral infections by the innate and adaptive immune response. *Annu Rev Immunol* 2001;19:65-91.
92. Paxton WA, Neumann AU, Kang S, Deutch L, Brown RC, Koup RA, et al. RANTES production from CD4+ lymphocytes correlates with host genotype and rates of human immunodeficiency virus type 1 disease progression. *J Infect Dis* 2001;183(11):1678-81.
93. Suni MA, Picker LJ, Maino VC. Detection of antigen-specific T cell cytokine expression in whole blood by flow cytometry. *J Immunol Methods* 1998;212(1):89-98.

94. McMichael AJ, Rowland-Jones SL. Cellular immune responses to HIV. *Nature* 2001;410(6831):980-7.
95. Brook GF, Butel JS, and Morse SA. *Medical microbiology*. Twenty-second ed. New York: Mc Graw-Hill companies Inc; 2001.
96. Rowland-Jones S, Tan R, McMichael A. Role of cellular immunity in protection against HIV infection. *Adv Immunol* 1997;65:277-346.
97. Borrow P, Lewicki H, Wei X, Horwitz MS, Peffer N, Meyers H, et al. Antiviral pressure exerted by HIV-1-specific cytotoxic T lymphocytes (CTLs) during primary infection demonstrated by rapid selection of CTL escape virus. *Nat Med* 1997;3(2):205-11.
98. Klein SA, Dobmeyer JM, Dobmeyer TS, Pape M, Ottmann OG, Helm EB, et al. Demonstration of the Th1 to Th2 cytokine shift during the course of HIV-1 infection using cytoplasmic cytokine detection on single cell level by flow cytometry. *AIDS* 1997;11(9):1111-8.
99. Appay V, Nixon DF, Donahoe SM, Gillespie GM, Dong T, King A, et al. HIV-specific CD8(+) T cells produce antiviral cytokines but are impaired in cytolytic function. *J Exp Med* 2000;192(1):63-75.
100. Kinter A, Arthos J, Cicala C. and Fauci AS. Chemokines, cytokine and HIV : a complex network of interactions that influence HIV pathogenesis. *Imm Rev* 2000; 177:88-98.
101. Kallas EG, Gibbons DC, Soucier H, Fitzgerald T, Treanor JJ, Evans TG. Detection of intracellular antigen-specific cytokines in human T cell populations. *J Infect Dis* 1999;179(5):1124-31.
102. Roitt I, Brostoff J, Male D. *Immunology*. 3th ed. London: Mosby-Year Book, Europe Ltd; 1993.
103. Katze MG, He Y, Gale M, Jr. Viruses and interferon: a fight for supremacy. *Nat Rev Immunol* 2002;2(9):675-87.
104. Samuel CE. Antiviral actions of interferons. *Clin Microbiol Rev* 2001;14(4):778-809, table of contents.
105. Levy DE. The virus battles:IFN induction of the antiviral state and mechanisms of viral evasion. *Cytokine Growth Factor Rev* 2001;12:143-156.

106. Callard RE, Gearing AJH. The cytokine facts book. London: Academic Press Limited; 1994.
107. Barre-Sinoussi F, Chermann JC, Rey F, Nugeyre MT, Chamaret S, Gruest J, et al. Isolation of a T-lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS). *Science* 1983;220(4599):868-71.
108. Markham PD, Salahuddin SZ, Veren K, Orndorff SH, Gallo RC. Hydrocortisone and some other hormones enhance the expression of HTLV-III. *Int J Cancer* 1986;37:67-72.
109. Voth R, Rossol S, Klein K, Hess G, Schutt KH, Schroder HC, et al. Differential gene expression of IFN-alpha and tumor necrosis factor-alpha in peripheral blood mononuclear cells from patients with AIDS related complex and AIDS. *J Immunol* 1990;144(3):970-5.
110. Krown SE, Niedzwiecki D, Bhalla RB, Flomenberg N, Bundow D, Chapman D. Relationship and prognostic value of endogenous interferon-alpha, beta 2-microglobulin, and neopterin serum levels in patients with Kaposi sarcoma and AIDS. *J Acquir Immune Defic Syndr* 1991;4(9):871-80.
111. Grunfeld C, Kotler DP, Shigenaga JK, Doerrler W, Tierney A, Wang J, et al. Circulating interferon-alpha levels and hypertriglyceridemia in the acquired immunodeficiency syndrome. *Am J Med* 1991;90(2):154-62.
112. Oka S, Hirabayashi Y, and Mouri H. Beta interferon and early stage HIV infection. *J AIDS* 1989;2:125-8.
113. Bach EA, Aguet M, Schreiber RD. The IFN gamma receptor: a paradigm for cytokine receptor signaling. *Annu Rev Immunol* 1997;15:563-91.
114. Boehm U, Klamp T, Groot M, Howard JC. Cellular responses to interferon-gamma. *Annu Rev Immunol* 1997;15:749-95.
115. Mosmann TR, Coffman RL. TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annu Rev Immunol* 1989;7:145-73.

116. Poli G, Introna M, Zanaboni F, Peri G, Carbonari M, Aiuti F, et al. Natural killer cells in intravenous drug abusers with lymphadenopathy syndrome. *Clin Exp Immunol* 1985;62(1):128-35.
117. Sukwit S, Chuenchitra T, Samakoses R, Songprasom K, Aree C, Akapirat S, et al. Flow cytometric detection of intracellular cytokines in peripheral blood of HIV-1 infected Thai children. *Asian Pac J Allergy Immunol* 2001;19(2):107-13.
118. Vyakarnam A, Matear P, Meager A, Kelly G, Stanley B, Weller I, et al. Altered production of tumour necrosis factors alpha and beta and interferon gamma by HIV-infected individuals. *Clin Exp Immunol* 1991;84(1):109-15.
119. Azzoni L, Papasavvas E, Chehimi J, Kostman JR, Mounzer K, Ondercin J, et al. Sustained impairment of IFN-gamma secretion in suppressed HIV-infected patients despite mature NK cell recovery: evidence for a defective reconstitution of innate immunity. *J Immunol* 2002;168(11):5764-70.
120. Nokta MA, Pollard RB. Patterns of interferon-gamma production by peripheral blood mononuclear cells from patients with human immunodeficiency virus infection. *J Interferon Res* 1990;10(2):173-81.
121. Taub DD. Chemokine-leukocyte interactions. The voodoo that they do so well. *Cytokine Growth Factor Rev* 1996;7(4):355-76.
122. Baggiolini M, Dewald B, Moser B. Human chemokines: an update. *Annu Rev Immunol* 1997;15:675-705.
123. Gao JL, Kuhns DB, Tiffany HL, McDermott D, Li X, Francke U, et al. Structure and functional expression of the human macrophage inflammatory protein 1 alpha/RANTES receptor. *J Exp Med* 1993;177(5):1421-7.
124. Wolpe SD, Cerami A. Macrophage inflammatory proteins 1 and 2: members of a novel superfamily of cytokines. *Faseb J* 1989;3(14):2565-73.
125. Cocchi F, DeVico AL, Garzino-Demo A, Arya SK, Gallo RC, Lusso P. Identification of RANTES, MIP-1 alpha, and MIP-1 beta as the major HIV-suppressive factors produced by CD8+ T cells. *Science* 1995;270(5243):1811-5.

126. Deng H, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhart M, et al. Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 1996;381(6584):661-6.
127. Dragic T, Litwin V, Allaway GP, Martin SR, Huang Y, Nagashima KA, et al. HIV-1 entry into CD4+ cells is mediated by the chemokine receptor CC-CKR-5. *Nature* 1996;381(6584):667-73.
128. Alkhatib G, Combadiere C, Broder CC, Feng Y, Kennedy PE, Murphy PM, et al. CC CKR5: a RANTES, MIP-1alpha, MIP-1beta receptor as a fusion cofactor for macrophage-tropic HIV-1. *Science* 1996;272(5270):1955-8.
129. Choe H, Farzan M, Sun Y, Sullivan N, Rollins B, Ponath PD, et al. The beta-chemokine receptors CCR3 and CCR5 facilitate infection by primary HIV-1 isolates. *Cell* 1996;85(7):1135-48.
130. Doranz BJ, Rucker J, Yi Y, Smyth RJ, Samson M, Peiper SC, et al. A dual-tropic primary HIV-1 isolate that uses fusin and the beta-chemokine receptors CKR-5, CKR-3, and CKR-2b as fusion cofactors. *Cell* 1996;85(7):1149-58.
131. Nagasawa T, Tachibana K, Kishimoto T. A novel CXC chemokine PBSF/SDF-1 and its receptor CXCR4: their functions in development, hematopoiesis and HIV infection. *Semin Immunol* 1998;10(3):179-85.
132. Feng Y, Broder CC, Kennedy PE, Berger EA. HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane, G protein-coupled receptor. *Science* 1996;272:872-7.
133. Ferbas J, Giorgi JV, Amini S, Grovit-Ferbas K, Wiley DJ, Detels R, et al. Antigen-specific production of RANTES, macrophage inflammatory protein (MIP)-1alpha, and MIP-1beta in vitro is a correlate of reduced human immunodeficiency virus burden in vivo. *J Infect Dis* 2000;182(4):1247-50.
134. Furci L, Scarlatti G, Burastero S, Tambussi G, Colognesi C, Quillent C, et al. Antigen-driven C-C chemokine-mediated HIV-1 suppression by CD4 (+) T cells from exposed uninfected individuals expressing the wild-type CCR-5 allele. *J Exp Med* 1997;186(3):455-60.

135. Rosenberg ES, Billingsley JM, Caliendo AM, Boswell SL, Sax PE, Kalams SA, et al. Vigorous HIV-1-specific CD4⁺ T cell responses associated with control of viremia. *Science* 1997;278(5342):1447-50.
136. Garzino-Demo A, Moss RB, Margolick JB, Cleghorn F, Sill A, Blattner WA, et al. Spontaneous and antigen-induced production of HIV-inhibitory beta-chemokines are associated with AIDS-free status. *Proc Natl Acad Sci U S A* 1999;96(21):11986-91.
137. Saha K, Bentsman G, Chess L, Volsky DJ. Endogenous production of beta-chemokines by CD4⁺, but not CD8⁺, T-cell clones correlates with the clinical state of human immunodeficiency virus type 1 (HIV-1)-infected individuals and may be responsible for blocking infection with non-syncytium-inducing HIV-1 in vitro. *J Virol* 1998;72(1):876-81.
138. Ullum H, Cozzi Lepri A, Victor J, Aladdin H, Phillips AN, Gerstoft J, et al. Production of beta-chemokines in human immunodeficiency virus (HIV) infection: evidence that high levels of macrophage inflammatory protein-1beta are associated with a decreased risk of HIV disease progression. *J Infect Dis* 1998;177(2):331-6.
139. Roger M. Influence of host genes on HIV-1 disease progression. *Faseb J* 1998;12(9):625-32.
140. Kinter AL, Ostrowski M, Goletti D, Oliva A, Weissman D, Gantt K, et al. HIV replication in CD4⁺ T cells of HIV-infected individuals is regulated by a balance between the viral suppressive effects of endogenous beta-chemokines and the viral inductive effects of other endogenous cytokines. *Proc Natl Acad Sci U S A* 1996;93(24):14076-81.
141. Verani A, Scarlatti G, Comar M, Tresoldi E, Polo S, Giacca M, et al. C-C chemokines released by lipopolysaccharide (LPS)-stimulated human macrophages suppress HIV-1 infection in both macrophages and T cells. *J Exp Med* 1997;185(5):805-16.
142. Dallman MJ, Montgomery RA, Larsen CP, Wanders A, Wells AF. Cytokine gene expression: analysis using northern blotting, polymerase chain reaction and in situ hybridization. *Immunol Rev* 1991;119:163-79.

143. Power CA, Grand CL, Ismail N, Peters NC, Yurkowski DP, Bretscher PA. A valid ELISPOT assay for enumeration of ex vivo, antigen-specific, IFN γ -producing T cells. *J Immunol Methods* 1999;227(1-2):99-107.
144. Czerkinsky C, Andersson G, Ekre HP, Nilsson LA, Klareskog L, Ouchterlony O. Reverse ELISPOT assay for clonal analysis of cytokine production. I. Enumeration of gamma-interferon-secreting cells. *J Immunol Methods* 1988;110(1):29-36.
145. Waldrop SL, Pitcher CJ, Peterson DM, Maino VC, Picker LJ. Determination of antigen-specific memory/effector CD4⁺ T cell frequencies by flow cytometry: evidence for a novel, antigen-specific homeostatic mechanism in HIV-associated immunodeficiency. *J Clin Invest* 1997;99(7):1739-50.
146. Cooper MA, Fehniger TA, Caligiuri MA. The biology of human natural killer-cell subsets. *Trends Immunol* 2001;22(11):633-40.
147. Trinchieri G. Biology of natural killer cells. *Adv Immunol* 1989;47:187-376.
148. Biron CA, Nguyen KB, Pien GC, Cousens LP, Salazar-Mather TP. Natural killer cells in antiviral defense: function and regulation by innate cytokines. *Annu Rev Immunol* 1999;17:189-220.
149. See DM, Khemka P, Sahl L, Bui T, Tilles JG. The role of natural killer cells in viral infections. *Scand J Immunol* 1997;46(3):217-24.
150. Barao I, Ascensao JL. Human natural killer cells. *Arch Immunol Ther Exp (Warsz)* 1998;46(4):213-29.
151. Yokoyama WM. Natural killer cells. 4th ed. Philadelphia: Lippincott-Raven; 1999.
152. Shea JO, and Ortaldo JR. The biology of natural killer cells: insights into the molecular basis of function. Oxford: University Press; 1992.
153. Rutishauser U, Acheson A, Hall AK, Mann DM, Sunshine J. The neural cell adhesion molecule (NCAM) as a regulator of cell-cell interactions. *Science* 1988;240(4848):53-7.

154. Winnock M, Garcia Barcina M, Lukomska B, Huet S, Saric J, Balabaud C, et al. Human liver-associated lymphocytes: an overview. *J Gastroenterol Hepatol* 1995;10 Suppl 1:S43-6.
155. Leibson PJ. Signal transduction during natural killer cell activation: inside the mind of a killer. *Immunity* 1997;6(6):655-61.
156. Solana R, Mariani E. NK and NK/T cells in human senescence. *Vaccine* 2000;18(16):1613-20.
157. Godfrey DI, Kronenberg M. Going both ways: immune regulation via CD1d-dependent NKT cells. *J Clin Invest* 2004;114(10):1379-88.
158. Chan WL, Pejnovic N, Liew TV, Lee CA, Groves R, Hamilton H. NKT cell subsets in infection and inflammation. *Immunol Lett* 2003;85(2):159-63.
159. Biron CA, Brossay L. NK cells and NKT cells in innate defense against viral infections. *Curr Opin Immunol* 2001;13(4):458-64.
160. MacDonald HR. Development and selection of NKT cells. *Curr Opin Immunol* 2002;14(2):250-4.
161. Skold M, Behar SM. Role of CD1d-restricted NKT cells in microbial immunity. *Infect Immun* 2003;71(10):5447-55.
162. van der Vliet HJ, Molling JW, von Blomberg BM, Nishi N, Kolgen W, van den Eertwegh AJ, et al. The immunoregulatory role of CD1d-restricted natural killer T cells in disease. *Clin Immunol* 2004;112(1):8-23.
163. Yoshimoto T, Paul WE. CD4pos, NK1.1pos T cells promptly produce interleukin 4 in response to in vivo challenge with anti-CD3. *J Exp Med* 1994;179(4):1285-95.
164. van der Vliet HJ, von Blomberg BM, Hazenberg MD, Nishi N, Otto SA, van Benthem BH, et al. Selective decrease in circulating V alpha 24+V beta 11+ NKT cells during HIV type 1 infection. *J Immunol* 2002;168(3):1490-5.
165. Motsinger A, Haas DW, Stanic AK, Van Kaer L, Joyce S, Unutmaz D. CD1d-restricted human natural killer T cells are highly susceptible to human immunodeficiency virus 1 infection. *J Exp Med* 2002;195(7):869-79.
166. Unutmaz D. NKT cells and HIV infection. *Microbes Infect* 2003;5(11):1041-7.

167. Pisegna S, Pirozzi G, Piccoli M, Frati L, Santoni A, Palmieri G. p38 MAPK activation controls the TLR3-mediated up-regulation of cytotoxicity and cytokine production in human NK cells. *Blood* 2004;104(13):4157-64.
168. Schmidt KN, Leung B, Kwong M, Zarembek KA, Satyal S, Navas TA, et al. APC-independent activation of NK cells by the Toll-like receptor 3 agonist double-stranded RNA. *J Immunol* 2004;172(1):138-43.
169. Sivori S, Falco M, Della Chiesa M, Carlomagno S, Vitale M, Moretta L, et al. CpG and double-stranded RNA trigger human NK cells by Toll-like receptors: induction of cytokine release and cytotoxicity against tumors and dendritic cells. *Proc Natl Acad Sci U S A* 2004;101(27):10116-21.
170. Lewis CE, and McGee JOD. The natural immune system (the natural killer cell). Oxford United States: Oxford University Press; 1992.
171. Fanger MW, Shen L, Graziano RF, Guyre PM. Cytotoxicity mediated by human Fc receptors for IgG. *Immunol Today* 1989;10(3):92-9.
172. Robertson MJ, Ritz J. Biology and clinical relevance of human natural killer cells. *Blood* 1990;76(12):2421-38.
173. Welsh RM, Brubaker JO, Vargas-Cortes M, O'Donnell CL. Natural killer (NK) cell response to virus infections in mice with severe combined immunodeficiency. The stimulation of NK cells and the NK cell-dependent control of virus infections occur independently of T and B cell function. *J Exp Med* 1991;173(5):1053-63.
174. Biron CA. Activation and function of natural killer cell responses during viral infections. *Curr Opin Immunol* 1997;9(1):24-34.
175. Rungrueangthakit KA, Duerr V, and Suriyanon L. Immune parameters among highly expressed persistently seronegative (HEPS) Thai women: In: Abstract of the XI Int conference on AIDS; Mo.A.4019 (Abstr.) 1:13.
176. Ratcliffe LT, Lukey PT, MacKenzie CR, Ress SR. Reduced NK activity correlates with active disease in HIV- patients with multidrug-resistant pulmonary tuberculosis. *Clin Exp Immunol* 1994;97(3):373-9.

177. Brenner BG, Gornitsky M, Wainberg MA. Interleukin-2-inducible natural immune (lymphokine-activated killer cell) responses as a functional correlate of progression to AIDS. *Clin Diagn Lab Immunol* 1994;1(5):538-44.
178. Schwartz O, Marechal V, Le Gall S, Lemonnier F, Heard JM. Endocytosis of major histocompatibility complex class I molecules is induced by the HIV-1 Nef protein. *Nat Med* 1996;2(3):338-42.
179. Bonavida B, Katz J, Gottlieb M. Mechanism of defective NK cell activity in patients with acquired immunodeficiency syndrome (AIDS) and AIDS-related complex. I. Defective trigger on NK cells for NKCF production by target cells, and partial restoration by IL 2. *J Immunol* 1986;137(4):1157-63.
180. Cai Q, Huang XL, Rappocciolo G, Rinaldo CR, Jr. Natural killer cell responses in homosexual men with early HIV infection. *J Acquir Immune Defic Syndr* 1990;3(7):669-76.
181. Chehimi J, Starr SE, Frank I, Rengaraju M, Jackson SJ, Llanes C, et al. Natural killer (NK) cell stimulatory factor increases the cytotoxic activity of NK cells from both healthy donors and human immunodeficiency virus-infected patients. *J Exp Med* 1992;175(3):789-96.
182. Scott-Algara D, Vuillier F, Cayota A, Dighiero G. Natural killer (NK) cell activity during HIV infection: a decrease in NK activity is observed at the clonal level and is not restored after in vitro long-term culture of NK cells. *Clin Exp Immunol* 1992;90(2):181-7.
183. Bryant J, Day R, Whiteside TL and Herberman RB. Calculation of lytic units for the expression of cell-mediated cytotoxicity. *J Immunol Methods* 1992;146:91-103.
184. Cox JH. Evaluation of natural killer cell activity. In Micheal NL, Kim JH, editors. *Methods in molecular medicine: HIV protocol*. NJ: Humana Press Inc; 1999.

185. Chehimi J, Bandyopadhyay S, Prakash K, Perussia B, Hassan NF, Kawashima H, et al. In vitro infection of natural killer cells with different human immunodeficiency virus type 1 isolates. *J Virol* 1991;65(4):1812-22.
186. Scott-Algara D, Vuillier F, Cayota A, Rame V, Guetard D, Moncany ML, et al. In vitro non-productive infection of purified natural killer cells by the BRU isolate of the human immunodeficiency virus type 1. *J Gen Virol* 1993;74 (Pt 4):725-31.
187. Toth FD, Mosborg-Petersen P, Kiss J, Aboagye-Mathiesen G, Zdravkovic M, Hager H, et al. Differential replication of human immunodeficiency virus type 1 in CD8- and CD8+ subsets of natural killer cells: relationship to cytokine production pattern. *J Virol* 1993;67(10):5879-88.
188. Reyburn H, Mandelboim O, Vales-Gomez M, Sheu EG, Pazmany L, Davis DM, et al. Human NK cells: their ligands, receptors and functions. *Immunol Rev* 1997;155:119-25.
189. Mendes R, Bromelow KV, Westby M, Galea-Lauri J, Smith IE, O'Brien ME, et al. Flow cytometric visualisation of cytokine production by CD3-CD56+ NK cells and CD3+CD56+ NK-T cells in whole blood. *Cytometry* 2000;39(1):72-8.
190. Bluman EM, Bartynski KJ, Avalos BR, Caligiuri MA. Human natural killer cells produce abundant macrophage inflammatory protein-1 alpha in response to monocyte-derived cytokines. *J Clin Invest* 1996;97 (12):2722-7.
191. Fehniger TA, Herbein G, Yu H, Para MI, Bernstein ZP, O'Brien WA, et al. Natural killer cells from HIV-1+ patients produce C-C chemokines and inhibit HIV-1 infection. *J Immunol* 1998;161(11):6433-8.
192. Scharton TM, Scott P. Natural killer cells are a source of interferon gamma that drives differentiation of CD4+ T cell subsets and induces early resistance to *Leishmania major* in mice. *J Exp Med* 1993;178(2):567-77.

193. Maino VC, Picker LJ. Identification of functional subsets by flow cytometry: intracellular detection of cytokine expression. *Cytometry* 1998;34(5):207-15.
194. Jung T, Schauer U, Heusser C, Neumann C, Rieger C. Detection of intracellular cytokines by flow cytometry. *J Immunol Methods* 1993;159(1-2):197-207.
195. Picker LJ, Singh MK, Zdraveski Z, Treer JR, Waldrop SL, Bergstresser PR, et al. Direct demonstration of cytokine synthesis heterogeneity among human memory/effector T cells by flow cytometry. *Blood* 1995;86(4):1408-19.
196. Prussin C, Metcalfe DD. Detection of intracytoplasmic cytokine using flow cytometry and directly conjugated anti-cytokine antibodies. *J Immunol Methods* 1995;188(1):117-28.
197. Peruzzi M, Azzari C, Rossi ME, De Martino M, Vierucci A. Inhibition of natural killer cell cytotoxicity and interferon gamma production by the envelope protein of HIV and prevention by vasoactive intestinal peptide. *AIDS Res Hum Retroviruses* 2000;16(11):1067-73.
198. Goicoa MA, Sen L, Iannitelli PS, Diez RA, Estevez ME. Natural killer suppressor factors in sera from HIV-infected subjects. *Scand J Immunol* 1995;41(6):523-8.
199. Cauda R, Tumbarello M, Ortona L, Kanda P, Kennedy RC, Chanh TC. Inhibition of normal human natural killer cell activity by human immunodeficiency virus synthetic transmembrane peptides. *Cell Immunol* 1988;115(1):57-65.
200. Bancroft GJ. The role of natural killer cells in innate resistance to infection. *Curr Opin Immunol* 1993;5(4):503-10.
201. Robinson WE, Jr., Mitchell WM, Chambers WH, Schuffman SS, Montefiori DC, Oeltmann TN. Natural killer cell infection and inactivation in vitro by the human immunodeficiency virus. *Hum Pathol* 1988;19(5):535-40.

202. Feremans WW, Huygen K, Menu R, Farber CM, de Caluwe JP, van Vooren JP, et al. Fifty cases of human immunodeficiency virus (HIV) infection: immunoultrastructural study of circulating lymphocytes. *J Clin Pathol* 1988;41(1):62-71.

APPENDIX

APPENDIX

A. Reagents for cell cultivation

A.1 Growth media for K-562 erythroleukemic cell line

10% fetal calf serum (FCS) (GIBCO BRL®, Grand Island, New York) in RPMI 1640 (GIBCO, USA) plus 100 units/ml penicillin and 100 µg/ml streptomycin.

A.1.1 5X RPMI 1640

Dissolve RPMI 1640 powder (GIBCO) with deionized tri-distilled water to make 5X solution; and sterile through filtration with 0.22 µm millipore membrane. The 5X media is frozen at -20°C until used.

A.1.2 1X RPMI 1640 (500 ml)

5X RPMI 1640	100	ml
1M N-2 hydroxyethylpiperazine-2-ethane sulfonic acid (HEPES)		
	1	ml
Penicillin (40,000 units/ml) (GIBCO)	1.25	ml
Streptomycin (40,000 units/ml) (GIBCO)	1.25	ml

Adjust to pH 7.2-7.4 with 5% NaHCO₃ and add deionized, tri-distilled water to 500 ml

A.1.3 10% FBS in 1X RPMI 1640 (growth media)

1X RPMI 1640	450	ml
Fetal bovine serum (Biochrom KG, Berlin, Germany)	50	ml

A.2 Cell freezing media (10% dimethylsulfoxide in 20% FBS RPMI 1640)**40% FBS in 1X RPMI 1640**

1X RPMI 1640	6	ml
FBS	4	ml

20% DMSO in 1X RPMI 1640

1X RPMI 1640	8	ml
DMSO (Sigma, St. Louis, Mo, USA)	2	ml

Working protocol

Suspend cells in cold 40% FBS RPMI 1640.

Add 20% DMSO in RPMI 1640 dropwise at an equal volume to achieve the final concentration of 10% DMSO and 20% FBS in RPMI media

A.3 Phosphate buffer saline Ca^{2+} and Mg^{2+} free (1X), pH 7.5 for cell washing, flow cytometry

NaCl	8	g
KCl	0.20	g
KH_2PO_4 (anhydrous)	0.12	g
Na_2HPO_4 (anhydrous)	0.91	g
Deionized, tri-distilled water	1,000	ml

Adjust to pH 7.5 by 1N NaOH. Sterilize by autoclaving at 121°C under pressure of 15 lbs/square inch for 15 minutes, then, store at room temperature

A.4 1M HEPES buffer

HEPES	253.3	g
Deionized, distilled water	1000	ml

A.4.1 Dissolve 253.3 g HEPES with deionized, distilled water to 1000 ml

A.4.2 Adjust to pH 7.4 with 1N NaOH

A.4.3 Sterilize through filtration with 0.22 μ m millipore membrane

A.5 0.4% Trypan blue

This reagent is used for dead cell staining in the process of cell counting by light microscope

Trypan blue dye (Sigma)	0.4	g
1X PBS	100	ml

A.5.1 Dissolve thoroughly 0.4 g trypan blue dye with 100 ml of 1X PBS

A.5.2 Filter the solution through Whatman filter paper and keep at room temperature

B. Reagents for flow cytometry**B.1 1% Paraformaldehyde**

Paraformaldehyde (sigma)	1	g
1X PBS	100	ml

Filter through 0.1 μ m filtered paper, store in the dark bottle at 4 °C.

B.2 FACS lysing solution

This reagent is used for lysing the red blood cells following direct immunofluorescence staining of human peripheral blood cells with monoclonal antibodies prior to flow cytometric analysis.

Working 1X FACS lysing solution

10X FACS lysing solution (Becton Dickson system, San Jose, CA, USA)	1	ml
Deionized, distilled water	9	ml

The prepared solution is stable for up to month when stored at room temperature.

B.3 FACS permeabilizing solution

This reagent is used for permeabilization of lymphocyte membrane prior to intracellular immunofluorescence staining with monoclonal antibodies.

Working FACS permeabilizing solution (1X)

10X concentrated FACS permeabilizing solution (Becton Dickinson)	1	ml
Deionized, distilled water	9	ml

Dilute concentrated FACS permeabilizing solution to 1:10 with deionized, distilled water. The prepared solution is stable for up to one month when stored at room temperature. Do not use this reagent if discoloration or precipitation is seen .

B.4 Phorbol 12-myristate 13-acetate (PMA)

This reagent is used to stimulate the cells to produce cytokine.

PMA stock solution (0.1 mg/ml)

PMA powder (Sigma)	1	mg
DMSO	10	ml

Store the stock PMA solution in small aliquots (e.g., 20 μ l) at -20°C ; do not repeat freeze-thaw cycle

PMA working solution (1 ng/ml)

Stock PMA solution	2	μl
Steriled PBS	198	μl

PMA working solution is used at concentration of 25 μl / 1ml whole blood or 12.5 μl of PMA/ 500 μl of whole blood to obtain the final concentration of 25 ng/ml

B.5 Ionomycin

This reagent is used to stimulate the cells to produce cytokine.

Ionomycin stock solution (0.5 mg/ml)

Ionomycin powder (Sigma)	1	mg
Ethanol (gold shield ethyl alcohol, 200 proof)	2	ml

Store the stock solution in small aliquots (e.g.,20μl) at -20 °C.

Ionomycin working solution (0.05 mg/ml)

Stock ionomycin solution	10	μl
Steriled PBS	90	μl

Ionomycin working is used at concentration of 20 μl/ ml of whole blood or 10 μl of ionomycin/ 500μl of whole blood to obtain the final concentration of 1μg/ ml

B.6 Brefeldin A (Sigma B-7651)

This reagent is used for blocking an intracellular protein release out of the cells during in vitro cell activation process.

Brefeldin A stock solution (5 mg/ml)

Brefeldin A powder (γ,4-Dihydroxy-2- [6-hydroxy-1-heptenyl] - 4 - cyclopentane-crotonic acid -λ lactone) (Sigma)	5	mg
DMSO	1	ml

Store the stock solution in small aliquots (e.g.,20 μl) at -20 °C ; do not repeat freeze-thaw cycle.

Brefeldin A working solution (0.5 mg/ml)

Stock Brefeldin A solution	20	μl
Sterile PBS	180	μl

BFA working solution is used at concentration of 0.5 μg/ μl of whole blood (5 μl BFA plus 500 μl whole blood) to achieve the final concentration of 10μg/ ml

C. Reagents for PBMC preparation**C.1 Standard Ficoll-hypaque (Robbins Scientific, Sunnyvale, CA, Norway)**

This reagent is used for PBMCs preparation from the EDTA blood. The bottle should be wrapped with aluminium foil and stored in a refrigerator.

D. Reagents for NK cytotoxicity assay**D.1 Sodium chromate [$\text{Na}_2^{51}\text{CrO}_4$ (^{51}Cr)]**

Just before use, dilute $\text{Na}_2^{51}\text{CrO}_4$ to the concentration of 1 mCi/ml of sterilized PBS. The amount of 200 μCi of ^{51}Cr per 1×10^6 K562 cells is required. The volume of ^{51}Cr is to be adjusted in each experiment according to the decay of the radioisotope.

D.2 10% Sodium dodecyl sulfate (SDS)

Purchase ready-made and sterile SDS from GIBCO (Gaithersburg, MD, USA). SDS has a long shelf life at room temperature.

BIOGRAPHY

NAME	MR. Sujin Assawawitoontip
DATE OF BIRTH	3 March 1970
PLACE OF BIRTH	Surin, Thailand
INSTITUTE ATTENDED	Mahidol University, 1992 Bachelor of Medical Technology Mahidol University, Doctor of Philosophy (Medical Microbiology)
PUBLICATION	Lymphocyte and NK cell subpopulations in HIV seronegative Thais. Asian Pac J Allergy Immunol 2003: 21; 95-103.
SCHOLARSHIP	The Royal Golden Jubilee Scholarships for Ph.D. Program, The Thailand Research Fund
RESEARCH ACTIVITY	Lymphocyte immunophenotype reference ranges in healthy Thai adults 1999. The Ninth Scientific Annual Meeting, 12 Nov 1999, The Virology Association (Thailand) Enumeration of natural killer cells and its activity as assayed in HIV-1 seronegative and HIV-1 infected persons. Siriraj Scientific Congress, 8 March 2000, Faculty of Medicine Siriraj Hospital Enumeration of natural killer cells and its activity as assayed in HIV-1 seronegative and HIV-1 infected persons. 5 th ASIA Pacific Congress of Medical Virology, 27 June 2000, Bali, Indonesia Natural killer cell numbers and their function in HIV-1 infected Thais. Second Symposium on Graduated Research Mahidol University, 27 April 2001, Faculty of Science, Mahidol University
HOME ADDRESS	264, Soi Bormrajchonnanee 4, Bang-plud, District, Bangkok, Thailand
POSITION&OFFICE	Faculty of Medical Technology, Mahidol University, Bangkok, Thailand 10700 Position : Medical Technologist