

เอกสารอ้างอิง

- ACGIH. Threshold limit values for chemical substances in the work environment adopted by ACGIH. *American Conference of Governmental Industrial Hygienists* 1985.
- Adam EM et al. Vapour toxicity of 1,1,1-trichloroethane (methyl chloroform) determined by experiments on laboratory animals. *Arch Ind Hyg Occup Med* 1950; 1: 225-236.
- Andrews LS, Snyder R. Toxic effects of solvents and vapors. In: Amdur MO, Doull J, Klaassen CD, eds. Casarett and Doull's Toxicology, 4th ed., pp 681-696. U.S.A., McGraw-Hill, 1992.
- Arito H, Takahashi M, Sotoyama M, Isuruta H, Ishikawa T. Electroencephalographic and autonomic responses to trichloroethylene inhalation in freely moving rats. *Arch Toxicol* 1993; 67: 193-199.
- Bai CL, Canfield PJ, Stacey NH. Individual serum bile acids as early indicators of carbon tetrachloride- and chloroform-induced liver injury. *Toxicology* 1992; 75: 221-234.
- Bai CL, Stacey NH. Mechanism of trichloroethylene-induced elevation of individual serum bile acids. II. In vitro and in vivo interference by trichloroethylene with bile acid transport in isolated rat hepatocytes. *Toxicol Appl Pharmacol* 1993; 121: 296-302.
- Barber-Riley G, Goetzee AE, Richards TG, Thomson JY. The transfer of bromsulphthalein from the plasma to the bile in man. *Clin Sci* 1961; 20: 149-159.
- Barret L, Torch S, Leray CL, Sarlieve L, Saxod R. Morphometric and

- biochemical studies in trigeminal nerve of rat after trichloroethylene or dichloroacetylene oral administration. *Neurotoxicology* 1992; 13: 601-614.
- Barret L et al. Trichloroethylene occupational exposure : elements for better prevention. *Int Arch Occup Environ Health* 1984; 53: 283-289.
- Berger ML, Sozeri T. Rapid halogenated hydrocarbon toxicity in isolated hepatocytes is mediated by direct solvent effects. *Toxicology* 1987; 45: 319-330.
- Bergmeyer HU et al. Recommendations of the german society for clinical chemistry. *Z Klin Chem Klin Biochem* 1970; 8: 659.
- Berk PD, Blaschke TF, Waggoner JG. Defective bromosulfophthalein clearance in patients with constitutional hepatic dysfunction (Gilbert ' s syndrome). *Gastroenterology* 1972; 63: 472-481.
- Berk PD, Potter BJ, Stremmel W. Role of plasma membrane ligand-binding proteins in the hepatocellular uptake of albumin-bound organic anions. *Hepatology* 1987; 7: 165-176.
- Buben JA, O'Flaherty EJ. Delineation of the role of metabolism in the hepatotoxicity of trichloroethylene and perchloroethylene : a dose - effect study. *Toxicol appl Pharmacol* 1985; 78: 105-122.
- Bull RJ, Templin M, Larson JL, Stevens DK. The role of dichloroacetate in the hepatocarcinogenicity of trichloroethylene. *Toxicol Lett* 1993; 68: 203-11.
- Cagen SZ, Klaassen CD. Evaluation of hepatic storage of sulfobromophthalien in rats and dogs. *Toxicology* 1982; 25: 261-270.
- Cherrick GR, Stein SW, Leevy CM, Davidson CS. Indocyanine green: observations on its physical properties, plasma decay and

- hepatic extraction. *J Clin Invest* 1960; 39: 592-600.
- Commission of the European Communities. 1,1,1-Trichloroethane, Trichloroethylene. In: Royal Society of Chemistry. Organo - chlorine Solvents. Health Risk to Workers, pp 55-71, 93-130. Luxembourg, 1986.
- Cornelius CE. Liver function tests in the differential diagnosis of hepatotoxicity. In: Meeks RG, Harrison SD, Bull RJ, eds. Hepatotoxicology, pp 181-213. U.S.A., CRC Press, 1991.
- Cosby NC, Dukelow WR. Toxicology of maternally ingested trichloroethylene (TCE) on embryonal and fetal development in mice and of TCE metabolites on in vitro fertilization. *Fundam Appl Toxicol* 1992; 19: 268-74.
- Cullen JM, Ruebner BH. A histopathologic classification of chemical - induced injury of the liver. In: Meeks RG, Harrison SD, Bull RJ, eds. Hepatotoxicology. U.S.A.: CRC Press, 1991.
- Cunningham ML, Gandolfi AJ, Brendel K, Sipes IG. Covalent binding of halogenated volatile solvents to subcellular macromolecules in hepatocytes. *Life Sciences* 1981; 29: 1207-1212.
- Dallas CE, Ramanathan R, Muralidhara S, Gallo JM, Bruckner IV. The uptake and elimination of 1,1,1-trichloroethane during and following inhalation exposures in rats. *Toxicol Appl Pharmacol* 1989; 98: 385-397.
- Davidson JW, Beliles, RP. Consideration of the target organ toxicity of trichloroethylene in terms of metabolite toxicity and pharmacokinetics. *Drug Metab Rev* 1991; 23: 493-599.
- Dees C, Travis C. The mitogenic potential of trichloroethylene in B6C3F1 mice. *Toxicol Lett* 1993; 69: 129-137.
- Diamandi AP, Diamandis EP, Siskos PA. Direct enzymic fluorimetric method for the determination of individual bile acids in bile.

Clinica Chimica Acta 1983; 134: 17-23.

Dossing M, Skinhoj P. Occupational liver injury. Review article. *Int Arch Occup Environ Health* 1985; 56: 1-21.

Dossing M, Ranek L. Isolated liver damage in chemical workers. *Br J Ind Med* 1984; 41: 142-144.

Dossing M, Soborg PA, Petersen LM, Ranek L. Liver damage associated with occupational exposure to organic solvents in house painters. *Eur J Clin Invest* 1983; 13: 151-157.

Driscoll TR, Hamdan HH, Wang G, Wright PFA, Stacey NH. Concentrations of individual serum or plasma bile acids in workers exposed to chlorinated aliphatic hydrocarbons. *Br J Ind Med* 1992, 49: 700-705.

Editorial. Serum bile acids in hepatobiliary disease. *Lancet* 1982; 2: 1136-1138.

Edling C, Tagerson C. Raised serum bile acid concentrations after occupational exposure to styrene : a possible sign of hepatotoxicity ? *Br J Ind Med* 1984; 41: 257-259.

Elcombe CR, Rose MS, Pratt IS. Biochemical, histological and ultrastructural changes in rat and mouse liver following the administration of trichloroethylene : possible relevance to species differences in hepatocarcinogenicity. *Toxicol Appl Pharmacol* 1985; 79: 365-76.

Elofsson SA. et al. Exposure to organic solvents. *Scand J Work Environ Health* 1980; 6: 239-273.

Enzyme Nomenclature-Recommendation (1964) of the International Union of Biochemistry. New York Elsevier 1965.

Fausa O, Gjone E. Serum bile acid concentrations in patients with liver disease. *Scand J Gastroent* 1976; 11: 537-543.

Fernicola C, Govoni S, Coniglio L, Daniele E, Trabucchi M. The use of

- 1,1,1-trichloroethane (methyl chloroform) in industrial operation : the neurotoxicity risk. *Med Lav* 1991; 82: 38-49.
- Fielder RJ, Williams SD, Sorrie GS, Bishop CM, Chandra FA, Gompertz D, Lucas EG, Van Dan Heuvel MJ, Mackay CJ. 1,1,1-Trichloroethane. Toxicity Review 9. Health and Safety Executive. London: Her Majesty's Stationary Office, 1984: 1-46.
- Fielder RJ et al. Trichloroethylene. Toxicity Review 6. Health and Safety Executive. London: Her Majesty's Stationary Office, 1982.
- Fischbein, A. et al. Industrial solvents and the liver. *Lancet* 1983; 15: 129.
- Forkert PG, Birch DW. Pulmonary toxicity of trichloroethylene in mice. Covalent binding and morphological manifestations. *Drug Metab Dispos Biol Fate Chem* 1989; 17: 106-13.
- Forkert PG, Sylvestre PL, Poland JS. Lung injury induced by trichloroethylene. *Toxicology* 1985; 35: 143-60.
- Fort DL, Stover EL, Rayburn JR, Hull M, Bantle JA. Evaluation of the developmental toxicity of trichloroethylene and detoxification metabolites using *Xenopus*. *Teratog Carcinog Mutagen* 1993; 13: 35-45.
- Franco G. New perspectives in biomonitoring liver function by means of serum bile acids: experimental and hypothetical biochemical basis. *Br J Ind Med* 1991; 48: 557-561.
- Franco G, Fonte R, Tempini G, Candura F. Serum bile acid concentrations as a liver function test in workers occupationally exposed to organic solvents. *Int Arch Occup Environ Health* 1986; 58: 157-164.
- Franco G, Santagostino G, Lorena M, Imbriani M. Conjugated serum bile acid concentrations in workers exposed to low doses of toluene and xylene. *Br J Ind Med* 1989; 46: 141-142.

- Gentile S, Persico M, Tiribelli C. Abnormal hepatic uptake of low doses of sulfobromophthalein in Gilbert's syndrome : the role of reduced affinity of the plasma membrane carrier of organic anions. *Hepatology* 1990; 12: 213-7.
- Gibaldi M, Perrier D. Pharmacokinetics. New York: Marcel Dekker, 1982; 60
- Gilli G, Bono R, Scursatone E, Guerrini E. Geographical and temporal patterns of air-borne and personal 1,1,1-trichloroethane exposure in Piedmont region (Italy). *Sci Total Environ* 1992; 116: 261-8.
- Goel SK, Rao GS, Pandya KP, Shanker R. Trichloroethylene toxicity in mice : a biochemical, hematological and pathological assessment. *Indian J Exp Biol* 1992; 30: 402-6.
- Gopinath C, Prentice DE, Street AE, Crook D. Serum bile acid concentration in some experimental liver lesions of rat. *Toxicology* 1980; 15: 113-127.
- Gottlieb ME, Stratton HH, Newell JC, Shah DM. Indocyanine green. Its use as an early indicator of hepatic dysfunction following injury in man. *Arch Surg* 1984; 119: 264-8.
- Hake CL et al. The metabolism of 1,1,1-trichloroethane by the rat. *Arch Environ Health* 1960; 1: 101-5.
- Hamdan H, Stacey NH. Mechanism of trichloroethylene - induced elevation of individual serum bile acids. I. Correlation of trichloroethylene concentrations to bile acids in rat serum. *Toxicol Appl Pharmacol* 1993; 121: 291-295.
- Henry RJ, Donald CC, James WW. Clinical chemistry: Principles and Technics, 2nd ed. Hargertown: Harper & Row, 1974.
- Herren-Freund SL et al. The carcinogenicity of trichloroethylene and its metabolites, trichloroacetic acid and dichloroacetic acid,

- in mouse liver. *Toxicol Appl Pharmacol* 1987; 90: 183-9.
- Hodgson E, Guthrie FE. Hepatotoxicity. Introduction to biochemical toxicology., Rev. *Biochem toxicol* 1987; 8: 341-356.
- Hodgson MJ, Heyl AE, Van Thiel DH. Liver disease associated with exposure to 1,1,1-trichloroethane. *Arch Intern Med* 1989; 149: 1793-1798.
- Hoffmann P, Breitenstein M, Toraason M. Calcium transients in isolated cardiac myocytes are altered by 1,1,1-trichloroethane. *J Mol Cell Cardiol* 1992; 24: 619-29.
- Honma T. Effects of trichloroethylene, 1,1,1-trichloroethane and carbon tetrachloride on plasma lipoproteins of rats. *Ind Health* 1990; 28: 159-74.
- Hougaard K, Ingvar M, Wieloch T, Siesjo BK. Cerebral metabolic and circulatory effects of 1,1,1-trichloroethane, a neurotoxic industrial solvent. I. Effects on local cerebral glucose consumption and blood flow during acute exposure. *Neurochem Pathol* 1984; 2: 39-53.
- Hurtuk BL, Krefetz RG. Enzymes. In: Michael LB, Janet LD, Edward PF, eds. Clinical chemistry : Principles, Procedure, Correlations, 2nd ed., pp 215-247. Philadelphia, J.B. Lippincott, 1992.
- Ikeda M, Ohtsuji H. A comparative study of the excretion of Fujiwara reaction positive substances in the urine of humans and rodents given trichloro- or tetrachloro- derivatives of ethane and ethylene. *Br J Ind Med* 1972; 29: 99-104.
- Jensen AL. Evaluation of fasting and postprandial total serum bile acid concentration in dogs with hepatobiliary disorders. *J Vet Med* 1991; A38: 247-254.
- Johnson SE, Rogers WA, Bonagura JD, Caldwell JH. Determination of serum bile acids in fasting dogs with hepatobiliary disease.

- Am J Vet Res* 1985; 46: 2048-2053.
- Kadry AM, Farghali H, Abdel-Rahman MS. Toxicity and metabolism of trichloroethylene in rat hepatocytes. *Arch Environ Contam Toxicol* 1989; 18: 888-94.
- Kaplowitz N, Kok E, Javitt NB. Postprandial serum bile acid for the detection of hepatobiliary disease. *JAMA* 1979; 225: 292-293.
- Karmen A, Wroblewski F, La Due JS. *J Clin Invest* 1955; 34: 126.
- Kawasaki S, Sugiyama Y, Iga T, Hanano M, Beppu T, Sugiura M, Sanjo K, Idezuki Y. Hepatic clearances of antipyrine, indocyanine green, and galactose in normal subjects and in patients with chronic liver diseases. *Clin Pharmacol Ther* 1988; 44: 217-224.
- Klaassen CD, Plaa GL. Comparison of the biochemical alterations elicited in livers from rats treated with carbon tetrachloride, chloroform, 1,1,2-trichloroethane and 1,1,1-trichloroethane. *Biochem Pharmacol* 1969; 18: 2019-2027.
- Klaassen CD, Plaa GL. Relative effects of various chlorinated hydrocarbons on liver and kidney function in mice. *Toxicol Appl Pharmacol* 1966; 9: 139-151.
- Klaassen CD, Watkins JB III. Mechanism of bile formation, hepatic uptake and biliary excretion. *Pharmacol Rev* 1984; 36: 1-67.
- Kobayashi H, Hobara T, Kawamoto T, Sakai T. Effects of 1,1,1-trichloroethane inhalation on heart rate and its mechanism: a role of autonomic nervous system. *Arch Environ Health* 1987; 42: 140-3.
- Korman MG, Hofmann AF, Summerskill WHJ. Assessment of activity in chronic active liver disease. Serum bile acids compared with conventional tests and histology. *N Engl J Med* 1974; 290: 1399-1402.
- Kostrzewski P, Jakubowski M, Kolacinski Z. Kinetics of

- trichloroethylene elimination from venous blood after acute inhalation poisoning. *J Toxicol Clin Toxicol* 1993; 31: 353-63.
- Kraft W, Lechner J, Vollmar AM, Reusch C, Warmbier M, Lohss E. Indocyanine green test in the dog. *Tierarztl Prax* 1991; 19: 439-446.
- Kroker R, Romer C. The significance of serum bile acid concentrations as indication of hepatic dysfunction in the mini-pig. *Zbl Vet Med* 1984; 31: 287-295.
- Kukongviriyapan V, Kukongviriyapan U, Stacey NH. Interference with hepatocellular substrate uptake by 1,1,1-trichloroethane and tetrachloroethylene. *Toxicol Appl Pharmacol* 1990; 102: 80-90.
- Liss GM, Greenberg RA, Tamburro CH. Use of serum bile acids in the identification of vinyl chloride hepatotoxicity. *Am J Med* 1985; 78: 68-78.
- Lundberg I, Ekdahl M, Kronevi T, Lidums V, Lundberg S. Relative hepatotoxicity of some industrial solvents after intraperitoneal injection or inhalation exposure in rats. *Environ Res* 1986; 40: 411-420.
- Lundberg I, Hakansson M. Normal serum activities of liver enzymes in Swedish paint industry workers with heavy exposure to organic solvents. *Br J Ind Med* 1985; 42: 596-600.
- MacFarland NH. Toxicology of solvents. *Am Ind Hyg Assoc J* 1986; 47: 704-707.
- Mashige F, Imai K, Ozuga T. A simple and sensitive assay of total serum bile acids. *Clinica Chimica Acta* 1976; 70: 79-86.
- Mazzullo M, Bartoli S, Bonora B, Colacci A, Lattanzi G, Niero A, Silingardi P, Grilli S. In vivo and in vitro interaction of trichloroethylene with macromolecules from various organs of rat and mouse. *Res Commun Chem Pathol Pharmacol* 1992; 76:

192-208.

- Mercier. 1,1,1 - Trichloroethane, JARC monographs on the evaluation of the carcinogenic risk of chemicals to human. 1979; 20: 515-530.
- Mori M, Oyamada M, Sakauchi F, Ogawa K. Effects of colchicine on the hepatocellular transport of indocyanine green in the rat. *Virchows Arch B Cell Pathol* 1987; 53: 37-43.
- Nagaya T, Ishikawa N, Hata H, Otobe T. Subclinical and reversible hepatic effects of occupational exposure to trichloroethylene. *Int Arch Occup Environ Health* 1993; 64: 561-3.
- Nakayama H, Kobayashi M, Takahashi M, Ageishi Y, Takano T. Generalized eruption with severe liver dysfunction associated with occupational exposure to trichloroethylene. *Contact Dermatitis* 1988; 19: 48-51.
- Novros JS, Koch TR, Knoblock EC. *Clin Chem* 1979; 25: 1891-1899.
- Okino T et al. Morphological and biochemical analyses of trichloroethylene hepatotoxicity : differences in ethanal- and phenobarbital-pretreated rats. *Toxicol Appl Pharmacol* 1991; 108: 379-89.
- Odum J, Foster JR, Green T. A mechanism for the development of clara cell lesions in the mouse lung after exposure to trichloroethylene. *Chem Biol Interact* 1992; 83: 135-53
- Ozuga T, Mitamura K, Mashige F, Imai K. Evaluation of fluorimetrically estimated serum bile acid in liver disease. *Clinica Chimica Acta* 1977; 75: 81-90.
- Pearlman FC, Lee RTY. *Clin Chem* 1974; 20: 477-453.
- Perbellini L, Olivato D, Zedde A, Miglioranzi R. Acute trichloroethylene poisoning by ingestion : clinical and pharmacokinetic aspects. *Intensive Care Med* 1991; 17: 234-5.

- Pessayre D et al. Hepatotoxicity of trichloroethylene - carbon tetrachloride mixture in rats. A possible consequence of potentiation by trichloroethylene of carbon tetrachloride - induced lipid peroxidation and liver lesions. *Gastroenterology* 1982; 83: 761-72.
- Plaa GL. Toxic responses of the liver. In: Amdur MO, Doull J, Klaassen CD, eds. Cassarett and Doull's toxicology. 4th edition. U.S.A.: McGraw-Hill, 1992.
- Plaa GL, Evans EA, Hine CH. Relative hepatotoxicity of seven halogenated hydrocarbon. *J pharmacol Exp Ther* 1958; 123: 224-229.
- Plaa GL, Hewitt WR. Detection and evaluation of chemically induced liver injury. In: Hayes AW, ed. Principle and Methods of Toxicology. 2nd Edition. New York: Raven Press, 1989; 599-628.
- Plaa GL, Larson RE. Relative nephrotoxic properties of chlorinated methane, ethane and ethylene derivatives in mice. *Toxicol Appl Pharmacol* 1965; 7: 37-44.
- Potter BJ, Blades BF, Shepard MD, Thung SM, Berk PD. The kinetics of sulfobromophthalein uptake by rat liver sinusoidal vesicles. *Biochim Biophys Acta* 1987; 898: 159-71.
- Priestly BE, Plaa GL. Temporal aspects of carbon tetrachloride-induced alteration of sulfobromophthalein excretion and metabolism. *Toxicol Appl Pharmacol* 1970; 17: 786-794.
- Prout MS, Provan WM, Green T. Species differences in response to trichloroethylene. I. Pharmacokinetics in rats and mice. *Toxicol Appl Pharmacol* 1985; 79: 389-400.
- Quast JF, Calhoun LL, Frauson LE. 1,1,1-Trichloroethane formation : a chronic inhalation toxicity and oncogenicity study in Fischer 344 rats and B6C3F1 mice. *Fundam Appl Toxicol* 1988; 11: 611-625.

- Ray K, Sahana CC, Chaudhuri SB, De GC, Chatterjee K. Effects of trichloroethylene anaesthesia on salivary paracetamol elimination. *Indian J Physiol Pharmacol* 1993; 37: 79-81.
- Reichling JJ, Kaplan MM. Clinical use of serum enzymes in liver disease. *Dig Dis Sci* 1988; 33: 1601-1614.
- Richards TG, Tindall VR, Young A. A modification of the bromsulphthalein liver function test to predict the dye content of the liver and bile. *Clin Sci* 1959; 18: 499-511.
- Ricker H, Christensen M, Arnfred T, Dige U, Thaysen EH. The diagnostic value of fasting serum bile acid concentration in patients with suspected liver disease. *Scand J Gastroenterol* 1982; 17: 565-570.
- Savolainen H et al. Trichloroethylene and 1,1,1-trichloroethane : Effects on brain and liver activity after five days intermittent inhalation. *Arch Toxicol* 1977; 38: 229-237.
- Schumann AM, Fox TR, Watanabe PG. A comparison of the fate of inhaled methyl chloroform (1,1,1-trichloroethane) following single or repeated exposure in rats and mice. *Fundam Appl Toxicol* 1982; 2: 27-32.
- Schumann AM, Fox TR, Watanabe PG. 14 C Methyl chloroform (1,1,1-trichloroethane) : Pharmacokinetics in rats and mice following inhalation exposure. *Toxicol Appl Pharmacol* 1982; 62: 390-401.
- Selden A, Hultberg B, Ulander A, Ahlborg GJ. Trichloroethylene exposure in vapour degreasing and the urinary excretion of N-acetyl-beta-D-glucosaminidase. *Arch Toxicol* 1993; 67: 224-6.
- Shibata H, Odani N. Blood clearance of ^{99m}Tc-phytate and indocyanine green in carbon tetrachloride treated dogs. *J Toxicol Sci* 1991; 16: 145-154.

- Sjogern B et al. Pulmonary reactions caused by welding-induced decomposed trichloroethylene. *Chest* 1991; 99: 237-8.
- Skender L, Karacic V, Basner B, Prpic MD. Assessment of exposure to trichloroethylene and tetrachloroethylene in the population of Zagreb, Croatia. *Int Arch Occup Environ Health* 1993; 65 (1 Suppl.): S 163-5.
- Skrede S, Solberg HE, Blomhoff JP, Gjone E. Bile acid measured in serum during fasting as a test for liver disease. *Clin Chem* 1978; 24: 1095-1099.
- Sotaniemi EA, Sutinen S, Sutinen S, Arrants AJ, Pelkonen RO. Liver injury in subjects occupationally exposed to chemicals in low doses. *Acta Med Scand* 1982; 212: 207-215.
- Stacey NH. Toxicity of mixtures of trichloroethylene, tetrachloroethylene and 1,1,1-trichloroethane : similarity of in vitro to in vivo response. *Toxicol Ind Health* 1989; 5:441-50.
- Stremmel W, Berk PD. Hepatocellular uptake of sulfobromophthalein and bilirubin is selectively inhibited by an antibody to the liver plasma membrane. *J Clin Invest* 1986; 78: 822-6.
- Takano T et al. Interaction of 1,1,1-trichloroethane with the mixed function oxidation system in rat liver microsomes. *Xenobiotica* 1988; 18: 1457-64.
- Tamburro CH, Greenberg R. Effectiveness of federally required medical laboratory screening in the detection of chemical liver injury. *Environ Health Persp* 1981; 41: 117-122.
- Thiele DL, Eigenbrodt EH, Ware AJ. Cirrhosis after repeated trichloroethylene and 1,1,1-trichloroethane exposure. *Gastroenterology* 1982; 83: 926-9.
- Van den Bergh AAH, Muller P. *Biochem* 1916; Z 77: 90-103.
- Wang G, Stacey NH. Elevation of individual serum bile acids on

exposure to trichloroethylene or α -naphthylisothiocyanate.

Toxicol Appl Pharmacol 1990; 105: 209-215.

Whitby H. Occupational liver disease. *Occup Health* 1986; 211-212.

Xia L, Yu T. Study of the relationship between the hepatotoxicity and free radical induced by 1,1,2-trichloroethane and 1,1,1-trichloroethane in rat. *Biomed Environ Sci* 1992; 5: 303-13.