

Omega-3 Fatty Acids in Thai Vegetables and Freshwater Algae

Lifeng Yang^{1*}, Onanong Phuseerit², Sirithon Siriamornpun³,
Srinual Jantathai³ and Li Hua⁴

¹ Department of Nutrition, Ningbo College of Health Science, Zhejiang 315000, China

² Department of Science and Technology, Faculty of Liberal Arts and Science, Roi-Et Rajabhat University, Selapoom, Roi-Et 45120, Thailand

³ Department of Food Science and Nutrition, Mahasarakham University, Mahasarakham, 44150, Thailand

⁴ Department of Cuisine and Nutrition Science, Yangzhou University, Jiangsu 225127, China

* Corresponding authors. E-mail addresses: lifeng_yang@126.com (L Yang)
Tel.: +86 15205746025

(Received: 8th May 2020, Revised: 9th June 2020, Accepted: 22nd June 2020)

Abstract - The aim of this study was to analyze polyunsaturated fatty acid (PUFA) composition in seventeen Thai vegetables, including six species of aquicolous, two freshwater algae, and nine terricolous species. For each sample, palmitic acid, linoleic acid and alpha-linolenic acid (18:3n-3) were the main fatty acids, which constituted nearly 90% of the total fatty acid composition of leaf tissues. The concentration ranges were 5.9 to 310 mg/100g (palmitic acid), 3 to 235 mg/100g (linoleic acid) and 2 to 227 mg/100g (alpha-linolenic acid), respectively. Green algae contained high level of C18 PUFA and low level of C20 PUFA. The algae and terricolous samples had ratios of n-6/n-3 ranged from 0.6 in Tao num to 2.0 in Phai, while aquicolous vegetable samples had a greater variation being ranged from 0.3 in Krached to 6.3 in Koon. Our findings on the fatty acid profile of vegetable showed n-3 fatty acid and ratio of n-6/n-3 might be of great nutritional interest.

Keywords: Thai vegetables, aquicolous, algae, terricolous, n-3 fatty acids

1. Introduction

Dietary and lifestyle patterns as well as energy balance have been reported to influence chronic non-communication diseases such as obesity (Kosti et al., 2008), diabetes mellitus, hypertension, stroke, cardiovascular disease (CVD) and some type of cancer, through a multitude of risk factors including lipids (Katcher et al., 2009), glucose tolerance, insulin tolerance (Oldroyd et al., 2006), eicosanoids (Kinsella, Broughton and Whelan, 1990), blood pressure (Sugiyama et al., 2007), coagulation and inflammation (Demosthenes et al., 2004). Current evidences support n-3 fatty acids supplementation, have multiple beneficial effects on plasma lipids (Laidlaw and Holub, 2003), blood pressure (Sirtori et al., 1998), cardiac function (Bao et al., 1998), arterial and venous thrombosis (Andriamampandry et al., 1999), endothelial functions (Engler et al., 2004) as well as potent anti-inflammatory qualities (Gil, 2002). Diets high in PUFA have a substantial effect on cholesterol and on decreases in coronary heart disease (CHD) mortality (Thomas et al., 1997; Okuda et al., 2005). However, total fat in the diet should not exceed 30% as recommended by WHO and FAO (WHO/FAO, 2002) and intake of dietary cholesterol should be kept below 300 g/day (WHO, 2003).

Many enzymes were considered to play a major role during de novo synthesis and subsequent modification of plant fatty acids. Such as chloroplast acetyl-CoA synthetase, Acetyl-CoA carboxylase (ACCase) which is the first committed step for fatty acid and acyl lipid synthesis, presumably inhibited by aryloxyphenoxypropionates (AOPPs) and the cyclohexanediones (CHDs). Sukenik and Livne (1991) found

the marine alga, *Isochrysis galbana* the in vitro activity of ACCase correlated well with changes in synthesis in response to light intensity, and desaturation enzymes such as $\Delta 9$, $\Delta 12$, $\Delta 15$ -desaturase were responsible for the major fatty acids of plants are the C18 compounds, oleic, linoleic and linolenic acids, which desaturated either by chloroplast (plastid) or by endoplasmic reticulum reviewed by Harwood (1996). Pan et al. (2007) revealed that heterologous expression of the cDNAs have ability to biosynthesize different fatty acid, recombinant SbDES2 converted palmitoleic acid (16:1 $\Delta 9$) to hexadecadienoic acid (16:2 $\Delta 9,12$), and that recombinant SbDES3 was capable of converting hexadecadienoic acid into hexadecatrienoic acid (16:3 $\Delta 9,12,15$) in *Saccharomyces cerevisiae*.

Environmental characteristics (temperature, mentals, and nature products) effects on plant lipids have been reviewed by previous studies, that the level of unsaturation of membrane fatty acids was inversely correlated with growth temperature. Williams and coworkers suggested that low temperature-induced desaturation C16 and C18 fatty acid in both cytosolic and the chloroplast resulted in higher levels of unsaturated fatty acids in the brassica napus leaf (Williams, Williams and Khan, 1992). The results from Falcone, Ogas and Somerville (2004) revealed key changes in the membrane fatty acid composition were decreasing in trienoic fatty acids and consistent increasing in unsaturated 16:0 and in dienoic 18:2 levels that occur in response to elevated temperature in Arabidopsis. It had been reported that metal supply such as cadmium resulted in a change of lipid content but did not affect fatty acids composition (Nouairi et al, 2006). Kabala et al. (2008) found

that exposure of plants to heavy metals such as Cd, Cu and Ni did not enhance the lipid peroxidation in the plasma membrane (PM) fraction. However, all metals caused an increase in the saturation of PM fatty acids and a decrease in the length of the fatty acid chain.

Evidence supported by Gao et al (2005) that uptake of organic chemicals such as phenanthrene and pyrened from soil and water exhibit significantly positive correlation with lipid contents (Gao, Zhu and Ling, 2005). Liang et al (2006) cultured the marine diatoms *Phaeodactylum tricorutum* and *Chaetoceros muelleri*, the result revealed that the maximum value of SFA was found in the ammonium treatment, that of MUFA in the nitrate treatment and for PUFA in the urea treatment irrespective of the UV radiation, respectively. Some nature products were associated with lipids and fatty acids synthesis. The results from Wang et al had indicate that eigalocatechin gallate (EGCG) the main component of tea polyphenols which is the specific inhibitor of fatty acid synthase from chicken liver, probably acting on β -ketoacyl reductase. The free carnitine occurred in the plant tissues of several species, such as cereals, legumes, and germinating seeds, which suggested to involve in diverse fatty acid metabolic pathways in Arabidopsis seedlings (Bourdin, Adenier and Perrin, 2007).

Good sources of n-3 fatty acids were fish, seafood and plant oils. Previous studies have been reported that edible vegetables such as mint (Pereira and Sinclair, 2001), purslane (Liu et al, 2000) and vervain (Guil, Torija, Gimenez and Rodriguez, 1996) content high percentage of n-3 fatty acids. Wild plants including

vegetables and seeds consumption have long been considered to have health benefits mainly due to vitamins, minerals and phytonutrients, various studies had been done in Thailand to assess the nutrients. Teaw had been proved having strong antioxidant properties (Maisuthisakul Pongsawatmanit and Gordon, 2007); Pak-paw, which was high in calcium (Kamchan et al, 2004), and guava, which showed the high anti-proliferative activity in vitro (Manosroi, Dhumtanom and Manosroi, 2006). This study reported for the first time, data on fatty acid composition and concentration of vegetables in Thailand, and provided an incentive to seek for an alternative nutritional source of PUFA.

2. Materials and methods

2.1 Materials

In this study, we collected seventeen species of vegetables purchased from local market Mahasarakham Thailand. Including six species of aquatic vegetables: Krached (*Neptunia oleracea* Lour), Bung Thai (*Ipomoea aquatica* Forsk), Chi nam (*Oenanthe stolonifera*), Koon (*Colocasia gigantea* Hook.f.), Bon (*Colocasia esculenta* Sahott) and Buo Sai (*Nymphaea lotus* L); two species of algae: Pum (*Wolffia arrhiza* L. Wimm) and Tao num (*Spirogyra* sp); nine species of terricolous vegetables: Khayang (*Limnophila aromatica* (Lak.) Merr), Kratin (*Leucaena leucocephala* de Wit), Kee-lek (*Cassia siamea* Lamk), Neamhuseu (*Coleus amboinicus* Lour), Kwinin (*Azadirachta indica* A. luss), Krad (*Spilanthes acmella* Murr), Phai (*Polygonum odoratum* Lour). All samples were conducted to analyze in more than triplicates ($n \geq 3$). For all the samples in this study, only the chloroplast-rich

leaves or heads were analyzed, and stems were omitted from analysis.

2.2 Methods

Approximately 10g of fresh vegetables were cut to pieces, and homogenized in 50 volumes of methanol-chloroform (2:1, v/v) containing 10 mg/L butylated hydroxytoluene and 0.1 mg/ml of tricosanoic acid (C23:0, Fluka, Germany) as an internal standard, extracted for overnight. Then each sample was filtered; the residue was extraction by 10 ml chloroform and methanol twice. The extracts were pooled and concentrated, and transferred into a 10ml methylation tube by chloroform containing 10 mg/L BHT.

Preparation of ester derivatives of fatty acids for chromatographic analysis: The fatty acid methyl esters (FAMES) of the total lipid extract were prepared by transesterification with 3 ml H₂SO₄ in methanol (0.9 mol/L) plus 1 ml toluene and incubating for 2 hours at 70°C. Before injected to GC, FAMES were filtered by Sep-pak silica column (Altech, Associates Inc., USA). Samples for gas chromatographic analysis was followed as we prior reported (Yang et al, 2006), the GC temperature program of analysis, was initial 150°C held for 1min, raised to 180°C at the ratio 25°C /min, raised to 220°C at the ratio of 2.5°C /min held for 4min, raised to 230°C at 15°C /min and then held this temperature for 3min.

3. Results and discussion

The fatty acids composition and concentration of total lipids determined by capillary gas chromatography were shown in (Table 1-4). There were 24 fatty acids identified in the samples examined. The

major fatty acids were palmitic acid (16:0), Oleic acid (18:1), Linoleic acid (18:2n-6) and linolenic acid (18:3n-3), which constituted nearly 90% of the total fatty acid composition of leaf tissues.

3.1 N-3 polyunsaturated fatty acids composition and concentration in commonly vegetables

Among the aquicolous vegetable and algae samples, we determined three n-3 fatty acids, which were hexadecatrienoic acid (16:3n-3), 18:3n-3 and eicosapentaenoic acid (20:5n-3). The most abundant fatty acid in these species was 18:3n-3, with the composition ranged from 5% of total fatty acids in Koon to 52.3% of total fatty acids in Krached. The 16:3n-3 and 20:5n-3 were present in small amounts in the aquicolous vegetables and algae samples, which were found in Buo Sai (2.9% and 8.5% of total fatty acids, respectively) and Pum (1.6% and 6.8% of total fatty acids, respectively); Pum was found mostly in tropical and subtropical water environments, exhibited the ability of absorbs large amounts of nitrogen and phosphorus (Fujita et al, 1999) which were associated with changes PUFA composition (Lin et al, 2007). There were no 16:3n-3 and 20:5n-3 presented in Krached, Chi nam, Koon and Bon, additionally. In (Table 1 and 2) showed that green algae had the highest level of C18 PUFAs and lowest in C20 PUFAs, the same results were found in algae from the Bohai Sea (Li et al, 2002). There were only two n-3 fatty acids identified in terricolous vegetable samples, they were 16:3n-3 and 18:3n-3. 18:3n-3 was the predominated n-3 fatty acid, detected in all terricolous samples, with the composition ranged from 19.5% in Wan to 35.8% in Krad, the concentration of 18:3n-3

ranged from 36.8 mg/100g in Neamhuseu to 282.1 mg/100g in Kratin. There was no 16:3n-3 identified in Neamhuseu, Wan and Tew Khoaw. Krad and Phai had the highest proportion of n-3 fatty acid, 35.8%

and 35.3%, respectively, while Kwinin and Kratin had the most concentration, 323.7 mg/100g and 289.1 mg/100g, respectively. In our study, we found Thai vegetable was a novel source of n-3 fatty acid.

Table 1. Fatty acid composition (% of total fatty acid) in Thai aquicolous vegetables and freshwater algae.

Fatty acids	Aquicolous					Algae		
	Krached	Buo Sai	Bung Thai	Chi nam	Koon	Bon	Pum	Tao num
12:0	ND	ND	2.3±0.1	2.1±0.5	ND	ND	ND	ND
14:0	ND	5.3±0.4	3.2±0.2	1.1±0.3	12.6±0.3	ND	2.1±0.2	12.0±0.9
16:0	12.7±0.2	23.1±0.5	25.5±1.5	26.8±2.9	29.3±0.2	36.2±1.4	16.7±1.4	28.1±1.2
17:0	4.6±0.5	2.8±0.6	3.1±0.1	3.3±0.4	2.0±0.3	ND	3.8±0.8	5.6±0.2
18:0	2.0±0.1	2.0±0.2	2.8±0.2	1.8±0.3	2.5±0.2	3.2±0.1	1.4±0.3	8.2±0.9
20:0	10.1±1.0	3.6±0.4	2.5±0.1	1.1±0.2	4.6±0.7	ND	3.2±0.4	ND
SFA	29.4±1.1	36.8±0.7	39.5±1.9	36.2±1.6	50.9±1.3	39.3±1.3	27.2±0.6	53.9±2.0
16:1	1.1±0.1	2.3±0.2	0.5±0.0	1.2±0.1	1.9±0.4	ND	2.5±0.2	2.7±0.3
18:1	1.9±0.2	3.0±0.5	1.1±0.3	4.9±0.4	10.4±2.2	7.7±0.8	3.5±0.3	3.2±0.4
MUFA	3.0±0.1	5.3±0.7	1.6±0.2	6.1±0.4	12.3±1.9	7.7±0.8	6.0±0.5	5.9±0.7
16:3n-3	ND	2.9±0.6	ND	ND	ND	ND	1.6±0.4	ND
18:3n-3	52.3±0.8	7.5±0.3	32.9±0.8	20.7±1.4	5.0±0.4	8.0±0.6	35.9±1.2	25.9±3.1
20:5n-3	ND	8.5±1.4	2.6±0.3	ND	ND	ND	6.8±1.1	ND
Total n-3	52.3±0.8	18.9±1.7	33.5±1.2	20.7±1.4	5.0±0.4	8.0±0.6	44.2±1.4	25.9±3.1
18:2n-6	14.3±0.4	35.4±0.9	21.2±0.7	37.0±0.3	29.2±3.0	45.0±1.0	22.5±1.0	12.6±0.5
18:3n-6	1.0±0.2	ND	ND	ND	ND	ND	ND	1.7±0.4
20:4n-6	ND	ND	2.2±0.1	ND	2.6±0.5	ND	ND	ND
Total n-6	15.3±0.4	35.4±0.9	23.4±0.8	37.0±0.3	31.8±2.3	45.0±1.0	22.5±1.0	14.3±0.6
PUFA	67.7±1.1	54.3±1.2	58.9±1.8	57.7±1.6	36.8±3.0	53.0±0.4	66.8±1.0	40.2±2.4
n-6/n-3	0.3±0.0	4.2±0.4	0.8±0.0	1.8±0.1	6.3±0.3	5.7±0.5	0.8±0.0	0.6±0.1

ND = not detected, SFA = saturated fatty acid, MUFA = monounsaturated fatty acid, PUFA = polyunsaturated fatty acid

Table 2. Fatty acid concentration (mg/100g) in Thai aquicolous vegetables and algae

Fatty acids	Aquicolous					Algae		
	Krached	Buo Sai	Bung Thai	Chi nam	Koon	Bon	Pum	Tao num
12:0	ND	ND	5.6±0.1	2.1±0.5	ND	ND	ND	ND
14:0	ND	3.1±0.1	7.8±0.6	1.1±0.3	5.2±0.7	ND	9.9±1.3	2.5±0.2
16:0	42.7±1.6	13.6±0.7	61.2±1.8	26.2±2.3	12.1±1.2	11.6±0.6	76.5±6.3	5.9±0.2
17:0	15.7±2.5	1.7±0.4	7.5±0.1	3.2±0.4	0.8±0.2	ND	17.5±3.7	1.2±0.0
18:0	6.6±0.1	1.2±0.2	6.7±0.2	1.8±0.3	1.0±0.2	1.0±0.1	6.3±1.0	1.7±0.2
20:0	34.0±3.7	2.1±0.4	6.1±0.1	1.1±0.2	1.9±0.4	ND	14.6±2.3	ND
SFA	99.0±6.7	21.7±1.5	94.8±1.9	35.5±0.7	21.2±2.8	12.6±0.8	124.8±5.5	11.4±0.3
16:1	3.7±0.3	1.4±0.2	1.2±0.0	1.2±0.1	0.8±0.1	ND	11.4±0.8	0.6±0.1
18:1	6.4±0.9	1.8±0.4	2.8±0.7	4.8±0.4	4.4±1.4	2.5±0.4	15.9±0.8	0.7±0.1
MUFA	10.0±0.9	3.1±0.6	4.0±0.7	5.9±0.4	5.1±1.4	2.5±0.4	27.3±1.6	1.2±0.1
16:3n-3	ND	1.7±0.4	ND	ND	ND	ND	7.5±1.8	ND
18:3n-3	176.4±8.0	4.4±0.1	79.1±4.6	20.3±1.8	2.1±0.1	2.6±0.1	164.7±9.9	5.5±0.7
20:5n-3	ND	5.0±0.9	6.3±0.9	ND	ND	ND	31.3±5.7	ND
Total n-3	176.4±8.0	11.1±1.1	85.4±4.9	20.3±1.8	2.1±0.1	2.6±0.1	203.5±11.4	5.5±0.7
18:2n-6	48.3±2.0	20.8±0.9	51.2±3.5	36.3±1.1	12.0±0.1	14.5±1.4	103.2±4.0	2.7±0.1
18:3n-6	3.3±0.8	ND	ND	ND	ND	ND	ND	0.4±0.1
20:4n-6	ND	ND	5.3±0.4	ND	1.1±0.3	ND	ND	ND
Total n-6	51.6±2.3	22.9±1.1	56.5±3.9	36.3±1.1	13.1±0.6	14.5±1.4	103.2±4.0	3.1±0.3
PUFA	227.9±10.7	36.9±2.0	147.6±9.5	56.6±2.9	15.2±0.3	17.1±1.4	306.6±13.7	8.5±0.6

ND = not detected, SFA = saturated fatty acid, MUFA = monounsaturated fatty acid, PUFA = polyunsaturated fatty acid

3.2 N-6 polyunsaturated fatty acids composition and concentration in commonly vegetables

The polyunsaturated fatty acids, linoleic acid and α -linolenic acid, are essential to the human diet because they cannot be synthesized by humans (Innis, 1996; Sinclair, 1990). Researches in human nutrition and health have shown the dietary importance of these essential fatty acids and their relationship to human health (Horrobin, 1993; Innis, 1996). Among the

polyunsaturated fatty acid in aquicolous vegetables and algae samples, the 18:2n-6 was the predominant n-6 fatty acid presented in these species. Bon was rich in 18:2n-6, which made up 45% of total fatty acids, in the other hand, Tao num was low in 18:2n-6, containing 12.6% of total fatty acids. Linoleic acid synthesized from oleic acid through $\Delta 12$ desaturation which is an integral membrane-bound protein in a wide range of plants and microorganisms (Zhang et al, 2007). The 18:3n-6 and arachidonic

acid (20:4n-6) were low presented in terricolous samples and 18:3n-6 was only detected in Krached and Tao num with concentration of 3.3 mg/100g and 0.4mg/100g for accounted 1% and 1.7% of total fatty acid, respectively. The 20:4n-6 was only identified in Koon and Bung Thai with the concentration at 1.1 mg/100g and 5.3 mg/100g, accounted for 2.6% and 2.2%. The total n-6 fatty acid composition and concentration varied from 14.3% in Tao num to 45% in Bon, and 3.1 mg/100g in Tao num to 103.2 mg/100g in Pum. Differ from aquicolous vegetables and algae samples, four n-6 fatty acids were present in terricolous samples, 18:2n-6, 20:4n-6, docosadienoic acid (22:2n-6) and docosatetraenoic acid (22:4n-6). Kee-lek had the highest percentage composition (35.2% of total fatty acid) and concentration (261.6 mg/100g) of 18:2n-6 which was the major n-6 fatty acid found in all terricolous samples we collected. Wan had the lowest composition of 18:2n-6 accounted for 13%, while Neamhuseu had the lowest concentration of 18:2n-6 accounted for 27.4 mg/100g. Kwinin was the only vegetable which was detected with 20:2n-6 and 22:2n-6, with percentage composition of 2.9% and 5.8%, accounted for 36.3 mg/100g and 71.9 mg/100g, respectively. 20:4n-6 and 22:4n-6 were only both identified in two species of terricolous vegetables, and Kee-lek, with small percent of composition, 10.3 mg/100g and 11.2 mg/100g in Kratin; 1.7% 12.7 mg/100g and 8.3 mg/100g in Kee-lek, respectively.

3.3 Total PUFA composition and concentration in commonly vegetables

The total PUFA percentage of composition ranged from 36.8% in Koon (aquicolous species) which represented the low concentration of 15.2 mg/100g to 67.7% in Krached (aquicolous species) which represented the concentration of 227.9 mg/100g; total PUFA concentration ranged from 8.5 mg/100g in Tao num (aquicolous species) which accounted for 40.2% of total fatty acids to 612.6 mg/100g in Kwinin (terricolous species) which accounted for 49.5% of total fatty acids. The metabolic pathways of PUFA biosynthesis as reviewed by Sprecher (2000), recent progress in the cloning and expression of $\Delta 6$ and $\Delta 5$ desaturases has conclusively established that position-specific desaturases are required to synthesize 20-carbon acids with their first double bond at position 5. Animal and human dietary intervention and epidemiological studies both in vitro and in vivo suggested balance of intake n-6 and n-3 fatty acid associated with chronic non-communication disease (Gazani et al, 2001; Tsipas and Morphake, 2003; Clouet et al, 1995; Hogberg et al, 2003). The ratio of n-6 and n-3 essential fatty acids was evolved on the human beings diet. A lower ratio of n-6/n-3 fatty acids was more desirable in reducing the risk of many of the chronic diseases of high prevalence in Western societies (Simopoulos, 2002). The ratios of n-6/n-3 in aquicolous vegetable samples varied from 0.3 in Krached to 6.3 in Koon; while smaller variations were found in algae and terricolous samples ranged from 0.6 in Tao num to 2.0 in Phai.

Table 3. Fatty Acid Composition (% of total fatty acid) in Thai Terricolous Vegetables

Fatty acids	Khayang	Kratin	Kee-lek	Neamhuseu	Kwinin	Krad	Phai	Wan	Tew Khoaw
12:0	ND	2.4±0.1	ND	5.5±0.3	1.5±0.0	ND	12.4±0.2	1.3±0.3	ND
14:0	18.9±5.8	ND	1.1±0.1	ND	2.4±0.2	ND	ND	ND	9.2±0.9
15:0	1.5±0.2	0.9±0.1	ND	ND	0.8±0.1	1.8±0.5	1.1±0.0	0.8±0.2	ND
16:0	21.1±2.4	21.7±0.9	23.1±0.2	37.2±2.6	25.0±0.2	27.3±0.4	18.1±0.1	19.0±0.6	18.3±0.4
18:0	1.7±0.3	3.6±0.3	2.4±0.0	3.0±0.2	3.2±0.1	1.3±0.0	1.9±0.2	3.1±0.1	3.3±0.0
20:0	ND	ND	ND	ND	ND	ND	0.6±0.0	1.9±0.6	ND
22:0	1.1±0.1	ND	ND	ND	0.8±0.1	ND	0.8±0.1	2.3±0.1	ND
SFA	44.3±3.0	26.2±0.8	26.6±0.1	45.7±2.7	33.7±0.4	30.4±0.8	34.9±0.3	27.2±0.2	30.8±0.5
14:1	4.3±0.8	ND	ND	ND	ND	ND	ND	ND	9.9±0.7
15:1	ND	ND	ND	ND	4.3±0.3	ND	ND	ND	ND
16:1	2.2±0.1	0.9±0.2	1.1±0.0	ND	2.9±0.1	1.8±0.1	1.5±0.1	0.8±0.1	ND
17:1	0.9±0.1	ND	ND	ND	ND	ND	ND	ND	ND
18:1	3.4±1.2	2.8±0.1	2.6±0.1	10.6±0.7	6.0±0.6	1.8±0.1	6.6±0.1	30.8±0.9	5.4±0.1
20:1	2.5±0.5	3.0±0.0	3.1±0.0	ND	0.8±0.1	1.4±0.2	0.6±0.1	1.4±0.2	ND
24:1	ND	ND	ND	ND	2.9±0.4	ND	ND	ND	ND
MUFA	13.3±1.4	6.7±0.2	6.8±0.1	10.6±0.7	17.0±0.4	5.0±0.3	8.7±0.0	33.0±0.6	15.2±0.8
16:3n-3	1.1±0.0	0.8±0.2	0.6±0.0	ND	6.4±0.4	2.0±0.3	2.1±0.1	ND	ND
18:3n-3	23.1±1.4	33.4±2.2	28.0±0.1	25.1±3.0	19.7±0.5	35.8±0.4	35.3±0.6	19.5±0.7	30.0±1.0
Total n-3	24.2±1.5	34.2±2.6	28.6±0.1	25.1±3.0	26.1±1.1	37.8±0.6	37.4±0.8	19.5±0.7	30.0±1.0
18:2n-6	18.2±2.2	28.0±2.0	35.2±0.0	18.6±0.6	14.6±0.1	26.7±0.9	19.1±0.1	13.0±1.0	23.9±0.3
20:2n-6	ND	ND	ND	ND	2.9±0.3	ND	ND	ND	ND
20:4n-6	ND	1.2±0.2	1.7±0.1	ND	ND	ND	ND	ND	ND
22:2n-6	ND	ND	ND	ND	5.8±1.2	ND	ND	ND	ND
22:4n-6	ND	1.3±0.1	1.1±0.1	ND	ND	ND	ND	6.1±0.1	ND
Total n-6	18.2±2.2	30.5±2.2	38.0±0.1	18.6±0.6	23.3±1.5	26.7±0.9	19.1±0.1	19.1±1.1	23.9±0.3
PUFA	42.4±2.3	64.7±0.9	66.6±0.1	43.7±3.5	49.5±1.2	64.5±0.9	56.5±0.7	38.5±0.4	53.9±1.3
n-6/n-3	1.3±0.2	1.1±0.1	0.8±0.0	1.3±0.1	1.1±0.1	1.4±0.1	2.0±0.0	1.5±0.2	1.3±0.0

ND = not detected, SFA = saturated fatty acid, MUFA = monounsaturated fatty acid, PUFA = polyunsaturated fatty acid

Table 4. Fatty Acid Concentration (mg/100g) in Thai Terricolous Vegetables

Fatty acids	Khayang	Kratin	Kee-lek	Neamhuseu	Kwinin	Krad	Phai	Wan	Tew Khoaw
12:0	ND	20.0±0.9	ND	8.1±0.9	18.9±0.6	ND	79.9±3.0	5.6±1.1	ND
14:0	42.5±23.8	ND	8.3±0.3	ND	29.5±2.4	ND	ND	ND	48.4±5.2
15:0	3.2±0.5	8.0±1.1	ND	ND	9.4±1.2	5.5±1.9	7.0±0.4	3.3±0.6	ND
16:0	44.5±6.6	182.2±9.3	171.9±10.9	55.0±8.7	310.7±5.5	83.6±5.5	116.9±6.2	80.2±5.5	95.7±5.3
18:0	3.7±0.4	30.2±4.1	18.1±1.2	4.4±0.7	39.6±1.6	4.1±0.3	12.1±0.9	13.0±1.0	17.2±0.8
20:0	ND	ND	ND	ND	ND	ND	3.7±0.2	8.0±2.5	ND
22:0	2.4±0.5	ND	ND	ND	9.5±1.6	ND	5.4±0.5	9.8±0.4	ND
SFA	96.3±31.3	240.5±14.2	198.3±12.0	67.6±10.3	417.5±9.2	93.1±7.6	225.0±8.9	119.8±2.6	161.3±8.1
14:1	9.4±3.6	ND	ND	ND	ND	ND	ND	ND	51.6±3.3
15:1	ND	ND	ND	ND	53.5±4.2	ND	ND	ND	ND
16:1	4.8±1.3	7.7±2.4	8.4±0.3	ND	36.6±1.5	5.6±0.1	9.7±0.8	3.3±0.5	ND
17:1	1.9±0.2	ND	ND	ND	ND	ND	ND	ND	ND
C18:1	6.9±1.0	23.5±1.3	19.5±0.7	15.7±2.4	75.1±7.7	5.5±0.7	42.3±2.0	129.7±8.2	28.1±1.7
20:1	5.1±0.3	25.2±2.4	22.8±1.1	ND	9.7±1.4	4.2±1.0	3.9±0.2	5.7±0.8	ND
24:1	ND	ND	ND	ND	36.6±5.0	ND	ND	ND	ND
MUFA	28.0±4.3	56.4±5.7	50.6±2.0	15.7±2.4	211.5±6.7	15.4±1.7	55.8±2.5	138.7±7.2	79.7±4.4
16:3n-3	2.5±0.7	7.0±1.8	4.3±0.6	ND	79.2±5.6	6.3±1.0	13.4±0.5	ND	ND
18:3n-3	49.9±15.2	282.1±42.2	208.1±11.3	36.8±2.3	244.5±7.4	109.7±6.7	227.7±14.1	81.8±5.2	156.9±10.7
Total n-3	52.4±16.3	289.1±44.5	212.4±12.5	36.8±2.3	323.7±10.5	115.9±9.2	231.1±16.2	81.8±5.2	156.9±10.7
18:2n-6	38.3±5.3	235.3±18.6	261.6±14.7	27.4±1.8	180.8±0.5	81.5±2.0	123.2±6.3	54.4±3.2	125.3±7.3
20:2n-6	ND	ND	ND	ND	36.3±3.0	ND	ND	ND	ND
20:4n-6	ND	10.3±0.9	12.7±0.8	ND	ND	ND	ND	ND	ND
22:2n-6	ND	ND	ND	ND	71.9±14.2	ND	ND	ND	ND
22:4n-6	ND	11.2±2.1	8.3±0.8	ND	ND	ND	ND	25.7±1.1	ND
Total n-6	ND	256.8±23.4	282.6±17.3	27.4±1.8	289.1±16.1	81.5±2.0	123.2±6.3	80.1±4.7	125.3±7.3
PUFA	80.7±21.2	546.0±56.2	495.1±27.9	64.1±2.9	612.6±5.0	197.5±8.9	364.2±20.7	161.9±5.6	282.2±17.7

ND = not detected, SFA = saturated fatty acid, MUFA = monounsaturated fatty acid, PUFA = polyunsaturated fatty acid

3.4 MUFA composition and concentration in commonly vegetables

Results from epidemiological and dietary intervention studies have shown that consumption of MUFA has beneficial effects

on reducing LDL cholesterol and fasting glucose, increasing immune response, which means having effects on reducing the risk of diabetes. In aquicolous vegetable and algae samples, there were two monounsaturated fatty acids (MUFA) found, palmitoleic acid (16:1) and Oleic acid (18:1), and there was

no 16:1 identified in Bon. The predominant MUFA was 18:1, with the composition ranged from 1.1% in Bung Jeen to 10.4% in Koon. The amount of total MUFA was appreciably low, ranged from 1.6% in Bung Jeen to 12.3% in Koon of total fatty acids; the concentration of 18:1 and total MUFA ranged from 0.7 - 1.2 mg/100g in Tao num, 15.9 - 27.3 mg/100g in Pum, respectively. In terricolous vegetable samples, oleic acid was detected in all the samples from 1.8%-30.8%, with concentration accounted for 5.5 -129.7 mg/100g, and total MUFA ranged from 5%-33% accounted for 15.4 -137.8 mg/100g in Krad and Wan, respectively. There were another six MUFA found in some species, myristoleic acid (14:1), 16:1, 17:1, eicosenoic acid (20:1) and nervonic acid (24:1). 14:1 was detected in Khayang (4.3%) and Tew Khoaw (9.9%). 15:1 and 24:1 were only examined in Kwinin with the ratio of total fatty acids composition of 4.3% and 2.9%. 16:1 and 20:1 were presented together in six species vegetables, Khayang (2.2% and 2.5%), Kratin (0.9% and 3%), Kee-lek (1.1% and 3.1%), Kwinin (2.9% and 0.8%), Krad (1.8% and 1.4%), Phai (1.5% and 0.6%) and Wan (0.8% and 1.4%).

3.5 Saturated fatty acids composition and concentration in commonly vegetables

A recent study by Kelly et al. (2001) reported that diet enriched in palmitic acid (16:0) resulted in an increased ex vivo collagen and ADP induced whole blood platelet aggregation (Kelly, 2001), and study from Li et al. (2003) reported that diet rich in SFA may influence TXA2 formation via effect on tissue proportion of 20:4n-6 (Li et al, 2003). Palmitic acid

was the most abundant saturated fatty acid in all species, ranging from 13% of total lipid in Krached to 37% in Neamhuseu, and presented as the majority fatty acid in four species of samples, including Koom (29%), Tao num (28%), Kwinin (25%) and Neamhuseu (37%). There were six saturated fatty acid presented in aquicolous vegetables, containing 12:0, myristic acid (14:0), 16:0, heptadecanoic acid (17:0), stearic acid (18:0) and arachidic acid (20:0), while seven saturated fatty acid identified in terricolous vegetables, including 12:0, 14:0, Pentadecanoic acid (15:0), 16:0, 18:0, 20:0 and docosanoic acid (22:0). The total SFA ranged 26% in Kratin 54% in Pum and Tao num. The concentration of 16:0 and total SFA in aquicolous species were lower than 100 mg/100g, ranged from 12 mg/100g in Bon to 61 mg/100g in Bung Jeen, and 12.6 mg/100g in Bon to 99 mg/100g in Krached, respectively. There were four species of samples in terricolous vegetables had the concentration of 16:0 more than 100 mg/100g, Kratin (182 mg/100g), Kee-lek (172 mg/100g), Kwinin (311 mg/100g), and Phai (170 mg/100g); six species of samples from both algae and terricolous vegetables had the concentration of total SFA more than 100 mg/100g, Pum (125 mg/100g), Kratin (241 mg/100g), Kee-lek (198 mg/100g), Kwinin (417 mg/100g), Phai (225 mg/100g), Wan (120 mg/100g) and Tew Khoaw (161 mg/100g).

4. Conclusions

We have demonstrated that vegetables contained a good profile of fatty acids. Unlike some nuts or legume seeds, the vegetable oils contain low content of saturates and high amount of PUFAs. In addition, some vegetable oils analyzed in

this study showed a substantial proportion of n-3 fatty acid especially 18:3 n-3. The desirable ratios of n-6/n-3 (0.3-6.3) might be of great nutritional interest. Apart from fiber and other nutrients, vegetable should be recommended as a good source for fatty acids.

5. Acknowledgement

The authors gratefully thank Mahasarakham University for the laboratory facilities.

6. Reference

- Andriamampandry, M.D., Leray, C., Freund, M., Cazenave, J.P. and Gachet, C. 1999. Antithrombotic effects of (n-3) polyunsaturated fatty acids in rat models of arterial and venous thrombosis. *Thrombosis Research*, 93, 9-16.
- Bao, D.Q., Mori, T.A., Burke, V., Puddey, I.B. and Beilin, L.J. 1998. Effects of dietary fish and weight reduction on ambulatory blood pressure in overweight hypertensives. *Hypertension*, 32, 710-717.
- Bourdin, B., Adenier, H. and Perrin, Y. 2007. Carnitine is associated with fatty acid metabolism in plants. *Plant Physiology and Biochemistry*, 45, 926-931.
- Clandinin, M.T., Foxwell, A., Goh, Y.K., Layne K. and Jumpson, J. A. 1997. Omega-3 fatty acid intake results in a relationship between the fatty acid composition of LDL cholesterol ester and LDL cholesterol content in humans. *Biochimica et Biophysica Acta (BBA)-Lipids and Lipid Metabolism*, 1346, 247-252.
- Clouet, P., Niot, I., Gresti, J., Demarquoy, J., Boichot, J., Durand, G. and Bezard, J. 1995. Polyunsaturated n-3 and n-6 fatty acid at a low level in the diet alter mitochondrial outer membrane. *Journal of Nutrition Biochemistry*, 6, 626-634.
- Engler, M.M., Engler, M.B., Malloy, M., et al. 2004. Docosahexaenoic acid restores endothelial function in children with hyperlipidemia: results from the EARLY study. *International Journal of Clinical Pharmacology and Therapeutics*, 42, 672-679.
- Falcone, D.L., Ogas, J.P. and Somerville, C.R. 2004. Regulation of membrane fatty acid composition by temperature in mutants of Arabidopsis with alterations in membrane lipid composition. *BMC Plant Biology*, 4(17), 1-15.
- Fujita, M., Mori, K. and Kodera, T. 1999. Nutrient removal and starch production through cultivation of *Wolffia arrhiza*. *Journal of bioscience and bioengineering*, 87, 194-198.
- Gao, Y.Z., Zhu, L.Z. and Ling, W.T. 2005. Application of the partition-limited model for plant uptake of organic chemicals from soil and water. *Science of The Total Environment*, 336, 171-182.
- Gazani, M.R., Smith, L., Haggarty, P., Fowler, P.A. and Templeton, A. 2001. High n-3: n-6 fatty acid ratios in culture medium reduce endometrial-cell survival in combined endometrial gland and stromal cell cultures from women with and without endometriosis. *Fertility and Sterility*, 76(4), 717-722.

- Gil, A. 2002. Polyunsaturated fatty acids and inflammatory diseases. *Biomedicine and Pharmacotherapy*, 56, 388-396.
- Gosslau, A. and Chen K.Y. 2004. Nutraceuticals, apoptosis, and disease prevention. *Nutrition*, 20, 95-102.
- Guil, J.L., Torija, M.E., Gimenez J.J. and Rodriguez, I. 1996. Identification of fatty acids in edible wild vegetables by gas chromatography. *Journal of Chromatography A*, 719, 229-235.
- Hamrouni, I., Salah, H.B. and Marzouk, B. 2001. Effects of water-deficit on lipids of safflower aerial parts. *Phytochemistry*, 58, 277-280.
- Harwood, J.L. 1996. Recent advances in the biosynthesis of plant fatty acids. *Biochimica et Biophysica Acta*, 1301, 7-56.
- Hogberg, A., Pickova, J., Andersson, K. and Lundstrom, K. 2003. Fatty acid composition and tocopherol content of muscle in pigs fed organic and conventional feed with different n6/n3 ratios, respectively. *Food Chemistry*, 80, 177-186.
- Horrobin, D.F. 1993. Omega-6 and omega-3 essential fatty acids in atherosclerosis. *Seminars in Thrombosis and Hemostasis*, 19, 129-137.
- Innis, S.M. 1996. Essential dietary lipids. In E.E. Ziegler and L.J. Filler (Eds), Present knowledge in nutrition (pp. 58-142). Washington, DC: ILSI Press.
- Kamchan, A., Puwastien, P., Sirichakwal, P.P. and Kongkachuichai, R. 2004. In vitro calcium bioavailability of vegetables, legumes and seeds. *Journal of Food Composition and Analysis*, 17, 311-320.
- Kabała, K., Janicka-Russak, M., Burzyński, M. and Kłobus, G. 2008. Comparison of heavy metal effect on the proton pumps of plasma membrane and tonoplast in cucumber root cells. *Journal of Plant Physiology*, 165, 278-288.
- Kinsella, J.E., Broughton, K.S. and Whelan, J.W. 1990. Dietary unsaturated fatty acids: interactions and possible needs in relation to eicosanoid synthesis. *The Journal of nutritional biochemistry*, 1(3), 123-141.
- Kosti, R.I., Panagiotakos, D.B., Tountas, Y., Mihas, C.C., Alevizos, A., Mariolis, T., Papathanassiou, M., Zampelas, A. and Mariolis, A. 2008. Parental Body Mass Index in association with The prevalence of overweight /obesity among adolescents in Greece; dietary and lifestyle habits in the context of the family environment: the Vyronas study. *Appetite*, 51(1):218-22.
- Laidlaw, M. and Holub, B.J. 2003. Effects of supplementation with fish oil-derived n 3 fatty acids and γ -linolenic acid on circulating plasma lipids and fatty acid profiles in women. *American Journal of Clinical Nutrition*, 77, 37-42.

- Li, D., Habito, R., Angelos, G., Sinclair, A.J. and Ball, M.J. 2003. Diet high in saturated fat decreases the ratio of thromboxane/prostacyclin in healthy men. *Biomedical and Environmental Science*, 16, 333-339.
- Li, X.C., Fan, X., Han, L.J. and Lou, Q.X. 2002. Fatty acids of some algae from the Bohai Sea. *Phytochemistry*, 59, 157-161.
- Liang, Y., Beardall, J. and Heraud, P. 2006. Effects of nitrogen source and UV radiation on the growth, chlorophyll fluorescence and fatty acid composition of *Phaeodactylum tricornutum* and *Chaetoceros muelleri* (Bacillariophyceae). *Journal of Photochemistry and Photobiology B: Biology*, 82(3), 161-172.
- Lin, Y.H., Chang, F.L., Tsao, C.Y., Leu, J.Y. 2007. Influence of growth phase and nutrient source on fatty acid composition of *Isochrysis galbana* CCMP 1324 in a batch photoreactor. *Biochemical Engineering Journal*, 37(15), 166-176.
- Liu, L.X., Howe, P., Zhou, Y.F., Xu, Z.Q., Hocart, C. and Zhang, R. 2000. Fatty acids and b-carotene in Australian purslane (*Portulaca oleracea*) varieties. *Journal of Chromatography A*, 893, 207-213.
- Maisuthisakul, P., Pongsawatmanit, R. and Gordon, M.H. 2007. Characterization of the phytochemicals and antioxidant properties of extracts from Teaw (*Cratoxylum formosum* Dyer). *Food Chemistry*, 100, 1620-1629.
- Manosroi, J., Dhumtanom, P. and Manosroi, A. 2006. Anti-proliferative activity of essential oil extracted from Thai medicinal plants on KB and P388 cell lines. *Cancer Letters*, 1, 114-120.
- Nouairi, I., Ammar, W.B., Youssef, N.B., D.D.B.M., Ghorbal, M.H. and Zarrouk, M. 2006. Comparative study of cadmium effects on membrane lipid composition of *Brassica juncea* and *Brassica napus* leaves. *Plant Science*, 3, 511-519.
- Okuda, N., Ueshima, H., Okayama, A., et al. 2005. Relation of long chain n-3 polyunsaturated fatty acid intake to serum high density lipoprotein cholesterol among Japanese men in Japan and Japanese-American men in Hawaii: the INTERLIPID study. *Atherosclerosis*, 178, 371-379.
- Oldroyd, J.C., Unwin, N.C., White, M., Mathers, J.C. and Alberti, K.G.M.M. 2006. Randomised controlled trial evaluating lifestyle interventions in people with impaired glucose tolerance. *Diabetes Research and Clinical Practice*, 72, 117-127.
- Pan, Z.Q., Rimando, A.M., Baerson, S.R., Fishbein, M. and Duke, S.O. 2007. Functional Characterization of Desaturases Involved in the Formation of the Terminal Double Bond of an Unusual 16:3 Δ 9,12,15 Fatty Acid Isolated from Sorghum bicolor Root Hairs. *Journal of Biological Chemistry*, 282(7), 4326-4335.

- Panagiotakos, D.B., Pitsavos, C., Chrysohoou, C., Skoumas, J. and Stefanadis, C. 2004. Status and management of blood lipids in Greek adults and their relation to socio-demographic, lifestyle and dietary factors: the ATTICA Study: Blood lipids distribution in Greece. *Atherosclerosis*, 173, 351-359.
- Pereira, C., Li, D. and Sinclair, A.J. 2001. The α -linolenic acid content of green vegetables commonly available in Australia. *International Journal Vitamin and Nutrition Research*, 71, 223-228.
- Simopoulos, A.P. 2002. The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomed Pharmacother*, 56, 365-379.
- Sinclair, H.M. 1990. Essential fatty acids: a historical perspective. *Biochemical Society Transactions*, 18, 756-761.
- Sirtori, C.R., Crepaldi, G., Manzato, E., Mancini, M., Rivellesse, A., Paoletti, R., Pazzucconi, F., Pamparana, F. and Stragliotto, E. 1998. One-year treatment with ethyl esters of n-3 fatty acids in patients with hypertriglyceridemia and glucose intolerance reduced triglyceridemia, total cholesterol and increased HDL-C without glycemic alterations. *Atherosclerosis*, 137, 419-427.
- Sprecher, H. 2000. Metabolism of highly unsaturated n-3 and n-6 fatty acids. *Biochimica et Biophysica Acta*, 1486, 219-231.
- Sugiyama, T., Xie, D., Graham-Maar, R.C., Inoue, K., Kobayashi, Y. and Stettler, N. 2007. Dietary and Lifestyle Factors Associated with Blood Pressure among U.S. Adolescents. *Journal of Adolescent Health*, 40(2), 166-172.
- Sukenik, A. and Livne, A. 1991. Variations in lipid and fatty acid content in relation to acetyl coA carboxylase in the marine prymnesiopyte *isochrysis galbana*. *Plant and Cell Physiology*, 32, 371-378.
- Tsipas, G. and Morphak, P. 2003. Beneficial effects of a diet rich in a mixture of n-3 essential fatty acids and of their metabolites on cyclosporine-nephrotoxicity. *Journal of Nutrition Biochemistry*, 14, 480-486.
- Wang, X. and Tian, W.X. 2001. Green tea epigallocatechin gallate: A natural inhibitor of fatty-acid synthase. *Biochemical and Biophysical Research Communications*, 288, 1200-1206.
- WHO/FAO. 2002. Diet, nutrition and the prevention of chronic diseases. Geneva, Switzerland. *WHO technical report series*, 916, 54-132.
- WHO. 2003. WHO global strategy on diet, physical activity and health: South-East Asia regional consultation meeting report. India: Nww Delhi.

- Williams, J.P., Williams, K. and Khan, M.U. 1992. Low temperature-induced fatty acid desaturation in *Brassica napus*: thermal lability of the process. *Biochimica et Biophysica Acta*, 1125(1), 62-67.
- Yang, L.F., Siriamornpun, S. and Li, D. 2006. Polyunsaturated fatty acid content of edible insects in Thailand. *Journal of Food Lipids*, 13, 277-285.
- Zhang, S., Sakuradani, E., Ito, K. and Shimizu S. 2007. Identification of a novel bifunctional D12/D15 fatty acid desaturase from a basidiomycete, *Coprinus cinereus* TD#822-2. *FEBS Letters*, 581, 315-319.