

## เอกสารอ้างอิง

- [1] Kudo, A., Omori, K. and Kato, H., A novel aqueous process for preparation of crystal form controlled and highly crystalline  $\text{BiVO}_4$  Powder from layered vanadates at room temperature and photophysical properties. *J. Chem. Soc.*, 1999, **121**, 11459-11467.
- [2] Tokunaga, S., Kato, H. and Kudo, A., Selective preparation of monoclinic and tetragonal  $\text{BiVO}_4$  with scheelite structure and their photocatalytic properties. *Chem. Mater.*, 2001, **13**, 4624-4628.
- [3] Bhattacharya, A. K., Mallick, K. and Hartridge, A., Phase transition in  $\text{BiVO}_4$ . *Mater. Lett.*, 1997, **30**, 7-13.
- [4] Zhou, Y., Vuille, K., Heel, A., Probst, B., Kontic, R. and Patzke, G. R., An inorganic hydrothermal route to photocatalytically active bismuth vanadate. *Appl. Catal. A-Gen.*, 2010, **375**, 140-148.
- [5] Zhang, Z., Wang, W., Shang, M. and Yin, W., Photocatalytic degradation of rhodamine B and phenol by solution combustion synthesized  $\text{BiVO}_4$  photocatalyst. *Catal. Commun.*, 2010, **11**, 982-986.
- [6] Bismuth vanadate. <http://spie.org/x35165.xml?pf=true&ArticleID=x35165>. (accessed Dec 12, 2011).
- [7]  $\text{BiVO}_4$ . <http://jsmchem.co.kr/sds/MSDS-YELLOW.doc>. (accessed Dec 12, 2011).
- [8] Phenol. <http://en.wikipedia.org/wiki/Phenol>. (accessed Dec 12, 2011).
- [9] 2-chlorophenol. <http://en.wikipedia.org/wiki/2-Chlorophenol>. (accessed Dec 12, 2011).
- [10] 2,4-dichlorophenol. <http://msds.pcd.go.th/searchName.asp?vID=275>. (accessed Dec 12, 2011).
- [11] Gotić, M., Music, S., Ivanda, M., Soufek, S. and Popovic, S., Synthesis and characterization of bismuth (III) vanadate. *J. Mol. Struct.*, 2005, **744**, 535-540.
- [12] Wood, P. and Glasser, F. P., Preparation and properties of pigmentary grade  $\text{BiVO}_4$  precipitated from aqueous solution. *Ceram. Int.*, 2004, **30**, 875-882.
- [13] Liu, J. B., Wang, H., Wang, H. and Yan, H., Hydrothermal preparation of  $\text{BiVO}_4$  powder. *Mat. Sci. Eng. B-Solid.*, 2003, **104**, 36-39.

- [14] Byrappa, K. and Yoshimura, M., *Handbook of hydrothermal technology: a technology for crystal growth and materials processing*, Noyes Publication, New York, 2001.
- [15] Riman, R. A., Pugh, R. J. and Bergstrom, L., The chemical synthesis of ceramic powder; in surface and colloidal chemistry in advanced ceramics processing. *Marc. Dek. Inc.*, 1995, **22**, 2310-2313.
- [16] Pecchi, G., Reyes, P., Sanhueza, P. and Villaseñor, J., Photocatalytic degradation of pentachlorophenol on TiO<sub>2</sub> sol-gel catalysts, *Chemosphere*, 2001, **43**, 141-146.
- [17] Sol-gel process. <https://www.llnl.gov/str/May05/Satcher.html>. (accessed Jan 27, 2012).
- [18] วราภรณ์ ชังฉิม, การสังเคราะห์และหาลักษณะเฉพาะของผงนาโนซิงค์ออกไซด์โดยวิธีโซลเจล, ปัญหาพิเศษวิทยาศาสตร์บัณฑิต, คณะวิทยาศาสตร์, มหาวิทยาลัยแม่โจ้, 2549.
- [19] กระบวนการโฟโตแคตาไลติกส์. <http://www.eng.ru.ac.th/envi/aj/EVE410/photocatalytic.pdf>. (accessed Jan 27, 2012).
- [20] Photocatalytic. <http://sichon.wu.ac.th/file/envi-shh-20090110-112240-pwrqR.pdf>. (accessed Jan 27, 2012).
- [21] Hayat, K., Gondal, M. A., Khaled, M. M. and Ahmed, S., Effect of operational key parameters on photocatalytic degradation of phenol using nano nickel oxide synthesized by sol-gel method, *J. Mol. Catal. A-Chem.*, 2011, **336**, 64-71.
- [22] Tücks, A. and Beck, H. P., The photochromic effect of bismuth vanadate pigments. Part I: synthesis, characterization and lightfastness of pigment coatings. *J. Solid State Chem.*, 2005, **178**, 1145-1156.
- [23] Photocatalytic. <http://www.titanpe.com/library/kb2501.htm>. (accessed Dec 12, 2011).
- [24] Zhang, X., Ai, Z., Jia, F., Zhang, L., Fan, X. and Zou, Z., Selective synthesis and visible-light photocatalytic activities of BiVO<sub>4</sub> with different crystalline phases. *Mater Chem. Phys.*, 2007, **103**, 162-167.
- [25] Photocatalytic. [http://puregreencoatings.com/pgc\\_science.aspx](http://puregreencoatings.com/pgc_science.aspx). (accessed Dec 12, 2011).
- [26] D'Oliveira, J. C., Al-Sayyed, G. and Phichat, P., Photodegradation of 2- and 3-chlorophenol in TiO<sub>2</sub> aqueous suspensions. *Environ. Technol.*, 1990, **24**, 1415-1417.

- [27] Ilisz, I., Dombi, A., Mogyorosi, K., Farkas, A. and Dekany, I., Removal of 2-chlorophenol from water by adsorption combined with  $\text{TiO}_2$  photocatalysis. *Appl. Catal. B-Environ.*, 2002, **39**, 247-256.
- [28] Suryaman, D. and Hasegawa, K., Biological and photocatalytic treatment integrated with separation and reuse of titanium dioxide on the removal of chlorophenols in tap water. *J. Hazard. Mater.*, 2010, **183**, 490-496.
- [29] Zhang, X., Zhang, Y., Quan, X. and Chen, S., Preparation of Ag doped  $\text{BiVO}_4$  film and its enhanced photoelectrocatalytic (PEC) ability of phenol degradation under visible light. *J. Hazard. Mater.*, 2009, **167**, 911-914.
- [30] Pookmanee, P., Pingmuang, K., Phanichphant, S. and Kangwansupamonkon, W., Characterization of bismuth vanadate powder synthesized by a hydrothermal method. *J. Micro. Soc. Thai.*, 2009, **32(1)**, 95-98.