

เอกสารอ้างอิง

- ชลอ ลี้มสุวรรณ. 2536. แนวทางการเลี้ยงกุ้งหน้าฝน. วารสารอะควาฟาร์มมิ่ง 5: 30-36.
- เทพรัตน์ อึ้งเศรษฐพันธ์ สุเทพ ปันธิวงศ์ สมบูรณ์ ใจปิ่นตา ประจวบ ฉายนุ สุดปราณี มณีศรี และรุ่งกานต์ อำไพพงศ์. 2545. แนวทางการจัดการปัญหาการผลิตและการตลาดปลาน้ำจืดจังหวัดเชียงใหม่. รายงานวิจัยสำนักงานกองทุนสนับสนุนการวิจัย PDG45N0008. 85 หน้า.
- ศูนย์สารสนเทศ. 2547. สถิติการประมงแห่งประเทศไทย พ.ศ. 2545. กรุงเทพฯ: กรมประมงกระทรวงเกษตรและสหกรณ์. 91 น.
- Casey, C. G., W.L. Steven, and V.Z. Paul. 2004. Instrumental versus sensory detection of off-flavors in farm-raised channel catfish. *Aquaculture* (236): 309-319.
- Farmer, L.J., J.M. McConnell, T.D.J. Hagan and D.B. Harper. 1995. Flavor and off-flavor in wild and farmed Atlantic salmon from locations around Northern Ireland. *Water Science and Technology*. 31(11): 259-264.
- Form, J. and V. Horlyck. 1984. Site of uptake geosmin a cause of earthy-flavor in rainbow trout (*Salmo gairdneri*). *Can. J. Fish. Aquat. Sci.* 41: 1224-1226.
- Izaurire, G., C.J. Hwang, S.W. Krasner and J. Micheal. 1982. Geosmin and 2-methylisoborneol from cyanobacteria in three water supply system. *App. Envi. Micro.* 43(3): 708-714.
- Johnsen, P.B. and S.W. Lloyd. 1992. Influence of fat content on uptake and depuration of the off-flavor 2-methylisoborneol by channel catfish (*Ictalurus punctatus*). *Can. J. Fish. Aquat. Sci.* 49 : 2406-2411.
- Johnson, P.B. and C.P. Dionigi. 1994. Physiology approaches to the management of off-flavor in farm-raised channel catfish (*Ictalurus punctatus*). pp. 141-161. In D. Tave and C.E. Tucker (eds.). *Recent Development in Catfish Aquaculture*. New York : The Haworth Press, Inc.
- Johnsen, P.B., S.W. Lloyd, B.T. Vingad and P.C. Dionigi. 1996. Effect of temperature on uptake and depuration of 2-methylisoborneol in channel catfish (*Ictalurus punctatus*). *J. World Aqua. Soc.* 27(1): 15-20.
- Klapper, H. 1991. *Control of Eutrophication in Inland Waters*. New York: Ellis Horwood.
- Lovell, R.T. and D. Broce. 1985. Cause of musty flavor in pond culture penaeid shrimp. *Aquaculture penaeid shrimp*. *Aquaculture* 50: 169-174.

- Martin, J.F., C.P. McCoy, W. Greenleaf and L.W. Bennett. 1987. Analysis of 2-methylisoborneol in water, mud and channel catfish (*Ictalurus punctatus*) from commercial culture ponds in Mississippi. *Can. J. Fish. Aquat. Sci.* 45: 909-912.
- Martin, J.T., L.W. Bennett and W.H. Graham. 1988. Off-flavor in the channel catfish (*Ictalurus punctatus*) due to 2-methylisoborneol and its Dehydration Products. *Water Sci. Technol.* 29 (8/9): 59-65.
- Martin, J.F., M.S. Plakas, H.J. Holley, J.V. Kitzman and A.M. Guaino. 1990. Pharmacokinetics and tissue disposition of the off-flavor compound 2-methylisoborneol in the channel catfish (*Ictalurus punctatus*). *Can. J. Fish. Aqua. Sci.* 47 : 544-547.
- Matsuyasu, N., O. Takahoro, K. Yoshiyuki, I. Noriyuki, I. Taichi, A. Akiro, S. Toshiaki, H. Euichi and S. Michio. 1996. Inhibitory effects of odor substances, geosmin and 2-methylisoborneol, on early development of sea urchins. Elsevier Science Ltd. PII: S0043-1354(96)00104-2.
- Milie, D.F., M.C. Baker, C.S. Tucker, B.T. Vinyard and C.P. Dionigi. 1992. High-resolution Airborne remote sensing of bloom-forming phytoplankton. *J. of Phytocology.* 28: 28-290.
- Persson, P.E. 1982. Muddy odor: a problem associated with extreme eutrophication. *Hydrobiologia* 89: 161p.
- Piumsombun S. 2001. Production, accessibility and consumption patterns of aquaculture products in Thailand. FAO Fisheries Circular No. 973.
- Rungreungwudhikrai, E. 1995. Characterization and classification of off-flavor of Nile tilapia. M.S. Thesis no. AE-95-24. Bangkok: Asian Institute of technology.
- Sivonen, K. 1982. Factor influencing odor production by actinomycetes. *Hydrobiologia.* 86: 165-170.
- Tabachek, J.L. and M. Yurkowski. 1976. Isolation and identification of blue-green algae producing muddy odor metabolites and 2-methylisoborneol in saline lake in Monitoba. *J. Fish Res. Board Can.* 33: 25-35.
- Tanchotikul, U. 1990. Studies on important volatile flavor compounds in Louisiana rangia clam (*Rangia cuneata*). Doctoral dissertation. Louisiana state university. 96 p.
- Tanchotikul, U. and T.C.Y. Hsieh. 1990. Methodology for quantification of geosmin and Levelin raggia clam (*Rangia cuneata*). *J. Food Sci.* 55(5): 235-312.

- Van Der Ploeg, M. 1989. Seasonal trends in flavor quality of channel catfish (*Ictalurus punctatus*) from commercial pond in Mississippi. *J. of Applied Aquaculture*. 2(3): 22-31.
- Van Der Ploeg, M. and C.E. Boyd. 1991. Geosmin production in cyanobacteria (blue green algae) in fish pond at Auburn, Alabama: *J. of the World Aquaculture Society* 22(4): 207-216.
- Yamada, N., N. Marakami, N. Kawamura and J. Sakakibara. 1994. Mechanism of an early lysis by fatty acid from *Axenic Phormidium tenue* (Musty odor-producing cyanobacterium) and its growth prolongation by bacteria. *Biol. Pharm. Bull.* 17(9): 1277-1281.
- Yamprayoom, J. and A. Noomhorm. 2000. Geosmin and Off-flavor in Nile Tilapia (*Oreochromis niloticus*). *J. of Aquatic product technology* 9(2): 29-41.
- Yurkowski, M. and J.L. TabachekL. 1974. Identification analysis and removal of geosmin from Muddy flavored trout. *J. Fish. Res Board. Can.* 31: 1851-1858.
- Yurkowski, M. and J.L. Tabachek. 1980. Geosmin and 2-methylisoborneol implicated as a cause of muddy odor and flavor in commercial fish from Cedar Lake. Manitoba. *Can. J. Fish. Aquat. Sci.* 37: 1449-1450.
- Zilberg B. 1966. Gastroenteritis in Salisbury European children – a five-year study. *Cent. Afr. J. Med.*, 12(9):164-168.