TABLE OF CONTENTS

	Page
TABLE OF CONTENTS	i
LIST OF TABLES	ii
LIST OF FIGURES	iii
LIST OF ABBREVIATIONS	vii
INTRODUCTION	1
LITERATURE REVIEW	3
MATERIALS AND METHODS	27
RESULTS	39
DISCUSSION	67
CONCLUSIONS	73
LITERATURE CITED	74
APPENDIX	89

LIST OF TABLES

Table		Page
1	Classification of animal lectins by Gudmundur Johann	
	Arason	16
2	Summary of lectin categories by Roger B. Dodd and Kurt	
	Drickamer	17
3	Properties of purified lectins from Penaeid shrimp	20
4	Hemagglutinating activity (HA) of serum from	
	Penaeus merguiensis against different types of mammalian	
	and animal erythrocytes	39
5	Effects of temperature on hemagglutination	40
6	Effects of pH on hemagglutination	40
7	Effects of divalent cation on hemagglutination	40
8	The sugar specificity of lectin from serum of Penaeus	
	mergueinsis	42
9	Purification of Penaeus merguiensis lectin	45
10	Partial amino acid sequences of the purified lectin subunits	
	from hemolymph of Penaeus merguiensis	48
11	Antibacterial activity in serum, hemolymph affinity and gel	
	filtration purified Penaeus merguiensis lectin incubated with	
	Vibrio spp. for 1 h. Percentage of inhibition was determined	
	from the appropriate serial dilution as colony forming unit	
	(CFU)	54
12	Protein expression in hemolymph of Penaeus merguiensis	
	after challenging with Vibrio harveyi	60
13	Hemagglutinating activity (HA) of serum hemolymph	
	haemocyte lysate supernatant from P. merguiensis against	
	different types of mammalian and animal erythrocytes	64

LIST OF FIGURES

Figure		Page
1	Geographical distribution for Penaeus merguiensis	3
2	Geographical distribution for Penaeus merguiensis in the	
	Upper Gulf of Thailand	4
3	Global capture production for Penaeus merguiensis	5
4	Global aquaculture production for Penaeus merguiensis	5
5	Thailand marine shrimp culture production during	
	1985-2004	6
6	General morphological features of Penaeid shrimp	
	(Penaeus latisulcatus)	7
7	Life cycle of shrimp showing distribution with depth, a) shrimp	
	eggs b) nauplius c) protozoea d) mysis e) postmysis f) juvenile	
	and g) mature adult shrimp	9
8	Affinity chromatography of the P. merguiensis lectin on mucin	
	CNBr-activated Sepharose 4-B	44
9	Gel filtration of the <i>P. merguiensis</i> lectin on Sephacryl S-200	44
10	SDS-PAGE (12.5%) of the <i>P. merguiensis</i> lectin eluted from	45
	PSM-Sepharose 4B column and Sephacryl S-200 column	
11	2-D gel performed with 200 µg of affinity purified lectin shows	
	the seperation of subunits by using Immobiline DryStrip pH	
	3-10 (13 cm)	47
12	Mass spectrum of spot no.1 by LC-MS/MS; precursor mass	
	974	49
13	Mass spectrum of spot no. 1 by LC-MS/MS; precursor mass	
	1047	49
14	Mass spectrum of spot no.1 by LC-MS/MS; precursor mass	
	1062	50

LIST OF FIGURES (Continued)

Figure		Page
15	Mass spectrum of spot no.1 by LC-MS/MS; precursor mass	
	1393	50
16	Mass spectrum of spot no. 2 by LC-MS/MS; precursor mass	
	671.86	51
17	Mass spectrum of spot no. 3 by LC-MS/MS; precursor mass	
	820.62	51
18	Mass spectrum of spot no. 4 by LC-MS/MS; precursor mass	
	591.10	52
19	Mass spectrum of spot no. 5 by LC-MS/MS; precursor mass	
	725.74	52
20	Mass spectrum of spot no. 6 by LC-MS/MS; precursor mass	
	620.88	53
21	Amino acid sequence which partial amino acid HWFSLFNPR	
	blase to NCBI (Mass spectrum of Spot 6 by LC-MS/MS;	
	precursor mass 602.88)	53
22	Bacterial clearance activity in banana prawn after challenge with	
	Vibrio harveyi	55
23	Protein profile of hemolymph from the P. merguiensis in 10%	
	SDS-PAGE (13 cm,) after challenged with Vibrio harveyi	57
24	2D maps of colloidal Coomassie brilliant blue G-250 stained	
	proteins of the affinity purified lectin after post injection with	
	V. harveyi at 1 min (A) and 120 min (B). The protein was	
	separated by 2-Dimensional Gel Electrophoresis. Isoelectric	
	focusing (IEF) was perfomed in the Immobiline TM DryStrip	
	pH3-10, 13 cm, IPG strips in a disposable cassette	58

LIST OF FIGURES (Continued)

Figure		Page
25	2D maps of colloidal Coomassie brilliant blue G-250 stained	
	proteins of the affinity purified lectin after post injection with	
	V. harveyi at 1 min (A) 15 min (B) and 120 min (C)	59
26	Light micrographs of shrimp hemocytes, Penaeus merguiensis.	
	(A) hyaline hemocytes; (B) a small granule hemocyte (SGH);	
	(C) large granule hemocytes (LGH)	62
27	Total and differential counts of hemocytes in Penaeus	
	merguiensis hemolymph (values are averge±SD, n=30) Total	
	hemocyte count (THC), small granule hemocyte (SGH), large	
	granule hemocyte (LGH)	62
28	12.5% SDS-PAGE of hemolymp (lane 2) serum (lane 3) and	
	hemocyte lysate supernatant (lane 4) proteins in reducing	
	condition. The protein in samples were treated with	
	dithiothreitol (DTT). Protein bands were visualized with	
	Silver staining	63
29	Antibacterial activity of serum, affinity purified lectin (AF)	
	and gel filtration purified lectin (GF) (~ 1.28 µg proteins)	
	and HLS (~ 2 µg proteins) from hemolymph and hemocyte of	
	Penaeus merguiensis	66
Appendi	x Figure	
1	2-D maps of colloidal Coomassie brilliant blue G-250 stained	
	proteins of the affinity purified lectin after post injection with	
	V. harveyi at 1 min (A) Green and red label spots are	
	responsed to V. harveyi infection	90

LIST OF FIGURES (Continued)

Appendix Figure		Page
2	2-D maps of colloidal Coomassie brilliant blue G-250 stained proteins of the affinity purified lectin after post injection with	
	V. harveyi at 15 min (B). Green and red label spots are	
	responsed to V. harveyi infection	91
3	2-D maps of colloidal Coomassie brilliant blue G-250 stained	
	proteins of the affinity purified lectin after post injection with	
	V. harveyi at 120 min (C). Green and red label spots are	
	responsed to V. harveyi infection	92

LIST OF ABBREVIATIONS

cm = centimeter

EDTA = ethylene diamine tetraacetic acid

 $\mu l = microliter$

M = Molar

mg = milligram

 $\mu g = microgram$

min = minute

ml = milliliter

mM = millimolar

NCBI = National Center for Biotechnology Information

rpm = rotations per minute

sec = second

 $egin{array}{lll} U & = & unit \\ V & = & volt \end{array}$

.

v = volume